



Management Practices, Perceived Constraints and Estimation of Biochemical Parameters in Indigenous Goats of Gwalior City of Central India

Anjali Gautam ^a, Kumar Govil ^{b*}, Jitendra Yadav ^c,
Amit Oad ^a, Anjani Kumar Mishra ^d,
Brijesh Kumar Ojha ^b and Abhilash Singh ^b

^a Department of Animal Husbandry and Dairying, Government of Madhya Pradesh, Rewa, M.P. 486001, India.

^b Department of Veterinary and Animal Husbandry Extension Education, College of Veterinary Science and Animal Husbandry, Rewa, M.P. 486001, India.

^c Department of Animal Nutrition, College of Veterinary Science and Animal Husbandry, Rewa, M.P. 486001, India.

^d Department of Livestock Production and Management, College of Veterinary Science and Animal Husbandry, Rewa, M.P. 486001, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJAEES/2023/v41i102199

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/106956>

Original Research Article

Received: 10/08/2023
Accepted: 28/09/2023
Published: 02/10/2023

ABSTRACT

Aim: A study was conducted in order to analyze goat rearing practices prevailing in the Gwalior City of Central India and to identify the constraints in goat farming.

*Corresponding author: E-mail: kumargovil22@gmail.com;

Study Design: Data of ninety six indigenous goats of 2 to 5 age group from thirty respondents from the district was collected.

Place and Duration of Study: The study was carried out in Gwalior City of Central India during July 2023.

Methodology: The data pertaining to goat rearing practices being followed and constraints being faced were collected with the help of pre-structured interview schedule after ensuring its reliability and validity. To evaluate the nutritional status of the animals the blood sample of twenty four randomly selected female goats of 2 to 5 years of age was taken from Jugular vein for the normal biochemical profile.

Results: It was observed that total protein and serum calcium level was significantly lower than normal value, while. Alkaline phosphates, SGOT and SGPT and blood urea levels are higher than the normal values.

Conclusion: It is concluded that goats reared on extensive system of rearing have poor reproductive performance (higher age at first kidding and kidding interval and low twinning rate) due to lack of scientific management and feeding practices by goat farmers of Gwalior City of Central India

Keywords: Goats; feeding; management; blood biochemistry; reproductive performance; constraints.

1. INTRODUCTION

With the increase in demand of goat products, the population of goats increased from 135.17 in 2012 to 148.88 million goat in 2019, showed a 10% growth rate over previous livestock census. Madhya Pradesh has fifth rank in goat population in India, population of goats increased from 8.3 million in 2012 to 11.6 millions in 2019, showed tremendous growth of 38% [1]. "Goat is a multipurpose animal and mostly reared by landless, poor, marginal and nomadic farmers under the most traditional managerial systems using low or no cost inputs" [2]. "Goats (*Capra hircus*) plays a significant role in the rural economy of India and acts as an insurance against crop failure to meet the immediate demand of finance, in addition it also provides an alternative source of livelihood to the farmers throughout the year. Goats in India are also known as poor man's cow because goat rearing is a great economic support to a large section of population in rural areas. India is a rich repository of goat genetic resources in the form of 28 well defined breeds" [3].

"At farmer level to obtain the maximum production, animals should be scientifically managed for feeding and breeding. Productive and reproductive traits are the most important traits in all animal production systems and reproductive efficiency is the most important factor affecting production rate and profit in livestock sector" [4]. "Thus, efforts should be made to improve these traits for efficient goat production, which are further influenced by genetic, environmental, and managerial

factors. Any deficiency of nutrients affecting the production and reproduction of the animal are well appeared blood biochemical profile. Due to the limited information about the local goat breeds about their blood chemistry, it is essential to conduct such a research especially because local people prefer them more than the commercial breeds. Thus, biochemical profiles are important preliminary step to be determined because they provide valuable information about the breed, sex and health status of the animal" [5]. Thus, keeping these views in mind the research was to observe the managerial practices followed at field level by farmers of Gwalior city of central India and investigation of blood chemistry of indigenous goats as a preliminary study, to assess the nutritional status of the animals.

2. MATERIALS AND METHODS

A study was conducted in order to analyze goat rearing practices prevailing in the Gwalior City of Central India and to identify the constraints in goat farming through collection of data of ninety six indigenous goats of 2 to 5 age group from thirty respondents from the district was collected. The data pertaining to goat rearing practices being followed and constraints being faced were collected with the help of pre-structured interview schedule after ensuring its reliability and validity. Blood samples were collected randomly from 24 from goat from selected goat farmers. The animals were restrained and blood samples were collected directly from jugular vein of the Goats under aseptic condition by using 15 gauge in vials uncoated and coated with EDTA vials for

blood biochemistry by blood analyser for estimation of blood biochemical constituents. Blood biochemical constituent's viz. blood Glucose, albumin, globulin, total protein, blood urea, SGPT, SGOT, alkaline phosphatase and calcium.

Each selected respondent was interviewed personally according to the prepared questionnaire for the collection of desired information about management, feeding and breeding information followed by the goat farmers of the area was collected and analyzed. The data of ninety six adult female goats (same age group of 2 to 5 years) about method of feeding, concentrate feeding, mineral mixture supplementation, deworming, age at first kidding (days), kidding Interval (days), twinning rate was collected from goat farmers.

3. RESULTS AND DISCUSSION

3.1 Management Practices and Perceived Constraints

The information regarding health care practices and nutritional status of the goats (n= 96) opted by the goat farmers of Gwalior city of Central India are depicted in Table 1. On the basis of data obtained it was reported that all the goats are reared on extensive system of rearing in which goats are allowed to graze in forest for average 6 to 8 hours a day only and no concentrate is fed to the animal, thus, all goats are solely depends on grazing to fulfill their nutritional demands. It might be due to intensive system of rearing requires initial high cost involvement and lack of knowledge about scientific goat rearing. On the other hand, maximum farmers (90%) of Burdwan district of West Bengal rearing goats in semi intensive system of rearing [6]. While data obtained in present study showed that farmers of the selected area does not fed concentrate to the goats i.e, 0% animal receiving concentrate and 6.25% goats received the mineral mixture, rest 93.75% goats did not receiving any mineral mixture.

As the goats are reared exclusively on extensive system of rearing, chances of parasitic infection are higher but only 18.75% animals were received the drug for control of endoparasites while remaining 81.25% animals were not received any drug for endoparasite control. Similarly, out of 39 goats examined, 23 found positive for helminth infestation in Jaipur District

of Rajasthan [7]. "Gastrointestinal parasitism is one of the major threats to small ruminant industry causing production losses and even mortality in severe cases" [8]. "Parasitic infestation is responsible to affect the growth, body weight, yield and reproductive performance of animal leading to economic loss of the farmer" [9]. "In sub-clinical cases also parasites continuously sucks host blood resulting in anemia, hypoproteinemia and lower blood glucose" [10].

In present study, all goats (100%) are not receiving any supplement ration or special feeding as per demand in pregnancy to meet the demand of growing foetus. All of the goat farmers kept pregnant and non pregnant animals together it might be due to either lack of knowledge about the isolation of pregnant does from others in the herd or may be due to less availability of space.

The findings presented in Table 1 revealed that, majority of goats showed history of foot and mouth disease (FMD), mastitis, metritis, prolapsed, dystocia, retained fetal membranes and disease incidence are 77.0%, 42.7 %, 31.25 %, 25.0 %, 21.8 % and 14.5 %, respectively, were reported.

On the basis of data obtained it was reported that overall means for age at first kidding, kidding Interval and twinning rate are 654 ± 94.4 days, 391.2 ± 31.5 days, 1.1 ± 0.1 , respectively. This variation might be attributed to different plan of nutrition as well as reproductive health.

Similarly, majority of respondents (64.67%) received 2 kids per birth, maximum number of goat farmers (64.67%) pointed out that kidding interval was 8-10 months, whereas 23.33 per cent raisers noticed kidding interval above 10 months in black bengal goat rearing in Burdwan district of West Bengal,. In addition 57.33% farmers in the district area vaccinated their goats against major diseases like Goat pox, PPR and FMD [6].

On the basis of data obtained in present study, it is found that indigenous goats showed higher age at first kidding, kidding interval and low twinning rate. It might be due to absence of concentrate and mineral mixture in feeding of goats along with no timely deworming of the animal. Similarly, production performance of Jakhara goats in its home tract in District Alwar of Rajasthan and reported average age at first kidding, kidding interval and twinning rate were

561.24 ± 9.83 days, 287.78 ± 9.89 days, and 1.54 ± 0.16 days in semi-intensive, and 632.3 ± 7.83 days, 332.86 ± 9.34 days and 1.32 ± 0.16 in extensive system of management respectively [11].

Patel and Pandey (2013) [12] also found age at first kidding (days), kidding Interval (days) 364.40± 4.32 and 716.52 ± 19.01 respectively in Mehsana goats. Similarly, the overall least squares means of age at first conception, age at first kidding, kidding interval, service period, gestation period and dry period were 513.20±20.94, 657.20±20.48, 292.03±4.14, 142.17±4.18, 149.95±0.07 and 144.00±4.09 days, respectively in Sirohi goats of Udaipur city of India [13]. Arun et al. [14], however, reported age at first kidding of 698.41 ± 1.49 days in Kutchi goats. The average at first kidding obtained in the present finding was in agreement to the findings of Acharya [15].

On the basis of information obtained from present study it was revealed that, most important constraints in the goat rearing practices-

1. Lack of pure breed buck (due to higher cost of castrated buck in market),
2. High incidence of diseases(due to lack of proper vaccination),

3. Improper feeding and management practices involves lack of mineral mixture feeding and timely deworming and vaccination),
4. Problem of lack of capital to start goat farm and
5. Lack of knowledge about scientific goat rearing.
6. Improper feeding management involves lack of mineral mixture and concentrate supplementation as per the requirement of different physiological stages along with timely deworming. Similar observations were also reported by Singh et al. [16], Guljar and Pathodiya [17] in Rajasthan and by Meganathan et al. [18] in Tamil Nadu.

3.2 Estimation of Blood Biochemical Parameters of the Indigenous Goats

The blood biochemical values of indigenous goats reared under extensive system of rearing are presented in Table 2. The overall mean of values showed the lower values of serum calcium and total protein levels, while, higher overall mean values of blood urea nitrogen (BUN), alkaline phosphatase, SGOT, SGPT levels were found.

Table 1. Health and nutritional status parameters of Indigenous Goat (*Capra hircus*)

S.No.	Parameters	Mean	
1	Type of rearing	Intensive	0
		Semi intensive	0
		Extensive	96 (100%)
2	Concentrate feeding	Yes	0
		No	96 (100%)
3	Special feeding of pregnant goat	Yes	0
		No	96 (100%)
4	Segregation of pregnant doe from others	Yes	12 (12.5%)
		No	84 (87.5%)
5	Feeding of mineral mixture	Yes	6 (6.25%)
		No	90 (93.75%)
6	Drugs used to control endoparasites	Yes	18 (18.75%)
		No	78 (81.25%)
7	The disease incidence in buffaloes	FMD	74 (77%)
		Mastitis	41 (42.7%)
		Metritis	30 (31.25%)
		Prolapse	24 (25.0%)
		Dystocia	21 (21.8%)
		ROP	14 (14.5%)
6	Age at first kidding (days)	654 ± 94.4	
7	Kidding Interval (days)	391± 31.5	
8	Twinning rate	1.1±0.1	

Table 2. Values of blood parameters of Indigenous Goat (*Capra hircus*)

S. No.	Parameters	Mean value	Normal value*
1	Glucose (mg/dl)	51.0± 9.0	50-75
3	Serum Ca (mg/dl)	8.1± 0.97	8.9-11.7
4	Total protein (g/dl)	5.0±0.62	6.4-7.0
5	Albumin (g/dl)	3.0± 0.26	2.7-3.9
6	Globulin (g/dl)	2.0± 0.68	2.7-4.1
7	A:G ratio	1.8± 0.77	-
8	T Bilirubin (mg/dl)	0.9± 0.09	-
9	Bilirubin direct (mg/dl)	0.3±0.08	-
10	Bilirubin indirect (mg/dl)	0.6±0.14	-
11	Alkaline phosphatase (IU/L)	121.9±7.81	93-387
12	SGOT (U/L)	67.3±12.00	167-513
13	SGPT (U/L)	82.6±12.81	6-19
14	Blood urea (mg/dl)	31.1± 4.78	10-20

* Whitbread, T. J. (2017) *Clinical Biochemistry MSD Veterinary manual* [19]

Blood biochemical profile can be used to evaluate the health, nutritional and physiological status of the goats. The glucose level is found within the normal range (51.0± 9.0 mg/dl). Blood glucose is one of the key nutrients affecting production and reproduction in farm animals and a minimum level of 40-60 mg/dl is required to maintain the physiological processes of the body [20]. Low blood glucose may cause infertility. The blood glucose can be used as indices of nutritional status during pregnancy in goats [21].

Proteins are the building blocks of all cells and body tissues. They act as transport substances of hormones, vitamins, minerals, lipids and other materials. Thus, protein act as working horses of the cell [22]. Baumgartner and Pernthaner (1994) [23] observed no effect of physiological stage on total protein. The total protein (g/L), albumin (g/L), urea (mg/dl), ALT (U/L), AST (U/L) in black bengal goats are 69.9 ± 1.03, 43.01 ± 0.75, 23.90 ± 0.89, 26.15 ± 1.72, 59.8 ± 2, respectively, while, values for Jamnapari goats are 78.53 ± 2.05, 47.36 ± 1.35, 35.85 ± 1.47, 17.33 ± 0.69, 47.16 ± 1.05, respectively [24].

Urea is made in the liver and passed out from the body in the urine. A blood urea nitrogen test is done to assess the function of kidneys. If kidneys are not able to remove urea from the blood normally, then BUN level rises. Alanine Aminotransferase (ALT or SGPT), is an enzyme found in many tissues, but the highest levels are found in liver and kidney tissues. Serum ALT has been recognized as a marker of hepatocellular injury since the 1950s [25]. Aspartate aminotransferase (AST or SGOT) is an enzyme involved in amino acid metabolism. Increased

serum AST activity is observed with both reversible and irreversible injury to hepatocytes and can be seen following hepatocellular injury and cholestasis, similar to serum ALT activity in dogs, swine, cats and goats.

The diagnostic sensitivity of serum AST activity in animals has been reported as 72% for hepatic necrosis and 100% for hepatic lipidosis [26]. In West African dwarf goat mean value of AST reported as 20.9 ± 1.2 U/L [27], while AST in wild goats reported as 235.3 ± 212.4 U/L [28].

Alkaline phosphatase enzyme is involved in energy transfer reactions its utility in growth has been established. The plasma alkaline phosphatase activity in healthy goats was found to be 432.78 ± 20.00 IU/L [29]. Bhooshan et al (2010) [30] found alkaline phosphatase in Barbari and Jamunapari goats irrespective of age and season of birth were 44.62±1.30 and 35.49±1.53 KA units respectively. In local Israeli goats [31] and in Black Bengal goats [32], alkaline phosphatase decreased with increase in age.

In present study the mean value of serum calcium (mg/dl) was found to be 8.1± 0.97, which is towards the lower side of normal range. Calcium is necessary for milk production, muscle contraction, nerve conduction and blood clotting. Milk is relatively high in calcium, and lactating goats need adequate levels of calcium for milk production. Serum calcium has a critical role in the metabolism of the body including the cells of reproductive system. An excess of calcium can cause abnormal bone growth [33]. Similarly, in smallholder cattle production feed shortages in the dry season, diseases and parasites were

major challenges that affected the production and reproduction performance of smallholder cattle production [34].

The feed and feeding strategies in goat rearing have significant impact on the quality and nutritional value of the meat. Therefore, it is important for producers to carefully manage feeding practices to ensure that their goats produce high quality meat [35].

4. CONCLUSION

It is concluded that goats are reared on extensive system in Gwalior city of Central India, lack of scientific feeding and management practices causing lower total protein and calcium level and higher levels of blood urea and SGPT levels in blood and poor reproductive performance (higher age at first kidding and kidding interval and low twinning rate), thus, causing economic losses to the farmers.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. DAHD. Dairy, animal husbandry department, Government of India; 2022. Available. <http://dahd.nic.in/about-us/divisions/cattle-and-dairy-development>
2. Patil SJ, Suryagandh SS, Kamble DK, Mandakmale SD. Factors affecting the age at first kidding in local, angora and their crossbred goats. *The Asian Journal of Animal Science*. 2009;3(2):193-195.
3. National Bureau of Animal Genetic Resources, India; 2018. Available:<http://www.nbagr.res.in>.
4. Yavarifard R, Gjhazi Hossein-Zadeh N, Shadparvar AA. Estimation of genetic parameters for reproductive traits in Mehraban sheep. *Czech Journal of Animal Science*. 2015;60(6):281-288.
5. Madan J, Sindhu S, Gupta M, Kumar S. Hematobiochemical profile and mineral status in growing beetal goat kids. *Journal of Cell and Tissue Research*. 2016;16(1):12-13.
6. Jana C, Rahman FH, Mondal SK and Singh AK. Management practices and perceived constraints in goat rearing in Burdwan district of West Bengal. *Indian Research Journal of extension education*. 2016;14(2):107-110.
7. Ahmed A, Da, MA, Bhat AA, Jena B, Mishra GK and Tiwari RP. Study on haemato-biochemical profile in goats suffering from gastrointestinal parasitism in Jaipur district of Rajasthan. *Journal of livestock Science*. 2015;6:52-55.
8. Tariq KA, Chishti MZ, Ahmad F, Shawl AS. *Journal of Helminthology*. 2010;84:93-97.
9. Sharma P, Sharma D, Dogra PK and Mandial RK. Comparative efficacy of fenbendazole and oxiclozanide-tetramisole combination against gastrointestinal nematodes in naturally infected Gaddi goats. *Veterinary Research International*. 2014;2(1):15-17.
10. Maiti SK, Rao VN, Ali SL. Clinicohaematological and therapeutic studies in parasitic gastroenteritis in sheep. *Indian Veterinary Journal*. 1999;76(5):435-437.
11. Rai B, Singh MK. Production performance of Jakhrana goats in its home tract. *Indian Journal of Animal Sciences*. 2005;75(10):1176.
12. Patel AC, Pandey DP. Growth, production and reproduction performance of Mehsana goat. *Journal of Livestock Science*. 2013;4:17-21.
13. Gautam L, Waiz HA, Nagda RK. Evaluation of the reproductive characteristics of Sirohi goats from Udaipur India. *J Entomol Zool Stud*. 2018;6:13-7.
14. Arun K, Tomar AKS, Mehta BS. Gestational performance of Kutchi goats under semi-arid condition of Rajasthan. *Indian J. Dairy Sci*. 2004;57(4):255-57.
15. Acharya RM. Goat genetic resources and their management. *Research in Goat: Indian Experience*. Central Institute for Research on Goat, Makhdoom, Farah, Mathura (UP). 1992; 1-21
16. Singh MK, Rai B, Singh P, Singh PK, Singh NP. Goat population status in different agro-climatic regions of India: An overview. *Indian J. of Small Ruminants*. 2008;14:48-70.
17. Guljar ML, Pathodiya OP. Constraints perceived by farmers in goat rearing in Mewar region of southern Rajasthan. *Indian Journal of Animal Sciences*. 2008;78:124-126.
18. Meganathan N, Selvakumar KN, Prabhu M, Saravan P, Senthil KG. Constraint

- analysis of tribal livestock farming in Tamil Nadu. Tamil Nadu Journal of Veterinary and Animal Sciences. 2010;6 (1): 12-18.
19. Whitbread TJ. Clinical Biochemistry. Abbey Veterinary Services, MSD and the MSD Veterinary Manual; 2017
 20. Dukes HH. The physiology of domestic Animals (c.f. Melvin J. Swenson and William O. Reece Edn.) P.42, Comstock Publishing Associates. Ithaca and London; 1996.
 21. Khan JR, Ludri RS. Changes in blood glucose, plasma non- esterified fatty acids and insulin in pregnant and non-pregnant goats. Tropical Animal Health Production. 2002;34(1):81-90. 5.
 22. Satyanarayana U. Biochemistry. 2ndEdn., Books and Allied (P) Ltd., Chintamani Das Lane, Kolkata. 2002;45.
 23. Baumgartner W and Pernthaner A. Influence of age, season, and pregnancy upon blood parameters in Austrian Karakul sheep. Small ruminant research. 1994 13(2): 147-151.
 24. Shaikat AH, Hassan MM, Khan SA, Islam MN, Hoque MA, Bari MS, Hossain ME. Haemato-biochemical profiles of indigenous goats (*Capra hircus*) at Chittagong, Bangladesh. Veterinary World. 2015;6(10):789.
 25. Sakha M, Shamesdini M, Mohamad-Zadeh F. Serum biochemistry values in Raini goat of Iran. Iranian Journal of Veterinary Medicine. 2008;6:1-7.
 26. West HJ. Evolution of total plasma bile acid concentrations for the diagnosis of hepatobiliary diseases in horses. Res. Vet. Sci. 1989; 46: 264-270.
 27. Daramola JO, Adeloje AA, Fatoba TA, Soladoye AO. Haematological and biochemical parameters of West African Dwarf goats. Livestock Research for Rural Development. 2005;17(8):95.
 28. Perez JM, Gonzalez FJ, Granados FJ, Perez MC, Fandos P, Sorignier RC et al. Haematologic and Biochemical Reference Intervals for Spanish Ibex. Journal Wildlife Diseases. 2003;39: 209-215.
 29. Sandhu AK, Saini A, Randhawa RS. Haematological studies in healthy goats. Indian Veterinary Journal. 2001;78(7): 590– 93.
 30. Bhooshan N, Kumar P. Profile of ovarian and thyroid hormones, alkaline phosphatase and cholesterol at oestrous cycle, gestation and early lactation in blood plasma of Marwari goats. Indian Journal of Animal Sciences. 2007;77 (12):1233–37
 31. Bogin E, Shimshony A, Avidar Y, Israli B. Enzymes, metabolites and electrolytes levels in the blood of local Israeli goats. Zentralblatt-fur-Veterinarmedizin-A. 1981;28(2):135–40.
 32. Kalita DJ, Mahapatra M. Serum constituents and serum enzyme activities of Black Bengal kids. Indian Journal of Animal Research. 1998;32:38–40.
 33. Hart S. Meat Goat Nutrition. Pages 58-83 in Proc. 23rd Ann. Goat Field Day, Langston University, Langston, OK; 2008
 34. Tenagne A, Taye M, Dessie T, Muluneh B, Kebede D, Tarekegn GM. Management practices, reproductive performances, and production constraints of indigenous cattle in north-western parts of Ethiopia. NJAS: Impact in Agricultural and Life Sciences. 2023;95(1):2211533.
 35. Gagaoua M, Alessandrini L, Das A, Lamri M, Bhattacharya D, Nanda PK, Das AK. Intrinsic and extrinsic factors impacting fresh goat meat quality: An overview. Scientific Journal" Meat Technology. 2023;64(1):20-40.

© 2023 Gautam et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/106956>