



Effect of Substrates on Germination, Growth and Flowering of Different Winter Annuals

Arvind Malik ^a, D. S. Dahiya ^a, A. K. Godara ^a,
Vikas Kumar Sharma ^b, S. K. Sehrawat ^a and Raveena ^{a*}

^a Department of Horticulture, CCS Haryana Agricultural University, Hisar, Haryana – 125004, India.

^b Department of Horticulture, Galgotias University, Noida, UP – 201310, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2024/v36i34403

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/99934>

Original Research Article

Received: 15/03/2023

Accepted: 19/05/2023

Published: 06/02/2024

ABSTRACT

This study was conducted at Centre for Quality Plant Material, Chaudhary Charan Singh Haryana Agricultural University, Hisar during 2015-16 to 2017-18 to determine the effect of different substrates and their combinations on vegetative growth and flowering of winter annuals (Larkspur, Antirrhinum, California Poppy and Dahlia). Six types of substrates (cocopeat, perlite, vermiculite, vermicompost, Farm Yard Manure and sand) were mixed by volume to create seven different treatments. The seeds were sown in the month of October in different treatments to record vegetative and floral parameters. The results showed that using substrate, that mixture of cocopeat with vermiculite and perlite following rate 3:1:1, gave higher and early germination and longer seedling length, seedling root length duration of flowering and earliest flowering and is found suitable for winter annuals nursery raising, vegetative growth and flowering.

*Corresponding author: E-mail: sainiraveena28@gmail.com;

Keywords: Substrates; antirrhinum; california poppy; flowering; winter annuals.

1. INTRODUCTION

Floriculture is a fast emerging industry, as progress both scientifically and commercially. Today commercial floriculture is the most profitable business and expanding rapidly all over the world [1]. As demand for flowers has gradually increased, floriculture has become an important trade in agriculture in the commercial sector [2]. The term "annuals" when applied to herbaceous ornamentals, refers to plants that are grown from seeds, produce flowers, set seeds and complete their life cycle for only one season [3] and serve as essential components in any landscape plan [4]. They are frequently used as bedding plants, garden plants, rockery plants, window basket, cut flowers, loose flowers and herbaceous border in gardens [5].

Growing media not only plays an important role for seed germination and root development but also act as a source of nutrient for quality seedling [6-7]. Cultivation of flower crops in soil is inexpensive, but it brings some risks like soil-borne disease, insect-pest and poor drainage, which suppressed the development of root system [8]. Generally, media for seedling are composed of soil, organic matter, pond soil and sand. Supplementing of the sand is aimed to make media more porous, while the organic matter (vermicompost) is added so as to enrich adequate nutrients for the seedling. Cocopeat is an agricultural by-product obtained after the extraction of fiber from the coconut husk with acceptable pH, EC (electrical conductivity) and other chemical attributes [7], which is mixable with high cation exchange and water holding capacities [9]. Vermicompost has high porosity, aeration, drainage, and water holding capacity [10] and also contains most nutrients in plant-available forms such as nitrates, phosphates, and exchangeable calcium and soluble potassium [11]. Keeping in view the effect of different substrate combinations was studied on growth and flowering in different winter annuals.

2. MATERIALS AND METHODS

The experiment was carried at Centre for Quality Plant Material, Chaudhary Charan Singh Haryana Agricultural University, Hisar (Haryana). In the study, Larkspur, Antirrhinum, California poppy and Dahlia were used during the year 2015-16 to 2017-18. Six substrates cocopeat, perlite, vermiculite, vermicompost, FYM and

send were used to create the seven different treatments in different proportion (by volume) for cultivation. The seven-substrate mixtures with five replications /treatments (with 5 plants/replications) were arranged in single rows on a greenhouse trough. The treatments were:

- S₁ – cocopeat + vermicompost + perlite (1:1:1)
- S₂ – cocopeat + vermiculite + perlite (3:1:1)
- S₃ – cocopeat + vermiculite + perlite (1:1:1)
- S₄ – sand + vermicompost (9:1)
- S₅ – sand + FYM (9:1)
- S₆ – sand + vermiculite (9:1)
- S₇ – sand (control)

The seeds of different winter annuals were cultivated with above media combination during October after treating with fungicide. The greenhouse with facility of controlling temperature, humidity and light with automation system for irrigation and fertigation was used. After recording of data of seedling plants, the plants were transplanted were kept under uniform condition during the study period where all the management practices were carried out as per the package of practices. The data were analyzed according to the procedure for analysis of factorial randomized block design (Factorial RBD) as given by Panse and Sukatme [12]. The overall significance of difference among the treatments was tested, using critical differences (C.D.) at 5% level of significance. The results were statistically analyzed with the help of a windows based computer package OPSTAT (Operational Statistics) [13].

3. RESULTS AND DISCUSSION

For all substrates and their combinations, the germination rates were below 85% in the all winter annuals (Table 1). Results showed that seed sown in cocopeat + vermiculite + perlite, 3:1:1 (S₂), had the highest germination rate (81.5%) followed by S₃ (77.2%), while lowest germination percentage was recorded in sand (63.4%), which was statistically at par with S₆ (65.4%). Larkspur, Antirrhinum and California poppy resulted into significantly higher germination percentage (82.2%, 80.5%, 84.3%) than Dahlia (79.0%) with cocopeat + vermiculite + perlite (3:1:1). California poppy seeds resulted in maximum germination percentage (72.6%), which was statistically at par with antirrhinum (71.7%). It can be observed that the seeds of California poppy seeds grown with T₂ showed

highest germination percentage (84.3%), which was statistically at par with S₂T₁ (82.2%), S₂T₂ (80.5%), S₃T₃ (80.4%) and S₂T₄ (79.0%).

Significant difference in days taken to 50% germination due to interactive effect of substrates and different winter annuals were observed and present in Table 2. Lowest days for 50% germination (7.5 days) was recorded with S₂ (cocopeat + vermiculite + perlite, 3:1:1) followed by S₃ (9.5 days). With the same media (Cocopeat + vermiculite + perlite, 3:1:1) antirrhinum and California poppy resulted into significantly lesser number of days taken to 50% germination (6.2 and 6.8 days) than larkspur and dahalia (7.6 and 9.2 days).

The results indicate that the effect of different substrate combinations on germination was significant. It is hardly possible to obtain a good crop without successful seedling establishment and the period of germination and seedling emergence is most vulnerable stage in a crop life [14]. The results also agreed with the earlier findings of Anderson [15], who observed that cocopeat is an important component and has a profound impact on physical, chemical and biological properties of substrates and is known to enhance seedbed conditions for desired seed emergence. Poor seedling emergence in sand based substrate might be due to poor moisture holding capacity [16] physical properties [17] and diurnal temperature fluctuation in root zone [16].

The length of seedling differed significantly due to combinations of substrates and type of winter annuals (Table 3). The maximum seedling length (9.6 cm) was reported from S₂ (cocopeat + vermiculite + perlite, 3:1:1), followed by S₃ (8.2 cm). California poppy resulted into significantly higher seedling length (12.2 cm) followed by dahalia (8.1 cm) and antirrhinum (7.4 cm), while lowest seedling length (4.6 cm) was found in larkspur. Conclusively, California poppy grown in S₂ (cocopeat + vermiculite + perlite, 3:1:1) produced largest seedlings (12.2 cm), followed by S₁T₃ (11.2 cm), whereas, shortest seedlings (2.9 cm) was observed from S₇T₁ (2.9 cm), which was statistically at par with S₆T₁ (3.3 cm) and S₅T₁ (3.5 cm). It could be due to the reason that combination of 3 cocopeat: 1 vermiculite: 1 perlite improved the water and nutrient consumption and maintained porosity. This result is also in agreement with the work of Godara et al [18] who suggested that soilless substrate caused better exchange of elements especially cations inside the substrate and they distributed

humidity properly around the root zone and it was finally effective in shoot length.

The difference in seedling root length of different winter annuals with respect to the substrates was found significant (Table 4). The highest seedling root length (6.1 cm) was reported from S₂ (cocopeat + vermiculite + perlite, 3:1:1) followed by S₃ (5.1 cm) and S₁ (3.9 cm), while lowest seedling root length (3.2 cm) was found in S₄ (sand + vermicompost, 9:1) followed by sand alone (3.6 cm). California poppy resulted into significantly higher seedling root length (6.4 cm) followed by antirrhinum (3.7 cm), whereas, lowest seedling root length (3.3 cm) was reported in both larkspur and dahalia. The data given in Table 4 also showed that the combined use of California poppy with S₂, produced maximum seedling root length (8.2 cm) followed by S₃T₃ (7.6 cm), whereas, minimum seedling root length was measured from S₄T₄ (2.0 cm), which was statistically at par with S₄T₁ (2.1 cm), S₆T₄ (2.1 cm), S₅T₁ (2.4 cm), S₅T₂ (2.4 cm) and S₇T₄ (2.4 cm).

It could be due to the alteration of physico-chemical properties (such as porosity, moisture content and air capacity) of raw material and hence the air and water balance in the root environment. These results were agreed with Sharma et al [19], who reported that soilless substrate showed better water retention, air filled porosity, gas diffusion and nutrient availability to the root development.

The days taken to initiate first flower exhibited significant difference due to different combinations of substrates, however the interactions between substrate and different winter annuals were non-significant (Table 5). Among all the substrate combinations, cocopeat + vermiculite + perlite (1:1:1) resulted into lesser number of days taken to first flower (57.8 days), followed by S₂ (63.2 days), whereas, highest number of days taken to first flower (74.8 days) was reported from control. Dahalia took minimum days (68.1 days) to first flower, which was statistically at par with antirrhinum (68.9 days) and California poppy (69.4 days), while maximum days to first flower were observed from larkspur (72.5 days).

The data given in Table 6 showed that among the different substrate combinations tried, S₂ (cocopeat + vermiculite + perlite, 3:1:1) resulted into longer duration of flowering (40.7 days) followed by S₃ (37.9 days), whereas, shorter duration of flowering (29.7 days) was reported

from sand only. In case of different winter annuals, dahalia showed highest duration of flowering (37.2 days), which was statistically at par with California poppy (36.5 days) and minimum duration for flowering was observed from larkspur (30.2 days). The interactive effect of substrate and plant type was found non-significant.

Table 1. Effect of substrates on per cent germination of different winter annuals under polyhouse conditions

Treatment	Germination (%)				Mean
	Larkspur (T ₁)	Antirrhinum (T ₂)	California Poppy (T ₃)	Dahalia (T ₄)	
S ₁	70.2	78.4	72.3	66.9	72.0
S ₂	82.2	80.5	84.3	79.0	81.5
S ₃	78.5	73.3	80.4	76.5	77.2
S ₄	68.9	71.8	76.3	71.5	72.1
S ₅	60.5	63.5	65.8	74.8	66.2
S ₆	63.4	66.3	67.9	64.1	65.4
S ₇	65.2	68.4	61.3	58.5	63.4
Mean	69.8	71.7	72.6	70.2	
CD (p = 0.05)	A = 2.87		B = 2.17		AxB = 5.74

Table 2. Effect of substrates on days taken to 50% germination of different winter annuals under polyhouse conditions

Treatment	Days taken to 50% germination				Mean
	Larkspur (T ₁)	Antirrhinum (T ₂)	California Poppy (T ₃)	Dahalia (T ₄)	
S ₁	12.4	10.8	7.5	11.7	10.6
S ₂	7.6	6.2	6.8	9.2	7.5
S ₃	9.6	7.5	9.9	10.9	9.5
S ₄	14.3	8.4	11.8	9.8	11.1
S ₅	15.2	12.2	8.6	10.2	11.6
S ₆	13.5	9.8	8.2	12.1	10.9
S ₇	10.7	13.6	10.8	13.3	12.1
Mean	11.9	9.8	9.1	11.0	
CD (p = 0.05)	A = 0.42		B = 0.32		AxB = 0.85

Table 3. Effect of substrates on seedling length (cm) of different winter annuals under polyhouse conditions

Treatment	Seedling length (cm)				Mean
	Larkspur (T ₁)	Antirrhinum (T ₂)	California Poppy (T ₃)	Dahalia (T ₄)	
S ₁	4.9	7.1	11.2	9.2	8.1
S ₂	7.0	9.2	12.2	10.2	9.6
S ₃	6.2	7.3	10.8	8.4	8.2
S ₄	4.5	8.5	9.7	8.1	7.7
S ₅	3.5	7.6	10.3	7.8	7.3
S ₆	3.3	6.2	6.7	7.3	5.9
S ₇	2.9	5.7	8.5	5.6	5.7
Mean	4.6	7.4	9.9	8.1	
CD (p = 0.05)	A = 0.33		B = 0.25		AxB = 0.65

Table 4. Effect of substrates on seedling root length (cm) of different winter annuals under polyhouse conditions

Treatment	Seedling root length (cm)				Mean
	Larkspur (T ₁)	Antirrhinum (T ₂)	California Poppy (T ₃)	Dahalia (T ₄)	
S ₁	2.5	3.8	5.4	3.8	3.9
S ₂	5.1	5.8	8.2	5.2	6.1
S ₃	3.4	4.7	7.6	4.7	5.1
S ₄	2.1	4.2	4.7	2.0	3.2
S ₅	2.4	2.4	6.8	3.2	3.7
S ₆	4.8	8.9	5.8	2.1	3.7
S ₇	3.1	2.8	6.2	2.4	3.6
Mean	3.3	3.7	6.4	3.3	
CD (p = 0.05)	A = 0.18		B = 0.14	AxB = 0.37	

Table 5. Effect of substrates on days to first flower of different winter annuals under polyhouse conditions

Treatment	Days to first flower				Mean
	Larkspur (T ₁)	Antirrhinum (T ₂)	California Poppy (T ₃)	Dahalia (T ₄)	
S ₁	65.8	65.9	64.7	66.9	65.8
S ₂	62.9	63.8	63.2	63.1	63.2
S ₃	69.4	68.2	68.9	64.9	57.8
S ₄	76.9	69.1	72.6	67.5	71.5
S ₅	75.3	69.0	72.8	71.3	72.1
S ₆	78.3	72.3	69.8	70.6	72.8
S ₇	78.9	74.2	73.9	72.1	74.8
Mean	72.5	68.9	69.4	68.1	
CD (p = 0.05)	A = 0.75		B = 2.08	AxB = N.S.	

Table 6. Effect of substrates on duration of flowering (days) of different winter annuals under polyhouse conditions

Treatment	Duration of flowering (days)				Mean
	Larkspur (T ₁)	Antirrhinum (T ₂)	California Poppy (T ₃)	Dahalia (T ₄)	
S ₁	33.5	38.1	38.2	37.4	36.8
S ₂	34.9	41.7	42.8	43.2	40.7
S ₃	71.7	39.6	41.3	39.1	37.9
S ₄	28.1	33.4	35.3	36.2	33.2
S ₅	29.8	34.3	34.6	35.5	33.5
S ₆	27.1	32.9	33.7	36.8	32.6
S ₇	26.2	30.6	29.8	32.1	29.7
Mean	30.2	35.8	36.5	37.2	
CD (p = 0.05)	A = 1.31		B = 0.99	AxB = N.S.	

4. CONCLUSION

In the present studies all substrates exhibited significant effect on precocity and duration of flowering in different winter annuals. Among the different combinations, the S₃ has perhaps created the most appropriate condition for days to first flower in all winter annuals. This is might

be due to the physiochemical properties of the growing media pose their effect on the plant growth and flowering (20-21) and the composition of growth media is very important factor to be taken under consideration [22]. Nourizadeh [23] has also reported the increase flowering in plants due to suitable conditions in soilless substrate by ventilation and water

maintenance. According to Riaz et al. [24], coconut coir based growing media can significantly improve days to first flower and flower duration in zinnia.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Mahalle SL, Roy P, Thakur S. Prospect and challenges of flower in India. *Agriculture Letters*. 2020;01(08):23-26.
- Datta SK. Present status of research on floriculture in India. *LS - An International Journal of Life Sciences*. 2019;08(02):71-93.
- Sindhu K, Naik BH, Kantharaj Y, Chandrashekar SY, Ganapathi M. Evaluation of winter annuals for suitability as cut and loose flowers under Hill Zone of Karnataka. *International Journal of Current Microbiology and Applied Sciences*. 2020;9(10):2412-2422.
- Brown SH. Colorful plant beds for South Florida and similar climates. Lee County Extension, Fort Myers, Florida; 2012.
- Love SL, Noble K, Parkinson S, Bell S. Herbaceous ornamentals: annuals, perennials, and ornamental grasses. Short-Season, high altitude gardening. 2009;1-16. Accessed 14 April 2022. Available: <http://www.cals.uidaho.edu/edcomm/pdf/bul/bul0861.pdf>
- Wilson SB, Stoffellea PJ, Graetz DA. Use of compost as a media amendment for containerized production of two subtropical perennials. *Journal of Environmental Horticulture*. 2001;19:37-42.
- Abad M, Noguera P, Puchades R, Maquieira A, Noguera V. Physio-chemical and chemical properties of some coconut dusts for use as a peat substitute for containerized ornamental plants. *Bio-resource Technology*. 2002;82:241-245.
- Beattle DJ, White JW. Liliium-hybrids and species. In: De Hertogh A. A. and Le Nard M. (eds.). *The Physiology of Flower Bulbs*. Elsevier, Amsterdam; 1992.
- Inden H, Torres A. Comparison of four substrates on the growth and quality of tomatoes. *Acta Horticulturae*. 2004;644:205-210.
- Edwards CA, Burrows I. The potential of earthworm compost as plant growth media. In: Edwards, C.A. Neuhauser, E.F. (Eds.), *Earthworms in Waste and Environmental Management*. SPB Academic Publishers, The Netherlands; 1988.
- Orozco SH, Cegarra J, Trujillo LM, Roig A. Vermicomposting of coffee pulp using the earthworm (*Eisenia fetida*) effects on C and N contents and the availability of nutrients. *Biology and Fertility of Soils*. 1996;22:162-166.
- Panse VG Sukatme PV. Statistical methods for Agricultural workers. Indian Council of Agricultural Research, New Delhi; 1984.
- Sheoran OP, Tonk DS, Kaushik LS, Hasija RC, Pannu RS. Statistical software package for agricultural research workers. Recent advances in information theory, statistics & computer applications by D.S. Hooda & R.C. Hasija Department of Mathematics Statistics, CCS HAU, Hisar; 1998.
- Onemli F. The effects of soil organic matter on seedling emergence in sunflower. *Plant, Soil and Environment*. 2004;50(11):494-499.
- Anderson TH. Microbial eco-physiological indicators to assess soil quality. *Agriculture, Ecosystems & Environment*. 2003;98:285-293.
- Forbes JC, Watson RD. Plant growth and development; seed and seedling. *Plants in Agriculture*. Cambridge Univ. Press., New York, USA; 1992.
- Stewart BW, Albrecht SL, Skirvim KW. Vigor transport vs. seed-soil contact in wheat germination. *Agronomy Journal*. 1999;91:783-786.
- Godara AK, Sharma VK, Beniwal V. Effects of substrates and containers on root system of hydroponically grown strawberry (*Fragaria x ananassa* Duch.) under greenhouse. *Bangladesh J. Bot*. 2016;45(5):1083-1089.
- Sharma VK, Godara AK, Kumatkar RB. Effect of soilless growing structures on vegetative growth and roots of strawberry cv. Sweet Charlie. *The Ecoscan*. 2015;9(1/2):89-91.
- Sharma VK, Godara AK. Effect of substrate mixtures on precocity and flower development in strawberry potted plants. *Journal of Pharmacognosy and Phytochemistry*. 2017;6(5):2406-2412.
- Wilkerson D. Growing media. *Texas greenhouse management handbook*.

- Texas Agriculture Extension Service; 2002. Accessed 14 April 2023. Available:<http://aggie-horticulture.tamu.edu/greenhouse/nursery/guides/green/media.html>
22. Ingram DL, Henley RW, Yeager TH. Growth media for container grown ornamental plants (BUL 241). Environmental Horticulture Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida; 2003.
23. Nourizadeh M. The effect of different substrate cultivation on the growth, performance and to evaluate the cold acclimation and freezing quality of greenhouse cucumber in without soil cultivation system. M.Sc thesis on Horticulture, Guilan University, Iran; 2003.
24. Riaz A, Arshad M, Younis A, Raza A, Hameed M. Effects of different growing media on growth and flowering of *Zinnia Elegans* cv. Blue Point. Pakistan Journal of Botany. 2008;40(4):1579-1585.

© 2024 Malik et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/99934>