



Efficacy of Plant Growth Regulators on Different Yield Parameters of Cucumber cv. Malini under Shade Net Conditions

A. Dinesh ^{a*}, P. Prasanth ^b, D. Lakshminarayana ^b,
K. Nagaraju ^a and P. Gouthami ^c

^a Department of Vegetable Science, College of Horticulture, Mojerla, SKLTS Horticultural University, Rajendranagar, Hyderabad – 500032, India.

^b Department of Floriculture and Landscaping, College of Horticulture, Mojerla, SKLTS Horticultural University, Rajendranagar, Hyderabad – 500032, India.

^c Department of Crop Physiology, College of Horticulture, Mojerla, SKLTS Horticultural University, Rajendranagar, Hyderabad – 500032, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJSSPN/2024/v10i1229

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/113610>

Original Research Article

Received: 13/12/2023

Accepted: 21/02/2024

Published: 26/02/2024

ABSTRACT

The experiment was conducted to study the effect of different plant growth regulators on yield of cucumber cv. Malini under shade net conditions. The experiment was carried out with 10 treatments GA3 at 75 ppm, 150 ppm and 250 ppm, Ethrel at 100 ppm, 200 ppm and 300 ppm, Salicylic acid at 75 ppm, 150 ppm and 250 ppm and control (water spray) in 3 replications. The yield parameters like number of fruits per plant, fruit weight, fruit length etc. were significantly influenced by plant growth

*Corresponding author: Email: dineshaccha@gmail.com;

regulators. Among all yield parameters superior values for maximum days taken for final harvest (86.49 days), maximum fruit weight (248.00 g) were recorded in treatment GA₃ at 250 ppm, treatment GA₃ at 75 ppm was recorded with the highest values for number of fruits per vine (9.46) and fruit diameter (4.56cm) and GA₃ at 150 ppm recorded with highest values for fruit length (21.33 cm) as compared to other treatments. Among all treatments, treatment with GA₃@ 75 ppm showed positive effect in terms of gross returns and over all net returns.

Keywords: Efficacy of plant; plant effect; plant growth; yield parameters.

1. INTRODUCTION

“Cucumber (*Cucumis sativus* L.) is a vine crop belonging to the family Cucurbitaceae having chromosome number $2n=14$. cultivation of cucumber is possible in field conditions as well as under shade net conditions, but crop grown under shade net conditions provide more yields compared to field grown crops and under protected conditions three crops in a year can be grown, cultivation under shade net conditions provide more favorable conditions for better growth and development. Mostly cucumber is preferred as a summer vegetable, immature fruits are consumed raw or along with salt in salads. Consumption of cucumber helps people suffering from indigestion, constipation and jaundice” [1]. “Cucumber is an annual plant species which is found to be day neutral in nature, Cucumber is a monoecious, trailing vine having hirsute or scabrous stems with triangular ovate shape leaves with acute curves, leaf axils with unbranched lateral tendrils. As the lateral branches develop, flower clusters appear at each leaf axils whereas fruits are warty and cylindrical. However, enhancement in yield and quality are important factors for getting higher returns in any crop, which will be improved by different horticultural techniques. Among these, use of different plant growth regulators at appropriate concentration increases the quality and yields of any crop” [1]. Plant growth regulators play a significant role in the growth and development of plants by affecting a variety of plant functions. Plant growth regulators can be a useful tool for boosting production and quality in monoecious crops like cucumbers that are cultivated under protected settings. These regulators can increase the femaleness of the crop and facilitate effective pollination. On the other hand, farmers lack the necessary knowledge and skills to apply plant growth regulators at the right times and dosages to achieve the intended effects.

2. MATERIALS AND METHODS

To determine the impact of various plant growth regulators on the yield of cucumber cv. Malini at

the College of Horticulture, Mojerla, the current study was conducted in the late rabi season of 2018–19 using the application of various concentrations of plant growth regulators.

The experiment was laid out in Completely Randomized Design (CRD) along with three replications. The experiment comprised of 10 treatments of different concentrations of GA₃(75 ppm, 150 ppm, 250 ppm), Ethrel (100 ppm, 200 ppm, 300 ppm) and salicylic acid (75 ppm, 150 ppm, 250 ppm) along with water spray (control). Seeds were sown on the well prepared beds two seeds per hill during early days of December month on beds with dimensions of 100 x 40 x 50 cm (width, height & distance between two beds) and size of the plot was 21m. Plant growth regulators were sprayed with 15 days interval at 20 and 35 days after sowing. The data to various yield parameters viz ., Number of fruits per vine, Days taken for final harvest, Fruit diameter (cm), Length of fruits (cm), Fruit weight (cm) was recorded during the course of investigation for the experiments conducted. Uniformly growing five plants were randomly selected and tagged in each treatment and replication for the purpose of recording observations and the data was subjected to statistical analysis.

3. RESULTS AND DISCUSSION

3.1 Yield Parameters

From the results it was observed that Treatment GA₃ at 75 ppm was recorded with the highest values for number of fruits per vine (9.46) and fruit diameter (4.56 cm), the increased number of fruits per plant due to increased flowering with application of gibberellins and increased fruit set due to increased availability of nutrients from leaves by GA₃. These findings are in agreement with the results reported by Ghani et al. [2] in bitter gourd and Increased fruit diameter with gibberellins is mainly due to their general function cell division and cell elongation and increased translocation of metabolites from source to sink. The findings were in agreement

Table 1. Effect of plant growth regulators on number of fruits per vine, days taken to final harvest, fruit diameter (cm) fruit length (cm) and fruit weight (g) of cucumber (*Cucumis sativus* L.) cv. Malini

Treatment No.	Treatment details	Number of fruits per vine	SD	Days taken to final harvest	SD	Fruit diameter (cm)	SD	Fruit length(cm)	SD	Fruit weight(g)	SD
T ₁	GA ₃ 75 ppm	9.46 ^a	0.12	82.26 ^b	0.93	4.56 ^a	0.02	17.83 ^{bcd}	0.63	239.39 ^b	2.12
T ₂	GA ₃ 150ppm	9.08 ^b	0.20	83.68 ^b	1.31	3.71 ^b	0.17	21.33 ^a	0.53	238.67 ^b	3.51
T ₃	GA ₃ 250 ppm	9.06 ^b	0.18	86.49 ^a	0.65	3.62 ^{bc}	0.01	18.33 ^{bcd}	0.11	248.00 ^a	5.14
T ₄	Ethrel 100 ppm	8.65 ^c	0.29	79.06 ^{cd}	1.48	3.14 ^{cd}	0.34	16.68 ^d	0.01	234.67 ^b	4.04
T ₅	Ethrel 200 ppm	8.46 ^c	0.06	80.14 ^c	1.03	3.37 ^{bcd}	0.25	17.90 ^{bcd}	0.01	239.33 ^b	1.53
T ₆	Ethrel 300 ppm	8.12 ^d	0.03	77.20 ^{de}	0.93	2.94 ^{de}	0.19	17.00 ^{cd}	0.01	236.96 ^b	2.66
T ₇	Salicylicacid 75ppm	8.40 ^{cd}	0.17	85.89 ^a	1.04	3.07 ^d	0.49	18.75 ^{bc}	0.01	233.00 ^b	7.97
T ₈	Salicylicacid 150ppm	8.60 ^c	0.31	83.15 ^b	1.02	3.04 ^d	0.28	19.00 ^b	0.49	237.36 ^b	5.00
T ₉	Salicylicacid 250ppm	8.64 ^c	0.29	83.05 ^b	1.52	3.06 ^d	0.17	17.00 ^{cd}	0.43	235.67 ^b	6.51
T ₁₀	Water spray (control)	7.27 ^e	0.15	75.60 ^e	1.15	2.48 ^e	0.35	16.60 ^d	0.02	216.00 ^c	6.56
	SEM±	0.11		0.65		0.18		0.63		2.84	
	CD at 5%	0.34		1.93		0.54		1.88		8.39	

with Jyoti et al. [3] in cucumber and Rahman et al [4] in cucumber reported the similar findings where application of GA₃ increased the number of fruits per vine. Whereas GA₃ at 250 ppm was recorded for maximum days taken for final harvest (86.49 days) and maximum fruit weight (248.00 g). This is mainly due to gibberellins' increased metabolic activity of plants which resulted in the enhancement of reproductive phase which resulted in prolonged the days of harvesting and this prolonged harvesting remains as crux in reaping out higher yields. The main reason associated with increased fruit weight is that treated plants with gibberellins remain more active physiologically and build up sufficient food for developing fruits which results in the increased fruit weight. Increase in size of individual fruits with application of GA₃ also reported by Vadigeri et al. [5] in cucumber, Chowdary et al. [6] in cucumber where GA₃ 200 ppm significantly increased fruit weight and Nagamani et al. [7] who reported that application of GA₃ significantly enhanced fruit weight in bitter gourd. Sapkota et al. [8] also observed similar findings in cucumber. Finally, the highest fruit length was recorded in treatment with GA₃ at 150 ppm. GA₃ promotes cell division and cell elongation which would have favored uptake of water and nutrients. Results were in line with Prabhu et al. [9] in ivy gourd, similar effect with Gibberellic Acid application was reported by Jyoti et al. [3]. Pandey et al. [10] also published similar findings in cucumber.

3.2 Economic Analysis

Among all treatments GA₃ at 75 ppm (T₁) recorded the highest gross return (Rs.16, 51,700), highest net return (Rs.11, 27,783.22) and best benefit cost ratio (2.15) followed by GA₃ at 250 (T₃) with benefit cost ratio (2.13), whereas the lowest benefit cost ratio (1.62) was recorded in water spray (control) (T₁₀). This might be due to the positive effect towards other yield attributing characters and lower cost of chemicals due to lower concentration compared to T₂ and T₃ [11].

4. CONCLUSION

The current study suggests that, in shade net conditions, plant growth regulators significantly impacted the production characteristics of the cucumber variety "Malini." Out of all the treatments for plant growth regulators, plants treated with 75 ppm of GA₃ had the best B:C ratio, maximum gross returns, net returns, and

positive effects on growth, phenological, yield, and quality indicators.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Bailey LH. Manual of cultivated plants. Macmillan Company, New York. 1969;1116.
2. Ghani MA, Amjad M, Iqbal Q, Nawaz A, Ahmad T, Hafeez O, Abbas M. Efficacy of plant growth regulators on sex expression, earliness and yield components in bitter gourd. Pak. j. Life. soc. Sci. 2013;11(3):218-224.
3. Jyothi S, Patel JB. Effect of growth regulators and stages of spray on seed yield and seed quality parameters of ridge gourd [*Luffa acutangula* (Roxb) L.]. J. Appl. & Nat. Sci. 2016;8(3):1551–1555.
4. Rahman MA, Sikder S, Bahadur MM, Pramanik SK. Influence of gibberellic acid (GA₃) on growth, flowering, and fruit yield of cucumber. J. Sci. Technol. (Dinajpur). 2020;18:33-42.
5. Vadigeri BG, Madalageri BB. Response of cucumber genotypes to ethrel and GA₃. Karnataka J. of Agric. Sci. 2001;2:176-178.
6. Chowdhury RS, Kumar V, Bhattacharya S, Mallick P, Ghosh A, Bhattacharjee S, Kothari SK. Effect of gibberellic acid (GA₃) on vegetative and reproductive growth and yield characters of cucumber (*Cucumis sativus*) under Coastal Region of West Bengal, India. Int. J. Plant Soil Sci. 2023;30(21):90-96.
7. Nagamani SA, Basu SA, Singh SA, Lal SKA, Behera TKC, Chakrabarty SKB, Talukdar AD. Effect of plant growth regulators on sex expression, fruit setting, seed yield and quality in the parental lines for hybrid seed production in bitter gourd (*Momordica charantia*). Int. J. Agr. Sci. 2015;85(9):1185–91.
8. Sapkota B, Dhital M, Shrestha B, Tripathi KM. Effect of plant growth regulators on flowering and fruit yield of cucumber (*Cucumis sativus* cv. malini) in chitwan, Nepal. Journal of Agriculture and Forestry University. 2020;4:161-167.
9. Prabhu M, Natarajan S. Effect of growth regulators on fruit characters and seediness in ivy gourd (*Coccinia grandis*

- L.). Agric. Sci. Digest. 2006;26(3):188–190.
10. Pandey P, Shukla. Upadhyay A. Effect of different plant growth regulators on growth, yield and quality parameters of cucumber (*Cucumis sativus* L.) cv. Kalyanpur green. Journal of Pharmacognosy and Phytochemistry. 2021;10(1):2681-2684.
11. Panse VG, Sukhatma PV. Staistical methods for agricultural workers. Indian Council of Agricultural Research, New Delhi; 1985.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/113610>