



# Comparative Evaluation of Different Biocontrol Agents against Anthracnose of Nendran Banana Caused by *Colletotrichum musae*

Arun. A.T. <sup>a\*</sup>, Pramod. R <sup>a</sup>, Radhakrishnan N.V <sup>a</sup>,  
Susha S. Thara <sup>a</sup>, Reji Rani O. P <sup>b</sup> and Anuradha. T. <sup>c</sup>

<sup>a</sup> Department of Plant Pathology, College of Agriculture, Vellayani, Kerala Agricultural University, Thrissur, (Kerala), India.

<sup>b</sup> Department of Entomology, College of Agriculture, Vellayani, Kerala Agricultural University, Thrissur, (Kerala), India.

<sup>c</sup> Department of Plant Biotechnology, College of Agriculture, Vellayani, Kerala Agricultural University, Thrissur, (Kerala), India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i6889>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/117308>

**Original Research Article**

**Received: 08/03/2024**

**Accepted: 10/05/2024**

**Published: 13/05/2024**

## ABSTRACT

India is the largest producer of banana in the world. Nendran, a French plantain, is the most important banana cultivar in Kerala. It is mainly affected by many diseases both in the field and after harvest. Being a freshly consumed fruit crop there are a lot of limitations in using chemicals for its disease control. Keeping this view, a study was conducted at the Department of Plant Pathology, College of Agriculture, Vellayani, to control anthracnose, a serious post harvest disease

\*Corresponding author: E-mail: [arunkaralmanna55@gmail.com](mailto:arunkaralmanna55@gmail.com);

**Cite as:** Arun. A.T., Pramod. R, Radhakrishnan N.V, Thara, S. S., Reji Rani O. P, & Anuradha. T. (2024). Comparative Evaluation of Different Biocontrol Agents against Anthracnose of Nendran Banana Caused by *Colletotrichum musae*. *Journal of Advances in Biology & Biotechnology*, 27(6), 302–309. <https://doi.org/10.9734/jabb/2024/v27i6889>

under tropical conditions. In this study comparative evaluation of three fungal biocontrol agents namely [*Trichoderma* sp. (KAU), *Trichoderma* sp. (TRPN-3), *Trichoderma* sp. (TRKR-2)] and bacterial biocontrol agents [*Pseudomonas fluorescens* (PN026), *Bacillus* sp. (B15), *Bacillus* sp. (B17)] were tested against *Colletotrichum musae*. Among the fungal biocontrol agents maximum mycelial growth inhibition was recorded with *Trichoderma* sp. (KAU) (77.44%) followed by *Trichoderma* sp. (TRKR-2) (77.11%). Minimum mycelial growth inhibition was recorded in *P.fluorescens* (PN026) (55.33%). Based on the results reveals that *Trichoderma* sp. (KAU) is a highly promising biocontrol agent for managing anthracnose of banana.

**Keywords:** Anthracnose; biocontrol; *Colletotrichum musae*; banana; *Trichoderma* sp; *Pseudomonas fluorescens*, *Bacillus* sp.

## 1. INTRODUCTION

“Banana is the second most important fruit crop in India next to mango. Its year round availability, affordability, varietal range, taste, nutritive and medicinal value makes it the favourite fruit among all classes of people. Besides India, other major banana producing countries are China, Philippines, Ecuador, Brazil and Indonesia” [1]. “India produces a total of 23 per cent of the entire world production of banana” [2].

“The area of banana cultivation in Kerala estimated about 60678 ha, with a production of 548425 tonnes” [3]. “Nendran, a French plantain cultivar belonging to Musa AAB group is the most important banana variety in Kerala. Owing to its unique taste and flavour it has great suitability for crispy chips. Being a climacteric fruit, post harvest loss in banana is huge and it is about 21.67%” [4].

“The genus *Colletotrichum* includes fungal species that are responsible for anthracnose diseases in different vegetable and fruit crops worldwide, including bananas” [5,6]. “Anthracnose of banana is a major catastrophic post harvest disease in India” [7]. “Symptoms of anthracnose in bananas caused by *C. musae* include black sunken lesions and the presence of salmon colored conidia” [8]. “In India, postharvest losses to disease, including anthracnose, have been estimated at 30-40%” [9]. “Anthracnose caused by *C.musae*, is a major constraint in the marketing of fruit intended for local as well as distant markets. It causes lesions on the fruit peel after ripening, as well as finger and crown rots. It affects fruit quality severely, resulting in significant economic losses in many parts of the tropics” [10].

“To increase the product yield, pesticides are overused and misused in spite of health concerns, environmental contamination and

export bans” [11]. “Build up of resistance to pathogens and severe environmental pollution, however, are some of the drawbacks from the prolonged application of chemical fungicides such as benomyl, prochloraz and azoxystrobin. A major concern for pesticide treated fruit is the issue of food safety that has received increased attention from the consumers in recent years. Hence, most overseas markets are no longer permitting chemical treatments for fruit entering their countries” [12]. There is an urgent need to find alternative ways to replace or reduce the use of synthetic chemicals.

“Biological control is a crop treatment which potentially can be used individually or in consortium with other compatible microbes, organic amendments to reduce losses caused by plant pathogens” [13]. “Fungal biocontrol agents, *Trichoderma* spp. are widely studied for its ability to produce mycoparasitic enzymes like chitinase,  $\beta$ -1,3-glucanase, pectinase, and cellulase, saprophytic ability, competition for space and nutrition, induction of systemic resistance in host helps in eco friendly management of plant diseases” [14]. “Bacterial biocontrol agents including *Pseudomonas* spp., *Bacillus* spp., and *Streptomyces* are mainly used for plant growth promotion and disease management and it includes various effective combination methods to control fruit decay” [15]. “Biological control has been recommended as an alternative method for controlling postharvest pathogens” [16,17]. In this study, comparative evaluation of fungal and bacterial biocontrol agents efficacy against *C.musae*, causing anthracnose of Nendran banana.

## 2. MATERIALS AND METHODS

### 2.1 Sampling and Isolation of Pathogen

The diseased Nendran banana samples were collected from Kuzhalmannam (10°43'17.1"N

76°35'47.0"E) Palakkad district of Kerala, India. After the symptomatology studies, the diseased fruits were subjected to pathogen isolation. "The diseased fruits were thoroughly washed in tap water, followed by washing with sterile water and allowed to dry. Inside the laminar air flow chamber, small bits (6 sq. mm area) of fruit peel were cut out at the margin where the diseased and healthy tissues met. The bits were placed in 0.1 per cent mercuric chloride for 60 seconds for surface sterilization and subsequently were transferred to sterile water and rinsed three times to remove residue of the chemical. The bits were allowed to dry over a sterile blotting paper. These bits were then transferred to Petri plates with solidified Potato Dextrose Agar (PDA) medium" [18]. The plates were wrapped using cling film and kept for incubation at room temperature (28±1°C). When mycelial growth initiation from the bits commenced, pure culturing was done. The hyphal tip method was used to obtain pure culture from incubated plates, the tip of the mycelia in the media was cut out using an inoculation needle or a cork borer and was inoculated into sterile petri plates with solidified PDA medium. The Petri plates were kept at 28±1°C. When pure cultures were obtained, the fungi were also transferred into PDA slants and kept at 4 °C further studies.

## 2.2 Morphological Characterization

Morphological features were used to select isolates resembling *Colletotrichum musae*. The selected isolate were cultured on PDA plates at 28±1 °C for 7days. 9 mm mycelial disc from colony margins were placed on the centre of 90 mm diameter Petri plate and incubated at 28±1°C. The colony colour and culture diameters (two perpendicular directions) were recorded at regular interval. Colony diameters were used to calculate hyphal growth rate (mm/day) and the length and width of conidia were measured.

## 2.3 Screening of different biocontrol agents against *C.musae*

Antagonistic activity of different fungal and bacterial biocontrol agents against anthracnose of banana pathogen *C.musae* was done by dual culture plate method [19]. "In dual culture, for fungal bio-agents, fifteen ml of PDA medium was poured into sterilized Petri plate and allowed for solidification. Nine mm diameter discs from actively growing colony of pathogen was cut with a sterile cork borer and placed near the periphery

of PDA plate. Similarly, bio-agents were placed on the other side *i.e.*, at an angle of 180°. Petri plate with pathogen served as control. The plates were incubated at 28±1°C for seven days. For each treatment, three replications were maintained. The extent antagonistic activity by bio-agents were recorded after incubation period of 7 days by measuring the growth of the test pathogen in dual culture and in control plates. For bacterial bio-agents nutrient agar medium was used". [20] Per cent inhibition over control was calculated according to formula given by Vincent [20].

$$\text{Percentage of inhibition (PI)} = \frac{C-T}{C} \times 100$$

Where,

PI = Per cent inhibition of mycelial growth  
C = Mycelial growth of pathogen in control (mm.)

T = Mycelial growth of pathogen in treatment (mm.)

## 2.4 Statistical Analysis

Experiments were performed in triplicates, and collected data were analysed by ANOVA using GRAPES (General R based Analysis Platform Empowered by Statistics) [21]. Experiment design was completely randomised design (CRD) for *in vitro* experiment.

## 3. RESULTS AND DISCUSSION

### 3.1 Sampling and Isolation

Anthrachnose infected banana samples were collected from Kuzhalmannam, Palakkad district of Kerala (Fig.1). *Colletotrichum musae*, the pathogen responsible for anthracnose disease was isolated from the collected sample and pure cultured. The culture were salmon in colour in Petri plate after seven days of growth and the cylindrical shaped conidia of the pathogen was observed under microscope (Fig. 2,3).

### 3.2 Morphological Characterization


The *C. musae* colonies typically grew rapidly on PDA. Initially, colonies appeared light pink colour and, later turned into salmon colour. The reverse side of culture plate appeared light saffron colour. In Petri plate there is clear visible of black dots *i.e.*, acervuli, which are visible to the naked eye. Conidia are typically hyaline (translucent) and single celled. Shape of conidia is in cylindrical and size is 15.82×4.85 µm.



**Fig. 1. Symptom of Anthracnose of Nendran Banana collected from Kuzhalmannam, Palakkad district of Kerala**



**Fig. 2. PDA culture plate showing 7 days old culture of *C. musae***



**Fig. 3. Microscopic view of *C. musae* (40X)**

### 3.3 Screening of Different Biocontrol Agents against *C. musae*

The present study was focused on comparative evaluation of fungal bio agents viz., *Trichoderma* sp. (KAU), *Trichoderma* sp. (TRPN-3), *Trichoderma* sp.(TRKR-2) and bacterial biocontrol agents includes *P. fluorescens* (PN026), *Bacillus* sp.(B15) and *Bacillus* sp.(B17) were tested *in vitro* against growth of *C. musae* (Table 1) (Fig. 4, 5).Maximum mycelial growth inhibition of the pathogen was recorded with *Trichoderma* sp. (KAU)(77.44 per cent) followed by *Trichoderma* sp. (TRKR-2) (77.11 per cent) by using fungal biocontrol agents. In case of screening with bacterial biocontrol agents maximum mycelial growth inhibition of the pathogen was recorded with *Bacillus* sp. (B17) (55.55 per cent) followed by *Bacillus* sp. (B15) (55.44 per cent). In mango the evaluation of six biocontrol agents viz., *T. hamatum*, *T. harzianum*, *T. koningii*, *T. longibrachiatum*, *T. pseudokoningii* and *T. viride* against *Colletotrichum gleosporioides* *in vitro* by dual culture technique. Among these isolates *T. harzianum* was the most effective, reducing of pathogen growth by 42 per cent. *T. viride*, *T. harzianum* and *T. koningii* were efficient in inhibiting the growth of *C. musae* [22,23].

“Antagonistic effect of *T. longibrachiatum*, *T. viride*, *T. asperellum* and *T. harzianum* against *C. gloeosporioides* in dual culture” [24]. “Similarly treatment with each *Trichoderma* species reduced radial growth of *C. gloeosporioides* in plates by 53–56 per cent in papaya. *Trichoderma harzianum* strain DGA01 has been reported to be effective in managing postharvest pathogens in banana, namely *Lasiodiplodia theobromae*, *C. musae*, *Thielaviopsis paradoxa* and *Fusarium verticillioides* during *in vitro* and *in vivo* evaluations” [25]. “Many species from *Trichoderma* are widely studied against destructive pathogens including *Alternation sesami* causing leaf spot in sesame, stem rot of groundnut caused by *Sclerotium rolfsii*” [26,27]. Recently, 11 species of native *Trichoderma* from 16 crop rhizosphere have been identified and their antagonistic potential were evaluated for the management of *Sclerotium rolfsi* and *Rhizoctonia solani* [28]. “To develop alternatives to chemical fungicides is triggered by consumers demand for food safety, environmental concern and increasing resistance of pathogens. The use of antagonistic microorganisms offers a sustainable and promising eco friendly alternative to chemicals like the genus *Trichoderma* spp. that manages major postharvest decays of fruits” [29].





Fig. 4. *In vitro* screening of different biocontrol agents against *C. musae*



**Fig. 5. Evaluation of different bio control agents T1- *P. fluorescens*, T2-*Trichoderma* sp. (KAU), T3-*Trichoderma* sp. (TRPN-3), T4-*Trichoderma* sp. (TRKR-2), T5-*Bacillus* sp. (B15), T6-*Bacillus* sp. (B17) T7-Pathogen inoculated control T8-Absolute control, against *C. musae***

**Table 1. *In vitro* screening of different biocontrol agents against *C.musae***

| S.No | Treatments                      | Mean Radial Growth (mm) | Per Cent Inhibition over Control (%) |
|------|---------------------------------|-------------------------|--------------------------------------|
| T1   | <i>P.fluorescens</i> (PN026)    | 40.20*                  | 55.33(48.06)**                       |
| T2   | <i>Trichoderma</i> sp. (KAU)    | 20.30                   | 77.44(61.64)                         |
| T3   | <i>Trichoderma</i> sp. (TRPN-3) | 20.80                   | 76.88(61.26)                         |
| T4   | <i>Trichoderma</i> sp. (TRKR-2) | 20.60                   | 77.11(61.42)                         |
| T5   | <i>Bacillus</i> sp. (B15)       | 40.10                   | 55.44(48.12)                         |
| T6   | <i>Bacillus</i> sp. (B17)       | 40.00                   | 55.55(48.19)                         |
| T7   | Pathogen inoculated control     | 90.00                   | 0.00(0.00)                           |
| T8   | Absolute control                | 0.00                    | 0.00(0.00)                           |
|      | CD 1%                           |                         | 1.406                                |
|      | SEM                             |                         | 0.469                                |

\*Average of three replications, \*\*Values in parentheses are angular transformed data

#### 4. CONCLUSION

Based on our present study, out of six biocontrol agents, *Trichoderma* sp. (KAU) was found most effective in inhibiting mycelial growth of *C.musae* followed by *Trichoderma* sp. (TRKR-2) and *Trichoderma* sp. (TRPN-3). *P.fluorescens* (PN026) was least effective in *in vitro* (Dual culture method) condition.

#### FUTURE SCOPE

Based on the obtained results from present study indicated that, *Trichoderma* sp. (KAU) exhibiting excellent mycelial growth control of anthracnose

pathogen *C.musae* in *in vitro* condition could be utilized in *in vivo* conditions for the management of anthracnose of Nendran banana.

#### ACKNOWLEDGEMENTS

Sincere gratitude is extended by the authors to College of Agriculture, Vellayani, Thiruvananthapuram for providing all the facilities required to carry out the reserach experiment.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

- Doss DA, Mazhar SH, Brahma U. Study on information needs regarding cultivation and marketing constraints of banana growers in Agasteeswaram Taluk of Kanyakumari district of Tamil Nadu. *International Journal of Advances in Agricultural Science and Technology*. 2020;7(10):27-33.
- Bhagat AA, Bhoge BSR. Markov chain analysis on the export prospectus of Banana in India; 2022.
- Agriculture statistics, A compilation and analysis of reports on Agriculture Statistics published by DES from 2005-06 to 2019-20, Directorate of economics and statistics, Govt.of Kerala; 2022.
- Saha CK, Ahamed MK, Hosen MS, Nandi R, Kabir M. Post-harvest losses of banana in fresh produce marketing chain in Tangail District of Bangladesh. *Journal of the Bangladesh Agricultural University*. 2021; 19(3):389-97.
- Cannon PF, Damm U, Johnston PR, Weir BS. *Colletotrichum*: Current status and future directions. *Studies in Mycology*. 2012;73(1):181-213.
- Rogério F, Ciampi-Guillard M, Barbieri MCG, Bragança CAD, Seixas CDS, Almeida AMR, Massola Jr NS. Phylogeny and variability of *Colletotrichum truncatum* associated with soybean anthracnose in Brazil. *Journal of Applied Microbiology*. 2017;122(2):402-415.
- Arun AT. Anthracnose of banana: A major catastrophic post harvest disease in India. *Agri Articles (e-magazine for Agricultural Articles)*. 2023;03(06).
- Freeman S, Katan T, Shabi E. Characterization of *Colletotrichum* species responsible for anthracnose diseases of various fruits. *Plant Disease*. 1998;82(6): 596-605.
- Dhua RS, Ghosh SK, Sen S. Standardisation of harvest maturity of banana cultivar Giant Governor in West Bengal. *Advances in Horticulture and Forestry*. 1992;2:24-35.
- Jeger M, Plumbly RA. The biology and control of *Colletotrichum* species on tropical fruit crops. *Plant Pathology*. 1990; 39:343-366.
- Panuwet P, Prapamontol T, Chantara S, Thavornyuthikarn P, Montesano MA, Whitehead Jr RD, Barr DB. Concentrations of urinary pesticide metabolites in small-scale farmers in Chiang Mai Province, Thailand. *Science of the Total Environment*. 2008;407(1):655-668.
- Jacobi KK, Giles JE. Quality of Kensington'mango (*Mangifera indica* Linn.) fruit following combined vapour heat disinfestation and hot water disease control treatments. *Postharvest Biology and Technology*. 1997;12(3):285-292.
- Sharma P, Jambhulkar PP, Raja M, Sain SK, Javeria S. *Trichoderma* spp. in consortium and their rhizospheric interactions. In: Sharma A, Sharma P (eds) *Trichoderma*. Springer. 2020;267–292.
- Sharma P, Sharma M, Raja M, Shanmugam V. Status of *Trichoderma* research in India: A review. *Indian Phytopathology* 2014;67:1–19.15. Janisiewicz, WJ, Korsten L; 2002.
- Biological control of postharvest diseases of fruits. *Annual Review of Phytopathology*. 40:411–441.
- Droby S, Wisniewski M, Teixidó N, Spadaro D, Jijakli, MH. The science, development, and commercialization of postharvest biocontrol products. *Postharvest Biology and Technology*. 2016;122:22-29.
- Dukare AS, Paul S, Nambi VE, Gupta RK, Singh R, Sharma K, Vishwakarma RK. Exploitation of microbial antagonists for the control of postharvest diseases of fruits: A review. *Critical Reviews in Food Science and Nutrition*. 2018;59(9):1498-1513.
- Rahimloo T, Ghosta Y. The occurrence of *Alternaria* species on cabbage in Iran. *Zemdirbyste-Agriculture*, 2015;102(3).
- Dennis, C, Webster J. Antagonistic properties of species-groups of *Trichoderma*: I. Production of non-volatile antibiotics. *Transactions of the British Mycological Society*. 1971;57(1):25-IN3.
- Vincent JM. Distortion of fungal hyphae in the presence of certain inhibitors. *Nature*. 1947;159(4051):850-850.
- Gopinath PP, Parsad R, Joseph B, Adarsh VS. GRAPES: General Rshiny Based Analysis Platform Empowered by Statistics; 2020. Available:https://www.kaugrapes.com/home. version 1.0.0. DOI: 10.5281/zenodo.4923220.
- Prabakar K, Raguchander T, Saravanakumar D, Muthulakshmi P, Parthiban VK, Prakasam V. Management of postharvest disease of mango anthracnose incited by *Colletotrichum*



- gleosporioides*. Archives of Phytopathology and Plant Protection. 2008;41(5):333-339.
23. Sangeetha G, Usharani S, Muthukumar A. Biocontrol with *Trichoderma* species for the management of postharvest crown rot of banana. *Phytopathologia Mediterranea*. 2009;48(2):214-225.
24. Valenzuela NL, Angel DN, Ortiz DT, Rosas RA, García CFO, Santos MO. Biological control of anthracnose by postharvest application of *Trichoderma* spp. on maradol papaya fruit. *Biological Control*. 2015;91:88-93.
25. Alvindia DG, Natsuaki KT. Evaluation of fungal epiphytes isolated from banana fruit surfaces for biocontrol of banana crown rot disease. *Crop Protection*. 2008;27(8): 1200 -1207.
26. Arun AT, Singh M. Management of *Alternaria* Leaf spot of Sesame (*Sesamum indicum* L.) in in vitro using different Biocontrol agents. *Biological Forum – An International Journal*. 2023;15(10):1639-1642.
27. Raja M, Sharma RK, Jambhulkar PP, Pandian RTP, Sharma P. Comparative evaluation of native *Trichoderma* species from groundnut rhizosphere against stem rot caused by *Sclerotium rolfsii* Sacc. *Indian Phytopathology* 2023; 76(459-471).
28. Jambhulkar PP, Singh B, Raja M. Genetic diversity and antagonistic properties of *Trichoderma* strains from the crop rhizospheres in southern Rajasthan, India. *Sci Rep*. 2024;14, 8610.
29. Janisiewicz WJ, Korsten L. Biological control of postharvest diseases of fruits. *Annual Review of Phytopathology*. 2002;40:411–441.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/117308>