



Assessment of Climate Change Impact and Adaptation Strategies for Rainfed Maize (*Zea mays* L.)

S. Akshaya^a, S. Kokilavani^{a*}, S. Pazhanivelan^b,
Ga. Dheebakaran^a, K. Boomiraj^a and S. Mohan Kumar^a

^a Agro Climate Research Centre, Tamil Nadu Agricultural University, Coimbatore – 641003, India.

^b Centre for Water and Geospatial Studies, Tamil Nadu Agricultural University, Coimbatore – 641003, India.

Authors' contributions

This work was carried out in collaboration among all authors. Authors SA and SK carried out verification of output, and wrote the protocol and the first draft of the manuscript. Authors SK and KB generated the outputs. Authors GD and SMK defined the methodology of the research and verification part. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i102620

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/104307>

Original Research Article

Received: 02/06/2023

Accepted: 04/08/2023

Published: 12/08/2023

ABSTRACT

The study was performed to assess the impact of climate change on spatiotemporal changes in rainfed maize yield. Climate projections data of MIROC-ESM-CHEM model from CMIP6 was used for future climatic scenarios in the maize growing areas of Dindigul and Perambalur districts of Tamil Nadu. The DSSAT model was used to simulate maize yield and evaluate adaptation strategies for base period (1991-2020), the mid (2040-2069) and end centuries (2070-2099) under SSP245 and SSP585 scenarios. The simulation finding shows that, in all scenarios maize yield declined in both Dindigul (7 to 9% and 11 to 12%) and Perambalur (6 to 9% and 11 to 13%) during mid and end centuries respectively from the base period (1991-2020). Following the adaptation strategies such as delayed sowing, the yield was increased in both Dindigul (5 to 6% and 4 to 5%)

*Corresponding author: E-mail: kokilavani.s@tnau.ac.in;

and Perambalur (4 to 5% and 5 to 6%) with respect to normal sowing date. The results of this study would help in developing adaptation strategies for minimizing the adverse effects of the projected climate in maize-growing districts of Tamil Nadu.

Keywords: SSP 245; SSP 585; climate change; impact; adaptation.

1. INTRODUCTION

Maize (*Zea mays L*) is the third most important crop grown next to rice and wheat in India. Globally maize (dry grain) is primarily used as feed, between 2020 and 2021, cereal production has grown by 64 million tonnes driven by 4.1 per cent increases in maize production [1]. The leading producers are USA (32.61%), followed by China (22.91%), Brazil (9.42%), European Union (8.41%), Argentina (5.41%) and India (4.1%) [2]. At the global level, maize (dry grain) is primarily used as feed (56% of production) [1]. In India the area under cultivation of maize is raised to 9.38 million hectares, yielding 28.752 million metric tonnes [3]. Maize grain contains approximately 72% starch, 10% protein, 4% fat and 365 Kcal/100g energy, whereas green fodder contains 10-12% of crude protein and 25-30% crude fibre. It is a day neutral plant with C4 type of photosynthesis and has very efficient utilization of solar radiation which is known as the "Queen of Cereals" due to its photo-thermo insensitive nature and the highest genetic potential for yield [4].

By 2050 the global population is projected to reach around 10 billion and global cereal equivalent food demand is expected to rise by around 10,094 million tonnes in 2030 and 14,886 million tonnes in 2050 [5]. As a result, enhancing crop production and productivity to meet rising demand is inevitable. Maize production is expected to approach around 50 million metric tonnes by 2025 to cope with the growing population [6]. Earlier studies in India generally confirm the trend of agricultural fall owing to climate change [7-10].

The Coupled Model Inter-comparison Project (CMIP) is the multi-model experimental framework by IPCC (Intergovernmental Panel on Climate Change) for bringing out information on changing climate. The IPCC sixth assessment report (AR6) featured a new state of CMIP6 models with a new set of emissions scenarios driven by various Shared Socioeconomic Pathways. With current land use, projections indicate a loss of 4-5 million tonnes production for every 1°C increase in temperature

during the growing season [10]. Rainfall influences maize growth during monsoon season by influencing soil moisture availability. Impact of elevated temperature is negated by increase in rainfall. Rainfall increases of 20% and 30% are expected to reimburse for the yield loss brought about by 1°C and 2°C temperature increases, respectively [11].

Climate change's impact on crop production has been extensively researched using statistical models and process-based crop models. The Decision Support System for Agrotechnology Transfer (DSSAT) is a windows based user friendly version that uses a variety of input variables, including crop, soil, and weather data that can simulate multiyear outcomes of crop management strategies for different crops [12]. Autonomous adaptation such as delayed date of sowing can be put forward for changing environment in order to meet the demands for future. This study was conducted to know the impact of climate change on maize production under normal and delayed date of sowing.

2. MATERIALS AND METHODS

2.1 Study Area

The most efficient cropping zones for maize are Theni, Perambalur, Dindigul, Salem, Trichy, Ariyalur, Erode and Tiruppur [13,14] in which Perambalur and Dindigul have been chosen as study areas (Fig. 1). Perambalur district spread between 11° 13' 59.99" N and 78° 52' 59.99" E has a total geographical area of 175736 ha and a net sown area of 2,16,422 ha. The soil is primarily red loamy and black. Average rainfall is 908 mm in which 52%, 34% and 14% of annual rainfall is contributed by northeast monsoon, southwest monsoon, and the winter hot weather period accounts respectively. The area and production of Perambalur is 67183 ha and 601209 tonnes respectively in 2021- 2022 [15]

Dindigul district lies between 10° 22' 2.3232" N and 77° 58' 49.0476" E with a total geographical area is 626664 ha, and the net sown area is 253541 ha, accounting for nearly 40% of the total geographical area. The district is dominated by




Fig. 1. Study area map

red and black soil, followed by red sandy soil. The average rainfall is 836.0 mm among which northeast monsoon, southwest monsoon, summer and winter period accounts for 50%, 26%, 18% and 5% of annual rainfall, respectively. The area and production of Dindigul is 27649 ha and 223741 tonnes respectively in 2021-2022 [15].

2.2 Methodology

For base period (1991-2020) maximum temperature, minimum temperature and precipitation data were taken from India Meteorological Department (IMD) and MIROC-ESM-CHEM model were chosen for future climatic projection, SSP245 represents the medium pathway of future greenhouse gas

emissions as an update to RCP4.5, with an additional radiative forcing of 4.5 W/m² by the year 2100 and SSP585 with an additional radiative forcing of 8.5 W/m² by the year 2100. It can be interpreted as an update to the CMIP5 scenario RCP8.5, now with socioeconomic considerations these two climatic scenarios were selected, the parameters of Tmax, Tmin, precipitation and solar radiation was taken for Dindigul and Perambalur district. The future climate projection was done for two-time period mid (2040-2069) and end centuries (2070-2099). This data was taken as an input to run the DSSAT model. CERES maize module of DSSAT performed well around the Asian conditions [16] was chosen to run maize COHM6 under normal date of sowing (30th June) and delayed date of sowing as adaptation strategy (15th July) to

project the future yield for both mid and end centuries in SSP245 and SSP585 scenarios in rainfed condition.

2.3 Weather Data

For the base period 1991-2020, data on daily maximum and minimum temperatures at $1^\circ \times 1^\circ$ resolution and daily rainfall at $0.25^\circ \times 0.25^\circ$ resolutions were obtained from the National Data Centre (NDC), India Meteorological Department (IMD), Pune.

The IPCC is the United Nations (UN) intergovernmental body empowered with regularly assessing the science of the Earth's changing climate and determining the state of insights into climate change. The World Climate Research Programme (WCRP) organised the Sixth Phase of the Coupled Model Intercomparison Project (CMIP6). CMIP is evolving into an integrated framework for organising a number of individual Model Intercomparison Projects (MIPs). MIPs are bundles of experiments and simulations deployed to test and compare specific aspects of climate models. These climate projections will be moulded by a new set of emissions and land use scenarios [17] designed around new societal development future pathways, the Shared Socioeconomic Pathways (SSPs) and RCPs. CMIP 6 global climate model data was collected from NASA Earth Exchange Global Daily Downscaled Projections at a resolution of $0.25^\circ \times 0.25^\circ$ for maximum temperature, minimum temperature and rainfall. Among the General Circulation Model (GCMs) MIROC-ESM-CHEM model was chosen for this study because study conducted by [18] showed that MIROC-ESM-CHEM model was best for precipitation and temperature projections for tropical riverbasin in India when compared to all other models. The daily weather data on maximum temperature ($^\circ\text{C}$), minimum temperature ($^\circ\text{C}$), solar radiation ($\text{MJ}/\text{m}^2/\text{day}$), and rainfall (mm) for SSP254 and SSP585 scenarios were converted into DSSAT weather file format using the Weatherman tool available in DSSAT for running CERES- Maize. In DSSAT CERES maize module performed well in Asian conditions [16].

2.4 Soil Data

District wise soil data was obtained from the database incorporates soil characteristic parameters of 623 Indian districts gleaned from the National Bureau of Soil Survey and Land Use

Planning (NBSS & LUP) soil database/map [19]. From the 27 Generic Soil Profiles presented by [20], compatible soil profiles for each district were allocated based on soil texture, rooting depth, and organic carbon content (proxy of fertility). HC27 soil profiles have been widely employed in crop modelling studies at the regional and global levels [21-24] this soil profile was used as an input database file in 'S Build' to create a soil file in DSSAT.

2.5 Crop Data

Maize COHM6 which is 110 days crop of parentage UMI 1200 x UMI 1230. It is grown in both rainfed and irrigated condition. Crop management information such as sowing date, spacing, planting depth, planting population fertilizer dosage and irrigation schedule is referred from crop production guide 2020. The crop management file (XFile) allows the inputs to the models to be simulated for each experiment to replicate the field situation with an assumption of pest and disease free condition.

2.6 Adaptation Strategies

In previous studies different adaptation strategies had been adopted to minimize the impact of future climate change. The adaptation strategies include different dates of sowing [11,25-28], increase in fertilizer dosage [29,30] and irrigation scheduling [31-30]. In this study changes in the sowing data was adopted, the normal date of sowing (30th June) was fixed from the information gathered by the farmers and late date of sowing (15th July).

3. RESULTS AND DISCUSSION


The average maize yield for Dindigul was 4553 kg/ha during the baseline period (1991–2020); however, under the SSSP245 scenarios, the yield was reduced by 3947 kg/ha (7%) and 3860 kg/ha (9%), and under the SSSP585 scenarios, the yield was reduced by 3777 kg/ha (11%) and 3742 kg/ha (12%) for the mid (2040-2069) and end centuries (2070-2099), respectively. In this district the projected mean Tmax (32°C) and Tmin (24°C) was increased from the baseline Tmax (30°C) and Tmin (22°C). During the base period (1991–2020), the average maize yield in Perambalur district was 4031 kg/ha; however, under the SSSP245 scenarios, the yield was reduced by 3785 kg/ha (6%) and 3655 kg/ha (9%), and under the SSSP585 scenarios, the yield was reduced by 3566 kg/ha (11%) and 3488 kg/ha (13%), respectively, for the mid

(2040–2069) and end centuries (2070–2099). The projected mean Tmax (35°C) and Tmin (26°C) was increased from the baseline Tmax (34°C) and Tmin (26°C). Under rainfed conditions, a 1°C increase in both Tmax and Tmin reduces crop yield by 1.55% and 0.88%, respectively [33]. It causes faster accumulation of growing degree days, resulting in a reduction in total crop duration; increases evaporation and transpiration, which increases total crop water requirement; and increases Tmin, resulting in a higher rate of night respiration. All of these have a negative impact on crop yield [30]. Figs. 2 & 3 depicts the projected changes in mean Tmax (°C) and Tmin (°C) for normal and delayed sowing dates in the mid and end of the century under SSP245 and SSP585 in comparison to the baseline during the crop growth period.

With respect to the normal date of planting, in delayed sowing under SSP245, Dindigul yield grew from 5% (4060 kg/ha) to 6% (4178 kg/ha), while Perambalur yield raised by 4% (3795 kg/ha) to 5% (3695 kg/ha) whereas in SSP585, the yield was increased by 4% (3929 kg/ha) and 5% (3923 kg/ha) for Dindigul, and by 5% (3741) to 6% (3695 kg/ha) for Perambalur for the mid

(2040-2069) and end (2070-2099) centuries, respectively. In both the district the Tmax and Tmin doesn't have any significant deviation during the crop growth period for normal and delayed sowing but there was an increase in average rainfall for the crop growth period (9 to 15% and 10 to 12% in Dindigul for SSP245 and SSP 585 respectively) (13 to 22% and 15 to 17% in Perambalur for SSP245 and SSP 585 respectively). Figs. 4 & 5 shows the projected changes in mean Rainfall (mm) and rainy days (days) in the mid and end of the century for (a) normal and (b) delayed sowing dates under SSP245 and SSP585 in comparison to the baseline during crop growth period. The reason for increase in projected rainfall is due to increase in rainy days 2 to 4 days in Dindigul and 4 to 5 days in Perambalur for both the scenarios. Previous studies conducted by [34] showed that delaying the planting date could increase the yield. Crop may benefit over delayed sowing by avoiding heat and drought stress during the early growth stage [35] and water stress during flowering time [36]. Saddique et al. [37] stated that planting date at July is a good adaptation practice to protect maize yield in future.

(a) Normal date of sowing



(b) Delayed date of sowing





Fig. 2. Depicts the projected changes in mean Tmax (°C) for (a) normal date of sowing and (b) delayed sowing dates in the mid and end of the century under SSP245 and SSP585 in comparison to the baseline during the crop growth period

(a) Normal date of sowing



(b) Delayed date of sowing





Fig. 3. Depicts the projected changes in mean Tmin (°C) for (a) normal date of sowing and (b) delayed sowing dates in the mid and end of the century under SSP245 and SSP585 in comparison to the baseline during the crop growth period

(a) Normal date of sowing



(b) Delayed date of sowing





Fig. 4. Shows the projected changes in mean Rainfall (mm) in the mid and end of the century for (a)normal and (b)delayed sowing dates under SSP245 and SSP585 in comparison to the baseline during crop growth period.

(a) Normal date of sowing




(b) Delayed date of sowing

Fig. 5. Shows the projected changes in mean Rainy days (days) in the mid and end of the century for (a) normal and (b) delayed sowing dates under SSP245 and SSP585 in comparison to the baseline during crop growth period

4. CONCLUSION

Assessing the effects of climate change on maize yields is essential for developing adaptation strategies to mitigate the negative effects of climate change and ensure food security. The DSSAT crop simulation model was used to estimate maize yield for future scenarios (SSP245 and SSP585) in efficient cropping districts of Dindigul and Perambalur. Delayed sowing was investigated as an adaptation strategy for the mid and late centuries. Because maize is a C4 crop, the effect of rising temperatures caused by climate change is far more significant than the effect of rising CO₂. The mean Tmax, Tmin, rainfall, and total rainy days increased consistently in the study areas for both future scenarios. Maize yield would be reduced in the future if adaptation strategies were not implemented. In both districts, the delayed date of sowing yielded a higher yield than the normal date of sowing. Adaptation options differed quantitatively depending on location and season. As a result, breeding cultivars with phenology in changed scenarios similar to current varieties is required. Plant breeders should consider the phenology variation of different maize genotypes when developing varieties for future climate change scenarios. This study has the potential to quantify future spatiotemporal changes in maize yield, as well as adaptation strategies to mitigate the negative effects of climate change.

ACKNOWLEDGEMENT

The authors are thankful to the Agro Climate Research Centre and Centre for Water and Geospatial Studies for providing the fund through ET-RS&GIS scheme to carry out the research work in a project mode and like to extend our sincere thanks to all staff members, seniors and my batchmates for their valuable comments and constructive suggestions on the manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. FAOSTAT. Food and Agriculture Organization of the United Nations; 2021. Available: <http://faostat.fao.org/site/567/default.aspx#ancor>
2. USDA. Foreign agricultural service. World agricultural production. Circular series WAP 8-20; 2020. Available: <https://apps.fas.usda.gov/psdonline/circulars/production.pdf>
3. IndiaAgristat. Area, production, and average yield, Directorate of Economics and Statistics, Department of Agriculture and Cooperation report, New Delhi. 2022; <http://www.indiaagrstat.com/>
4. Hulmani S, Salakinkop SR, Somangouda G. Productivity, nutrient use efficiency,

- energetic, and economics of winter maize in south India. *Plos one*. 2022; 17(7):e0266886.
5. Islam S, Karim Z. World's Demand for Food and Water: The Consequences of Climate Change [Online First], Intech Open. DOI: 10.5772/intechopen. 85919.
 6. Sandhu R, Irmak S. Assessment of AquaCrop model in simulating maize canopy cover, soil-water, evapotranspiration, yield, and water productivity for different planting dates and densities under irrigated and rainfed conditions. *Agricultural Water Management*. 2019;224:105753.
 7. Saseendran SA, Singh KK, Rathore LS, Singh SV, Sinha SK. Effects of climate change on rice production in the tropical humid climate of Kerala, India. *Climatic Change*. 2000 Mar;44:495-514.
 8. Aggarwal PK, Mall RK. Climate change and rice yields in diverse agro-environments of India. II. Effect of uncertainties in scenarios and crop models on impact assessment. *Climatic Change*. 2002;52(3):331-43.
 9. Mall RK, Aggarwal PK. Climate change and rice yields in diverse agro-environments of India. I. Evaluation of impact assessment models. *Climatic Change*. 2002;52:315-30.
 10. Aggarwal PK. Global climate change and Indian agriculture: impacts, adaptation and mitigation. *Indian Journal of Agricultural Sciences*. 2008;78(11):911.
 11. Byjesh K, Kumar SN, Aggarwal PK. Simulating impacts, potential adaptation and vulnerability of maize to climate change in India. *Mitigation and adaptation strategies for global change*. 2010;15: 413-31.
 12. Venkatesan M, Pazhanivelan S. Estimation of Maize yield at spatial level using DSSAT crop simulation model. *Madras Agricultural Journal*. 2018;1:105.
 13. Kokilavani S, Geethalakshmi V. Identification of efficient cropping zone for rice, maize and groundnut in Tamil Nadu. *Indian Journal of Science and Technology*. 2013;6(10):5298-301.
 14. Abinaya J, Janaki Rani A, Kokilavani S, Nirmala Devi M, Gangaiselvi R. Examining the Prospective zones for Maize and Sorghum in Tamil Nadu. *Biological Forum – An International Journal*. 2022; 14(2a): 209212.
 15. Season and crop report, Department of Economics and Statistics, Government of Tamil Nadu, 2021-22.
 16. Harinarayanan MN, Dheebakaran Ga. Usability of monthly ERFs (Extended Range Forecast System) to predict maize yield using DSSAT (Decision Support System for Agro-technology Transfer) model over Erode District of Tamil Nadu. *Journal of Applied and Natural Science*. 2022;14(SI):244-50.
 17. Riahi K, Van Vuuren DP, Kriegler E, Edmonds J, O'Neill BC, Fujimori S, Bauer N, Calvin K, Dellink R, Fricko O, Lutz W. The Shared Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions implications: An overview. *Global Environmental Change*. 2017;42: 153-68.
 18. Jose DM, Dwarakish GS. Ranking of downscaled CMIP5 and CMIP6 GCMs at a basin scale: case study of a tropical river basin on the South West coast of India. *Arabian Journal of Geosciences*. 2022;15(1):120.
 19. Bhattacharyya T, Ray SK, Sarkar D, Balbuddhe DV, Dasgupta D, Chandran P, Tiwari P, Pal DK, Mandal C, Nimje AM, Deshmukh AS. Soil resource information of different agro-eco subregions of India for crop and soil modelling. National Project on climate change (ICAR Network Project), National Bureau of Soil Survey and Land Use Planning, Nagpur, Maharashtra; 2011.
 20. Koo J, Dimes J. HC27 Generic soil profile database. Harvard Dataverse ver. 4. Washington, DC: International Food Policy Research Institute; 2013.
 21. Nelson GC, Rosegrant MW, Koo J, Robertson R, Sulser T, Zhu T, Ringle C, Msangi S, Palazzo A, Batka M, Magalhaes M. Climate change: Impact on agriculture and costs of adaptation. *Intl Food Policy Res Inst*; 2009.
 22. Dourte DR, Fraise CW, Uryasev O. WaterFootprint on AgroClimate: A dynamic, web-based tool for comparing agricultural systems. *Agricultural Systems*. 2014;125:33-41.
 23. Müller C, Robertson RD. Projecting future crop productivity for global economic modeling. *Agricultural Economics*. 2014; 45(1):37-50.
 24. Kim S, Myoung B, Stack D, Kim J, Hatzopoulos N, Kafatos M. Sensitivity of simulated maize crop yields to

- regional climate in the Southwestern United States. InAGU Fall Meeting Abstracts. 2013:GC11D1035.
25. Boomiraj K, Wani SP, Aggarwal PK, Palanisami K. Climate change adaptation strategies for agro-ecosystem– A review. *Journal of Agrometeorology*. 2010;12(2): 145-60.
 26. Lashkari A, Alizadeh A, Rezaei EE, Bannayan M. Mitigation of climate change impacts on maize productivity in northeast of Iran: a simulation study. *Mitigation and adaptation strategies for global change*. 2012;17:1-6.
 27. Jalota SK, Kaur H, Kaur S, Vashisht BB. Impact of climate change scenarios on yield, water and nitrogen balance and-use efficiency of rice–wheat cropping system. *Agricultural Water Management*. 2013; 116:29-38.
 28. Rao AS, Shanker AK, Rao VU, Rao VN, Singh AK, Kumari P, Singh CB, Verma PK, Kumar PV, Bapuji Rao B, Dhakar R. Predicting irrigated and rainfed rice yield under projected climate change scenarios in the eastern region of India. *Environmental Modeling & Assessment*. 2016;21:17-30.
 29. Pramod VP, Rao BB, Ramakrishna SS, Singh MM, Patel NR, Sandeep VM, Rao VU, Chowdary PS, Rao VN, Kumar PV. Impact of projected climate on wheat yield in India and its adaptation strategies. *Journal of Agrometeorology*. 2017;19(3): 207-16.
 30. Rao AS, Chandran MS, Bal SK, Pramod VP, Sandeep VM, Manikandan N, Raju BM, Prabhakar M, Islam A, Kumar SN, Singh VK. Evaluating area-specific adaptation strategies for rainfed maize under future climates of India. *Science of The Total Environment*. 2022;836:155511.
 31. Xu CC, Wu WX, Ge QS, Zhou Y, Lin YM, Li YM. Simulating climate change impacts and potential adaptations on rice yields in the Sichuan Basin, China. *Mitigation and adaptation strategies for global change*. 2017;22:56594.
 32. Ma L, Ahuja LR, Islam A, Trout TJ, Saseendran SA, Malone RW. Modeling yield and biomass responses of maize cultivars to climate change under full and deficit irrigation. *Agricultural Water Management*. 2017;180:88-98.
 33. Srivastava RK, Panda RK, Chakraborty A. Assessment of climate change impact on maize yield and yield attributes under different climate change scenarios in eastern India. *Ecological Indicators*. 2021;120:106881.
 34. Deb P, Shrestha S, Babel MS. Forecasting climate change impacts and evaluation of adaptation options for maize cropping in the hilly terrain of Himalayas: Sikkim, India. *Theoretical and Applied Climatology*. 2015;121:649-67.
 35. Kassie BT, Asseng S, Rotter RP, Hengsdijk H, Ruane AC, Van Ittersum MK. Exploring climate change impacts and adaptation options for maize production in the Central Rift Valley of Ethiopia using different climate change scenarios and crop models. *Climatic Change*. 2015;129:145-58.
 36. Chen C, Baethgen WE, Robertson A. Contributions of individual variation in temperature, solar radiation and precipitation to crop yield in the North China Plain, 1961–2003. *Climatic change*. 2013;116:767-88.
 37. Saddique Q, Cai H, Xu J, Ajaz A, He J, Yu Q, Wang Y, Chen H, Khan MI, Liu DL, He L. Analyzing adaptation strategies for maize production under future climate change in Guanzhong Plain, China. *Mitigation and adaptation strategies for global change*. 2020; 25:1523-43.

© 2023 Akshaya et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/104307>