



Etiology, Diagnosis and Treatment for Puncta and Canalicular Obstruction: A Comprehensive Analysis

Foivos Papadopoulos ^{a*}

^a *Ophthalmic Plastic Center of Athens, Greece.*

Author's contribution

The sole author designed, analyzed, interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/JAMMR/2024/v36i55419

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/115075>

Review Article

Received: 22/01/2024

Accepted: 25/03/2024

Published: 06/04/2024

ABSTRACT

Puncta and canalicular obstruction is a common ophthalmological condition that can lead to significant morbidity and impaired tear drainage. This comprehensive analysis aims to provide an overview of the etiological factors and risk factors associated with this condition, along with a detailed examination of the anatomy of the lacrimal drainage system. Diagnostic techniques for assessing puncta and canalicular obstruction, including clinical evaluation, patient history, imaging techniques, and tear drainage tests, are discussed. Additionally, various treatment options for managing this condition are presented, including the management of specific etiological factors such as inflammation, infections, neoplastic conditions, and anatomical abnormalities. The article also addresses potential complications and provides insights into the prognosis of puncta and canalicular obstruction. By elucidating the multifaceted aspects of this condition, this comprehensive analysis aims to contribute to the understanding and management of puncta and canalicular obstruction, ultimately improving patient outcomes.

*Corresponding author: E-mail: phoevosnt5@gmail.com;

Keywords: Puncta obstruction; canalicular obstruction; lacrimal drainage system; etiology; anatomical abnormalities.

1. INTRODUCTION

Puncta and canalicular occlusion are serious problems that can interfere with the lacrimal drainage system's normal operation, resulting in compromised tear drainage and ocular discomfort. Tears enter the lacrimal drainage system through the puncta, tiny apertures situated at the inner corner of the eyelids. The canaliculi, which link the puncta to the lacrimal sac and subsequently to the nasal cavity, are where tears continue their journey from there.

The natural flow of tears is disrupted when the puncta or canaliculi become blocked or occluded, leading to a variety of clinical symptoms. Epiphora, recurring eye infections, persistent inflammation, and discomfort are all possible effects of this illness. Injury, infections, inflammatory conditions, structural anomalies, and trauma are only a few causes of puncta and canalicular blockage [1].

In ophthalmology, understanding the underlying causes and processes of puncta and canalicular blockage is crucial and has big therapeutic consequences. It makes it possible for medical professionals to correctly identify and treat the problem, relieving the symptoms in afflicted people and averting consequences. The development of novel therapeutic approaches and surgical procedures aimed at reestablishing regular tear drainage and enhancing patients' quality of life is also facilitated by research in this field.

This study intends to address important goals and queries in light of the importance of puncta and canalicular blockage. Investigating the etiology and risk factors connected to puncta and canalicular blockage, finding innovative diagnostic techniques, and assessing the effectiveness of current treatment options are the main goals. This study also aims to investigate potential preventative approaches and provide customized therapies to deal with certain blockage sources.

This study aims to offer insightful information that can inform clinical treatment and improve patient outcomes by illuminating the underlying processes and therapeutic approaches for puncta and canalicular blockage. Ultimately, increasing our knowledge of this problem will

open the door to more individualized and efficient methods for diagnosing, treating, and preventing puncta and canalicular blockage, which will be beneficial to those who are impacted by this ocular disorder.

2. ANATOMY OF THE LACRIMAL DRAINAGE SYSTEM

The lacrimal drainage system is essential for preserving ocular surface health by enabling appropriate tear drainage from the ocular surface into the nasal cavity. The puncta, canaliculi, lacrimal sac, and nasolacrimal duct are all associated components in this system.

The puncta, also known as lacrimal puncta or lacrimal canaliculi, are tiny apertures in the eyelids' inner corners. Each eye has two puncta located on the top and lower eyelids. These microscopic orifices allow tears to enter the lacrimal drainage system. Tears, which are generated by the lacrimal glands, include a combination of water, electrolytes, proteins, and other components that are required to maintain a healthy ocular surface.

Tears enter the canaliculi from the puncta, which are thin tubular structures that stretch from the puncta to the lacrimal sac. Through peristaltic motions, the canaliculi are lined with specialized epithelial cells that assist drive tears towards the sac. The canaliculi are classified into two types: horizontal canaliculi that run closer to the eyelid edge and vertical canaliculi that extend more vertically towards the lacrimal sac [2].

2.1 Overview of the Lacrimal Sac and Nasolacrimal Duct

The tears then enter the lacrimal sac via the canaliculi. The lacrimal sac is a dilated structure that serves as a tear reservoir. It's in the lacrimal fossa, a bony dip on the side of the nose next to the medial canthus. The lacrimal sac is coated with a mucous membrane and contains muscle walls that contract when you blink, allowing tears to flow into the nasolacrimal duct.

The nasolacrimal duct is a thin, bony tube that extends from the lacrimal sac and enters into the nasal cavity's inferior meatus. Its main purpose is to allow tears from the lacrimal sac to flow into the nasal cavity. This drainage process keeps

excess tears, dirt, and foreign particles off the ocular surface, preserving its clarity and health.

Unobstructed tear drainage is critical for the general health of the ocular surface. Efficient tear drainage prevents tears from accumulating on the ocular surface, which can cause visual abnormalities, discomfort, and possible corneal and conjunctival damage. Tears are constantly renewed and distributed to assist lubricate the ocular surface, wipe away foreign bodies and detritus, and maintain the optimum hydration and nutrients essential for optimal eye health [3].

The natural flow of tears is disturbed when there is an impediment or blockage in any section of the lacrimal drainage system. Trauma, infections, inflammatory illnesses, structural anomalies, and iatrogenic causes such as surgical procedures can all produce puncta and canalicular blockage. When the puncta or canaliculi get clogged, tears cannot drain correctly, resulting in excessive weeping (epiphora), which can be both socially and functionally inconvenient.

Furthermore, a clogged lacrimal drainage system can result in stagnant tears, allowing bacteria and other germs to flourish. Recurrent eye infections, persistent inflammation of the conjunctiva and eyelids, and pain can all result from this. In extreme cases, the blockage can cause dacryocystitis, an infection of the lacrimal sac characterized by discomfort, edema, and redness in the medial canthal area [4].

Finally, the lacrimal drainage system, which includes the puncta, canaliculi, lacrimal sac, and nasolacrimal duct, is critical for the health and operation of the ocular surface. The puncta serve as entrance sites for tears into the system, while the canaliculi use peristaltic motions to transport tears towards the lacrimal sac. The lacrimal sac collects tears before draining them into the nasal cavity via the nasolacrimal duct.

2.2 Importance of Unobstructed Tear Drainage

Unobstructed tear drainage is essential for a variety of reasons. For starters, it guarantees that excess tears, dirt, and foreign particles are eliminated from the ocular surface quickly, reducing visual disruptions and preserving ocular clarity. The continual replenishment of tears also lubricates the ocular surface, avoiding dryness and discomfort. Tears supply nutrients and hydration for the cornea and conjunctiva, encouraging its health and integrity.

The natural flow of tears is disturbed when puncta and canalicular blockage develop, resulting in a variety of clinical symptoms. Excessive crying, or epiphora, can occur when tears are unable to drain adequately from the ocular surface. This can be both socially and functionally inconvenient, lowering an individual's quality of life. Furthermore, stagnant tears on the ocular surface might foster bacterial development, raising the risk of recurring eye infections and chronic inflammation [5].

Additionally, untreated puncta and canalicular blockage might lead to more serious consequences. The occlusion can lead to infection of the lacrimal sac, culminating in dacryocystitis. Dacryocystitis causes discomfort, swelling, and redness in the medial canthal area and need immediate medical attention. To avoid problems and reduce symptoms, it is critical to appropriately detect and manage puncta and canalicular blockage.

It is critical to understand the architecture and function of the lacrimal drainage system in order to establish suitable diagnostic and therapeutic procedures. Clinicians may now reliably determine the location and type of the blockage because to advances in diagnostic modalities such as lacrimal probing, irrigation, and imaging methods. This understanding enables targeted interventions ranging from conservative measures such as warm compresses and topical medicines to surgical procedures such as dacryocystorhinostomy (DCR) or canalicular repair [6].

This research tries to achieve a number of goals. It aims to examine the underlying causes and risk factors related with puncta and canalicular blockage, shedding light on the condition's origin. Furthermore, researchers try to create novel diagnostic tools and imaging techniques that allow for early and accurate diagnosis of the blockage. This information can assist physicians in tailoring treatment approaches and improving patient outcomes.

Furthermore, the effectiveness of existing treatment options for puncta and canalicular occlusion is a critical topic of investigation. Researchers can enhance and optimize treatment regimens by examining the outcomes of various approaches, increasing success rates and reducing problems. This study also looks into preventive methods for puncta and canalicular blockage, such as identifying

modifiable risk factors and applying suitable preventive techniques [5].

To summarize, knowing the architecture and significance of unrestricted tear drainage through the lacrimal drainage system is critical for ocular surface health. Tear flow can be disrupted by puncta and canalicular blockage, resulting in excessive tearing, recurrent infections, and pain. The goal of this research is to discover the reasons, enhance diagnostic tools, refine treatment options, and investigate prevention approaches, with the goal of eventually improving patient outcomes and quality of life.

3. RISK FACTORS FOR PUNCTA AND CANALICULAR OBSTRUCTION

Puncta and canalicular blockage can be caused by a number of circumstances, including age-related changes, trauma, inflammatory and viral causes, systemic disorders, and certain drugs. Understanding these risk variables is critical for puncta and canalicular blockage diagnosis, therapy, and prevention. This chapter will go through these risk factors in depth.

Age-related variables influence the development of puncta and canalicular blockage. The structural and functional alterations of the lacrimal drainage system occur as people age. The puncta may become smaller and less open, preventing normal tear drainage. Furthermore, the canaliculi may lose flexibility and become more prone to blockage. Furthermore, changes in the composition and quality of tears with age might lead to the creation of clogs in the drainage system. These characteristics, taken together, enhance the incidence of puncta and canalicular blockage in the elderly [1].

Puncta and canalicular blockage can result from trauma and injury to the eye and adjacent tissues. Direct trauma to the eye or nasal area, for example, might cause injury to the puncta or canaliculi, leading in scarring or constriction of these structures. Inadvertent obstructions can also be caused by surgical operations affecting the eyelids or the lacrimal drainage system. Furthermore, foreign materials or chemical irritants in the eye might cause inflammation and eventual blockage. Prompt examination and treatment of traumatic injuries is critical for avoiding long-term consequences such puncta and canalicular blockage [7].

Puncta and canalicular blockage are frequently connected with inflammatory and infectious

causes. Inflammation of the eyelids (blepharitis) or conjunctiva (conjunctivitis) can cause excessive mucus or debris to form, which can clog the puncta or canaliculi. Chronic inflammatory ailments such as granulomatous disorders (such as sarcoidosis) or autoimmune diseases (such as Sjögren's syndrome) can also lead to blockage development. Recurrent styes or dacryocystitis, for example, can cause scarring and constriction of the lacrimal drainage system, impeding tear flow. Early detection and treatment of these inflammatory and infectious disorders are critical for avoiding long-term consequences [8].

Certain systemic disorders and drugs have been linked to puncta and canalicular blockage. Systemic diseases such as diabetes and thyroid problems can impair the integrity and function of the lacrimal drainage system, increasing the risk of blockage. Anticholinergic medications or those that promote mucous membrane dryness, such as antihistamines and some antidepressants, might affect tear generation and quality, predisposing people to blockage. To reduce the incidence of puncta and canalicular blockage, healthcare practitioners must consider these criteria while prescribing drugs and managing patients with systemic disorders [9].

To summarize, a number of risk factors lead to the formation of puncta and canalicular blockage. The pathophysiology of this syndrome is influenced by age-related changes, trauma, inflammatory and viral causes, as well as systemic disorders and drugs. Recognizing these risk factors is critical for identifying at-risk individuals, aiding early diagnosis, and adopting effective management options. More study in this area is required to better understand the processes behind these risk factors and to create tailored therapies to successfully prevent and cure puncta and canalicular blockage.

4. COMMON ETIOLOGICAL FACTORS

Puncta and canalicular blockage can be caused by a variety of acquired and congenital conditions, as well as foreign substances or anatomical abnormalities. Understanding these aspects is critical for making an accurate diagnosis and treating individuals with lacrimal drainage system dysfunction [10]. In this chapter, we will go over each of these etiological elements in greater depth.

Acquired Causes:

1. **Stenosis:** Stenosis is defined as the narrowing or constriction of the puncta or canaliculi, which prevents proper tear flow. Chronic inflammation, scarring, or damage to the lacrimal drainage system can all cause this. Chronic blepharitis, recurrent conjunctivitis, and past ocular operations are all common causes of acquired punctal or canalicular stenosis. Stenosis can cause insufficient tear drainage and tear buildup on the ocular surface, resulting in symptoms such as excessive weeping, discharge, and pain (Dolin and Hecht, 1986).
2. **Inflammation:** Punctal and canalicular blockage can be caused by inflammatory disorders such as dacryocystitis or canaliculitis. Dacryocystitis is defined by inflammation and infection of the lacrimal sac, which is frequently caused by an obstruction in the nasolacrimal duct. If the infection is not treated, it might spread to the canaliculi, resulting in canaliculitis. Inflammation caused by these disorders can cause scarring and fibrosis, resulting in puncta or canaliculi blockage (David Starr Jordan and Stoica, 2020).
3. **Tumors:** Although uncommon, tumors of the lacrimal drainage system can lead to punctal and canalicular blockage. Tumors that originate in the conjunctiva, lacrimal sac, or adjacent tissues might infiltrate the puncta or canaliculi and cause mechanical blockage. When compared to benign tumors, malignant tumors such as squamous cell carcinoma or adenoid cystic carcinoma are more prone to induce blockage. To avoid irreparable damage to the lacrimal system, it is critical to diagnose and treat lacrimal drainage system cancers as soon as possible (Heindl et al., 2010).

4.1 Congenital Causes

Developmental Abnormalities: Congenital lacrimal drainage system defects can cause punctal and canalicular blockage from birth or later in life. These anomalies may include punctal atresia, in which the puncta do not form or are missing, or congenital nasolacrimal duct obstruction (CNLDO), in which the nasolacrimal duct is blocked. CNLDO is a common congenital disorder characterized by severe tearing that, in some cases, resolves spontaneously. Persistent blockage, on the other hand, might lead to problems such as recurring infections or dacryocystitis (Dantas, 2010).

Obstruction due to Foreign Bodies or Anatomical Variations (Bourkiza and Lee, 2012):

1. **Foreign Bodies:** Mechanical blockage can be caused by foreign bodies stuck in the puncta or canaliculi. Eyelashes, cosmetic debris, dust, and other microscopic particles are examples of foreign bodies. When a foreign body obstructs the puncta or canaliculi, tear drainage is disturbed, resulting in tearing, pain, and the possibility of infection. To restore normal tear flow and avoid difficulties, the foreign body must be removed as soon as possible.
2. **Anatomical Variations:** Punctal and canalicular blockage can also be caused by variations in the architecture of the lacrimal drainage system. Anatomical problems such as unusually thin canaliculi, improper puncta placement, or an irregular canaliculi course can all obstruct appropriate tear drainage. These differences may be present from birth or emerge over time as a result of age-related changes or other causes.

Understanding the etiology of punctal and canalicular blockage is critical for appropriate therapy. A thorough study of the patient's medical history, clinical examination, and maybe imaging investigations such as lacrimal syringing or dacryocystography are required for proper diagnosis. Treatment options are determined on the underlying cause and the degree of the blockage [11].

For acquired reasons, conservative measures such as warm compresses, eyelid cleaning, and topical anti-inflammatory medicines may be tried first. Procedures such as punctal dilation or canalicular intubation may be used to restore normal tear outflow in situations of stenosis or scarring. Inflammatory disorders frequently need specific antibiotic or anti-inflammatory medication, as well as steps to address the underlying source of the inflammation. In situations of persistent dacryocystitis or substantial scarring, surgical intervention, such as a dacryocystorhinostomy (DCR), may be required [12].

Most congenital cases of CNLDO resolve spontaneously within the first year of life. Consistent blockage, on the other hand, can be handled with conservative treatments such as massage, topical antibiotics, and monitoring. When conservative treatment fails or difficulties

emerge, surgical intervention may be necessary. To create appropriate tear drainage, procedures such as probing and irrigation, balloon dacryoplasty, or DCR might be explored [13].

Foreign body obstructions must be removed as soon as possible to restore normal tear flow and avoid difficulties. Typically, foreign body removal is accomplished with irrigation or mild probing. Referral to an ophthalmologist may be required in situations with complicated or affected foreign bodies.

Anatomical differences that lead to blockage may necessitate surgical treatment in order to restore normal tear drainage. Canaliculoplasty, punctoplasty, or canalicular reconstruction may be used to relieve the blockage and increase tear flow (Dantas, 2010).

Finally, punctal and canalicular blockage can occur owing to a variety of acquired and congenital reasons, as well as foreign substances or anatomical differences. Understanding and recognizing these etiological elements is critical for successful treatment. Depending on the underlying reason and degree of the blockage, treatment techniques range from conservative to surgical treatments. Individuals affected with punctal and canalicular blockage can benefit from prompt and precise diagnosis, as well as appropriate therapies, to help reduce symptoms, prevent complications, and enhance overall quality of life.

5. DIAGNOSTIC TECHNIQUES FOR PUNCTA AND CANALICULAR OBSTRUCTION

Accurate diagnosis of puncta and canalicular obstruction is essential for effective management and treatment planning. Various diagnostic techniques are employed to evaluate the lacrimal drainage system and identify the underlying cause of obstruction. This unit will explore the diagnostic methods used in the assessment of puncta and canalicular obstruction, including clinical evaluation, patient history, imaging techniques, and tear drainage tests.

5.1 Clinical Evaluation and Patient History

In examining individuals with suspected puncta and canalicular blockage, a comprehensive clinical assessment is essential. The ophthalmologist or oculoplastic surgeon will

conduct a thorough examination that may include the following elements [14].

- a. *External examination:* The lacrimal drainage system's outward appearance is evaluated for evidence of inflammation, edema, or discharge.
- b. *Lid evaluation:* The lids are examined for symptoms of ectropion, entropion, or any other lid malposition that may lead to lacrimal drainage failure.
- c. *Punctal examination:* The puncta's patency, location, and integrity are evaluated. Probing may be used to assess punctal stenosis or blockage.
- d. *Canaliculus evaluation:* The canaliculi are checked for stenosis, blockage, or lacrimal plugs.
- e. *Tear meniscus evaluation:* Abnormalities in the height and quality of the tear meniscus may suggest a clog in the drainage system.

A complete patient history is required in addition to the clinical examination. The ophthalmologist will ask about symptoms including epiphora (excessive tearing), recurring infections, and a history eye trauma or prior surgery. Understanding the length, intensity, and symptoms associated with the blockage might give useful insights into the underlying cause.

5.2 Imaging Techniques

a. *Dacryocystography (DCG):* DCG is a radiographic imaging procedure in which a contrast agent is injected into the lacrimal drainage system. The anatomy is then seen with X-ray scans to discover any obstructions or anomalies. DCG can offer precise information regarding the position and amount of the blockage, which can help with treatment planning.

b. *Lacrimal Scintigraphy:* A radioactive tracer is injected into the lacrimal drainage system during lacrimal scintigraphy. Gamma cameras are used to follow the tracer's movement, allowing the study of tear drainage dynamics. This approach can aid in determining the location and severity of a blockage, assessing tear flow rates, and distinguishing between functional and anatomical obstacles [15].

5.3 Tear Drainage Tests

a. *Jones Test:* The Jones test is a straightforward and widely used diagnostic method for assessing

lacrimal drainage. It entails injecting a drop of fluorescein dye into the conjunctival sac. The patient is then encouraged to softly blow their nose. The presence of the dye in the nasal cavity or nasopharynx implies a patent lacrimal drainage system. The absence of color indicates a barrier in the drainage channel [16].

b. *Fluorescein Dye Disappearance Test*: The rate at which fluorescein dye fades from the tear film is measured in this test to determine tear clearing and drainage. A particular quantity of fluorescein dye is administered to the patient, and the time it takes for the dye to dissipate is recorded. Delayed dye fading indicates obstructed tear drainage or other underlying problems [17].

Clinical evaluation, patient history, imaging techniques such as dacryocystography and lacrimal scintigraphy, and tear drainage tests such as the Jones test and fluorescein dye disappearance test all provide valuable information for the accurate diagnosis of puncta and canalicular obstruction. Combining these methods assists in determining the location, severity, and cause of the blockage, as well as in developing an effective treatment strategy. Each diagnostic approach provides distinct insights into the underlying pathophysiology, allowing for a thorough evaluation of the lacrimal drainage system [18].

The clinical assessment and patient history give critical preliminary information regarding the patient's symptoms, duration of symptoms, and any pertinent circumstances such as trauma or past procedures. The ophthalmologist can discover any evident indicators of blockage or abnormalities by closely inspecting the exterior appearance, lids, puncta, canaliculi, and tear meniscus. The evaluation of lid malposition, such as ectropion or entropion, aids in determining whether or not lid abnormalities contribute to the blockage [19].

Imaging methods are essential for further investigating and visualizing the lacrimal drainage system. A contrast agent is used in dacryocystography (DCG) to delineate the anatomy of the lacrimal drainage system. This method generates comprehensive radiographic pictures that can pinpoint the exact position, extent, and kind of blockage. DCG is very helpful in detecting strictures, fistulas, and other anatomical anomalies [20].

In contrast, lacrimal scintigraphy involves the injection of a radioactive tracer into the lacrimal

drainage system. This approach enables dynamic imaging of tear flow, revealing tear drainage rates and patterns. Lacrimal scintigraphy can distinguish between functional and anatomical blockages, assisting in the identification of the underlying source of the obstruction [21].

Tear drainage tests are non-invasive methods used to assess the lacrimal drainage system's patency and function. The Jones test, which employs fluorescein dye, determines the drainage pathway's continuity. The presence or absence of blockage can be detected by monitoring whether the dye emerges in the nasal cavity or nasopharynx when the patient blows their nose. This test is useful for finding common blockage locations, such as the canaliculi or the nasolacrimal duct [16].

The fluorescein dye disappearance test determines how quickly fluorescein dye vanishes from the tear film. The tear clearing and drainage efficiency are assessed in this test. A prolonged disappearance of the dye indicates inadequate tear drainage, which might be caused by puncta or canalicular occlusion [17].

Ophthalmologists can acquire a full examination of the lacrimal drainage system by combining various diagnostic procedures. The combination of clinical evaluation, patient history, imaging techniques such as dacryocystography and lacrimal scintigraphy, and tear drainage tests such as the Jones test and fluorescein dye disappearance test improves diagnostic accuracy and aids in the selection of the best treatment approach.

Finally, correct identification of puncta and canalicular blockage is critical for effective therapy. Clinical examination, patient history, imaging methods, and tear drainage tests are all important in evaluating the lacrimal drainage system and determining the underlying reason of blockage. Ophthalmologists can establish an accurate diagnosis, facilitate successful treatment planning, and improve patient outcomes by applying these diagnostic tools in a complete and integrated way.

6. TREATMENT OPTIONS FOR PUNCTA AND CANALICULAR OBSTRUCTION

Treatment approaches for puncta and canalicular blockage seek to restore normal lacrimal drainage system function, relieve symptoms, and

enhance patient quality of life. Treatment is determined on the underlying etiology, the degree of the blockage, and individual patient variables. To obtain the best results, many times a mix of non-surgical and surgical procedures are used.

For mild to severe instances of puncta and canalicular blockage, non-surgical procedures are the initial line of treatment. Conservative treatments include warm compresses, eyelid cleanliness, and topical lubricants. These therapies serve to alleviate symptoms and improve tear drainage by reducing inflammation, clearing debris, and promoting tear film stability. Non-surgical techniques are frequently regarded as first-line treatments, particularly in situations of partial blockages or functional problems [22].

Another non-surgical strategy used to treat obstructive disorders is lacrimal drainage system dilation and irrigation. To enlarge the narrow or stenotic passageways, lacrimal probes or dilators are inserted into the puncta and canaliculi. Tear flow is restored by dilating the blocked segments, allowing for better drainage. Following dilatation, sterile saline solution irrigation may be used to clean out any collected debris or mucus. This treatment may be performed in the office setting under local anesthetic with little pain.

When non-surgical procedures fail to provide satisfactory results or the blockage is severe, surgical measures may be required. Dacryocystorhinostomy (DCR) is a surgical technique that is widely used to treat puncta and canalicular blockage. It entails constructing a new drainage conduit by circumventing the partially clogged lacrimal drainage system. During DCR, a link is formed between the lacrimal sac and the nasal cavity, allowing tears to flow directly into the nasal cavity and avoid the blockage. DCR can be done externally or endoscopically, depending on anatomical concerns and the surgeon's inclination [23].

Another surgical technique for treating canalicular blockages is canalicular stenting. A silicone or polyurethane stent is inserted into the afflicted canaliculus to preserve patency and promote normal tear drainage. The stent acts as a scaffold, keeping the canaliculus open and enabling tears to flow freely. Canalicular stenting is very beneficial when there are canalicular strictures or recurring blockages [24].

Endoscopic approaches for the treatment of puncta and canalicular blockage have grown in

favor in recent years. Endoscopic dacryocystorhinostomy (endo-DCR) is a procedure that involves the use of specialized equipment and an endoscope to establish a new drainage channel without the need for exterior incisions. This minimally invasive method has advantages such as less scarring, faster healing, and better cosmesis. Endoscopic laser dacryocystorhinostomy (endo-L-DCR) combines the benefits of endoscopy with laser energy to allow for precise tissue ablation and promote healing. Hanife Tuba Akcam and Onur Konuk, [25].

The most appropriate treatment strategy is determined by a number of criteria, including the location and degree of the obstruction, the patient's general health, and the surgeon's skill. To select the best care strategy for each patient, a comprehensive examination and tailored approach are required.

Clinical assessment and patient history are critical during the diagnostic phase in determining the existence of puncta and canalicular blockage. However, without the use of imaging methods and tear drainage tests, the diagnosis procedure is inadequate. These approaches give useful information on the location and degree of the blockage, assisting in the selection of the best treatment approach.

Dacryocystography and lacrimal scintigraphy are typical imaging procedures used to study the architecture and function of the lacrimal drainage system. Dacryocystography is injecting a contrast agent into the lacrimal system, followed by imaging to see how the dye flows along the drainage routes. This approach aids in determining the location and nature of the blockage, allowing for more targeted treatment planning. Lacrimal scintigraphy, on the other hand, evaluates tear outflow and identifies any abnormalities in the lacrimal system using a radioactive tracer.

Tear drainage tests, such as the Jones test and fluorescein dye disappearance test, give important information regarding the lacrimal drainage system's patency and performance. The Jones test is inserting a saline solution into the conjunctival sac and evaluating the appearance of fluid at the nasal aperture of the lacrimal system. A positive Jones test implies that there is a functional blockage, whereas a negative test indicates that there is a mechanical obstruction. The fluorescein dye disappearance

test includes injecting fluorescein dye into the conjunctival sac and monitoring its disappearance over time. The delayed disappearance of dye implies a blockage in the lacrimal drainage system [26].

Once a puncta and canalicular occlusion diagnosis has been established, relevant treatment options can be considered. Non-surgical therapies are frequently considered first, particularly in situations with mild to moderate blockages. Conservative treatment, such as warm compresses and eyelid cleanliness, reduces inflammation and clears debris from the lacrimal system, which helps ease symptoms and enhance tear outflow. Topical lubricants are also used to improve tear film stability and relieve dryness.

Lacrimal drainage system dilation and irrigation is a non-surgical technique done under local anesthetic. Lacrimal probes or dilators are inserted into the puncta and canaliculi to enlarge the tight passageways and increase tear flow. To drain out any collected debris or mucus, irrigate with sterile saline solution at the same time.

If non-surgical procedures do not produce satisfactory outcomes or if the blockage is severe, surgical measures may be required. Dacryocystorhinostomy (DCR) is a common surgical technique that includes establishing a new tear drainage channel. An external method, in which an incision is made on the skin around the medial canthus, or an endoscopic technique, in which specialized instruments and an endoscope are used to reach the nasal cavity and form a link between the lacrimal sac and the nasal mucosa, can be utilized to accomplish this. DCR has a high success rate in symptom relief and restoring tear drainage [27].

Canalicular stenting is another surgical treatment option for canalicular blockages. It entails inserting a silicone or polyurethane stent into the afflicted canaliculus to preserve patency and adequate tear drainage. The stent works as a scaffold, preventing collapse of the canaliculus and enabling tears to flow freely.

Endoscopic procedures have transformed the management of puncta and canalicular blockage. Endoscopic dacryocystorhinostomy (endo-DCR) is a minimally invasive treatment that involves inserting an endoscope and specialized equipment into the nose to construct a new drainage conduit without the need for exterior

incisions. This method has various advantages, including less scarring, faster healing, and better aesthetic effects. Endoscopic procedures, which allow for precise imaging and manipulation of the lacrimal system, are especially advantageous in patients who have already undergone nasal or sinus surgery.

Endoscopic laser dacryocystorhinostomy (endo-L-DCR) combines the advantages of endoscopy with the use of laser energy to enhance tissue ablation and improve healing. The laser is utilized to construct a bone ostium and remove any blocking tissues, allowing a functional drainage channel to be established. Endo-L-DCR has a high success rate and can be especially effective in circumstances when standard surgical procedures have failed or are contraindicated [28].

The surgical intervention chosen is determined by a number of criteria, including the location and degree of the blockage, the patient's features, the surgeon's expertise, and the available resources. The purpose of surgery is to restore normal lacrimal drainage system function, increase tear clearing, and relieve symptoms including epiphora (excessive weeping) and recurring infections [29].

To get the greatest results, a mix of surgical and non-surgical therapies may be required in some circumstances. To treat both the mechanical obstruction and the underlying anatomical defects, a patient with significant canalicular obstruction may have canalicular stenting followed by endoscopic dacryocystorhinostomy [30].

Postoperative care is critical for ensuring positive results and minimizing complications. Patients are often offered antibiotics to avoid infection and encouraged to do frequent saline irrigation to keep the lacrimal system functioning properly. Follow-up appointments are planned to monitor the healing process, evaluate tear drainage, and address any concerns or issues that may occur [31].

It is crucial to note that treatment success rates for puncta and canalicular blockage differ depending on the underlying reason and degree of the obstruction. Some individuals may have total symptom relief and restoration of normal tear drainage, whilst others may only have partial recovery or require further procedures. To provide comprehensive and tailored care for

patients with puncta and canalicular blockage, ophthalmologists, otolaryngologists, and other healthcare specialists must work closely together.

Finally, the therapeutic options for puncta and canalicular occlusion include both non-surgical and surgical treatments. Conservative management, dilation, and irrigation are frequently used as first-line treatments. If these methods fail or the blockage is severe, surgical therapies such as dacryocystorhinostomy, canalicular stenting, endoscopic techniques, or laser-assisted operations may be required. The treatment method used is determined on the unique features of the blockage, patient circumstances, and surgeon ability. To maximize results and enhance patients' quality of life, puncta and canalicular blockage necessitate a multidisciplinary approach, tailored treatment, and long-term follow-up.

7. MANAGEMENT OF SPECIFIC ETIOLOGICAL FACTORS

Puncta and canalicular blockage can be caused by a variety of underlying etiological reasons, and treatment frequently depends on addressing the specific cause. In this part, we will look at some of the common etiological variables related with puncta and canalicular blockage.

7.1 Treating Inflammation and Infections

Inflammation and infections can both play a role in the formation of puncta and canalicular blockage. When the blockage is caused by an inflammatory disease, such as chronic dacryocystitis or canaliculitis, the primary goal of therapy is to reduce the inflammation and manage the infection.

To target the pathogenic organisms, treatment usually consists of a mix of topical and systemic medicines. Surgical intervention may be required in some situations to drain abscesses or remove granulation tissue that is impeding the drainage channel. Treatment of the underlying inflammation and infection is critical to preventing recurring blockage and maintaining the function of the lacrimal drainage system.

7.2 Addressing Neoplastic Conditions

Neoplastic diseases, such as lacrimal sac or canalicular tumors, can restrict the lacrimal drainage channel mechanically. These situations

require a multidisciplinary approach, which includes consultation with oncologists and ophthalmic surgeons.

The treatment strategy is determined on the kind, location, and stage of the tumor. Surgical excision is frequently used as the primary therapeutic strategy, with the goal of removing the tumor and restoring normal anatomical structure. Adjuvant treatments such as radiation or chemotherapy may be needed in some circumstances to guarantee total tumor cell elimination.

In patients with neoplastic-related obstruction, regular follow-up and surveillance are required to monitor for recurrence or metastasis. Prompt identification and treatment of neoplastic diseases are critical for improving results and reducing the risk of complications.

7.3 Correcting Anatomical Abnormalities

Puncta and canalicular blockage can be caused by anatomical abnormalities such as nasolacrimal duct strictures or congenital defects. Surgical correction is frequently used to restore the normal architecture and function of the lacrimal drainage system in these circumstances.

Depending on the anatomical anomaly, surgical therapies may include dacryoplasty, dacryocystorhinostomy, or canalicular repair. The purpose of surgery is to construct or repair a patent drainage channel that allows tears to flow freely.

The effectiveness of anatomical abnormality surgical treatment is determined by various factors, including the severity of the blockage, the existence of accompanying comorbidities, and the surgical method used. To guarantee maximum healing and functional results, close postoperative monitoring and adequate postoperative care are required.

7.4 Complications and Prognosis

Untreated or improperly managed puncta and canalicular blockage can result in a number of problems. Chronic tearing (epiphora), recurring infections, and even the creation of a lacrimal sac abscess might ensue from persistent blockage. Chronic tear buildup in the lacrimal sac can also result in dacryocystitis, a painful infection of the lacrimal sac that need rapid medical intervention.

Furthermore, long-term occlusion can cause irreparable damage to the lacrimal drainage system, resulting in chronic functional impairment and limited tear clearing if left untreated. These consequences can have a substantial influence on a patient's quality of life, causing pain, vision abnormalities, and social shame.

The severity and length of the blockage, the success of therapy, and the existence of any accompanying ocular or systemic disorders all influence the prognosis of puncta and canalicular obstruction. Timely intervention and effective therapy enhance the prognosis greatly, with a better possibility of symptom remission and restoration of normal lacrimal drainage function.

The majority of individuals with puncta and canalicular blockage can have their symptoms resolved and their tear drainage restored if they get prompt and appropriate treatment. In circumstances when the blockage is slight or transient, non-surgical therapies like as conservative management, dilatation, and irrigation can give relief. These conservative therapies try to relieve symptoms and enhance natural tear clearing via the lacrimal system.

When conservative methods fail or the blockage is chronic and severe, surgical intervention may be required. Dacryocystorhinostomy (DCR) and canalicular stenting are popular procedures used to develop alternate tear drainage paths or to bypass the clogged part of the lacrimal system. DCR entails constructing a new drainage conduit between the lacrimal sac and the nasal cavity, allowing tears to flow through the blockage. Canalicular stenting is inserting tubes or stents into the afflicted canaliculus to keep it open and allow tears to drain.

Endoscopic and laser-assisted procedures have improved the treatment options for puncta and canalicular blockage. Endoscopic DCR, which uses endoscopic guidance and specialized equipment, is a less invasive method with success rates equivalent to classic open DCR. Endonasal laser dacryocystorhinostomy, for example, uses laser energy to establish the new drainage conduit, minimizing the need for external incisions and perhaps leading to speedier recuperation.

Complications related to the treatment of puncta and canalicular blockage are uncommon but can arise. Infection, hemorrhage, scarring, and stent-related complications are all possibilities. Close

monitoring and adequate postoperative care can assist to reduce problems and provide the best possible outcomes.

To summarize, the treatment of puncta and canalicular occlusion requires a personalized strategy depending on the underlying cause. The treatment of inflammation and infections, the treatment of neoplastic disorders, and the correction of anatomical defects are all important factors in the management process. Timely intervention, whether non-surgical or surgical, is critical in reducing complications and improving the prognosis. Healthcare practitioners can successfully relieve symptoms, restore normal tear drainage, and improve the quality of life for patients with puncta and canalicular blockage by addressing particular etiological reasons and adopting the most appropriate therapy methods.

8. CONCLUSION

Puncta and canalicular blockage is a multifaceted illness that necessitates a thorough study of its origin, diagnostic procedures, and treatment choices. This study work has offered a thorough examination of the numerous factors related with puncta and canalicular blockage, with the goal of improving knowledge and management of this problem.

Inflammation, infections, neoplastic diseases, and anatomical anomalies were found as prevalent etiological causes in this investigation. These variables can contribute to puncta and canalicular system occlusion, resulting in poor tear drainage. When evaluating and managing patients with puncta and canalicular blockage, these criteria must be taken into account.

Diagnostic procedures are critical in determining the severity of this illness. The clinical assessment and patient history give important information about the symptoms and probable underlying causes. Dacryocystography and computed tomography, for example, help in seeing the lacrimal drainage system and determining the location and extent of the obstruction. Tear drainage tests, such as the Jones I and II tests, aid in determining the system's functional integrity.

Treatment for puncta and canalicular blockage is determined on the underlying etiology and severity of the problem. Specific etiological variables are managed by treating inflammation and infections with medical therapy such as

topical antibiotics and anti-inflammatory medicines. Neoplastic diseases may necessitate surgical excision or radiation treatment. Surgical procedures, such as dacryocystorhinostomy or canalicular repair, can be used to rectify anatomical defects.

Complications linked with puncta and canalicular blockage should be examined since they can affect a patient's prognosis. Recurrent infections, lacrimal sac abscess, and dacryocystitis are examples of consequences. Early detection and control of problems are critical for attaining positive outcomes.

In conclusion, this complete discussion of the etiology, diagnostic procedures, and treatment choices for puncta and canalicular blockage has offered a thorough assessment. By better understanding this illness, healthcare providers may make more educated decisions about patient care, resulting in better outcomes and quality of life for people suffering with puncta and canalicular blockage.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. Mueller AJ, Scofield-Kaplan SM, Tenzel P. a, Winward B, Clair BC, Mancini R. Endoscopic dacryocystorhinostomy as an option for symptomatic epiphora in isolated monocanalicular obstruction. *Orbit*. 2020; 40(5):419–422. DOI:https://doi.org/10.1080/01676830.2020.1799420.
2. Fiorino MG, Quaranta-Leoni C. Quaranta-Leoni FM. Proximal lacrimal obstructions: a review. *Acta Ophthalmologica*. 2021;99(7):701–711. DOI:https://doi.org/10.1111/aos.14762.
3. Tabbara KF, Bobb AA. Lacrimal System Complications In Trachoma. 1980;87(4): 298–301. DOI:https://doi.org/10.1016/s0161-6420(80)35234-2.
4. McNab AA. Lacrimal canalicular obstruction associated with topical ocular medication. 1998;26(3):219–223. DOI:https://doi.org/10.1111/j.1442-9071.1998.tb01315.x.
5. Quaranta-Leoni FM, Fiorino MG, Serricchio F, Quaranta-Leoni F. Management of proximal lacrimal obstructions: a rationale. *Acta Ophthalmologica*. 2020;99(4). DOI:https://doi.org/10.1111/aos.14632.
6. Bohman E, Kugelberg M, Dafgård Kopp E. Long-term outcome of lacrimal stent intubation for complete acquired lacrimal drainage obstructions. *Acta Ophthalmologica*. 2019;98(4):396–399. DOI:https://doi.org/10.1111/aos.14263.
7. Yulish M, Pikkell J. Presumed Virus-Induced Punctal Occlusion; 2014. DOI:https://doi.org/10.1155/2014/809851.
8. Corak F. (2022). The relationship between histopathological features and surgical success in primary acquired punctal stenosis - Fatma Corak Eroglu, Irem Inanc, Bizden Sabuncuoglu, Emine Sen, Burcu Kazanci, [online] *European Journal of Ophthalmology*; 2022. Available:https://journals.sagepub.com/doi/abs/10.1177/11206721211033479?journalCode=ejoa [Accessed 1 Jun. 2023].
9. Briscoe D, Edelstein E, I. Zacharopoulos, Yoram Keness, Kilman A, Zur F, Assia EI. Actinomyces canaliculitis: diagnosis of a masquerading disease. 2004;242(8):682–686. DOI:https://doi.org/10.1007/s00417-004-0871-5.
10. Lin CH, Wang CY, Shen YC, Wei LC. Clinical Characteristics, Intraoperative Findings, and Surgical Outcomes of Canalicular Laceration Repair with Monocanalicular Stent in Asia. 2019;1–6. DOI:https://doi.org/10.1155/2019/5872485.
11. Enright NJ, Brown S, Rouse H, McNab AA, Hardy T. Nasolacrimal Sac Diverticulum: A Case Series and Literature Review; 2019. DOI:https://doi.org/10.1097/iop.0000000000001156.
12. Qazi Y, Aggarwal S, Hamrah P. Image-guided evaluation and monitoring of treatment response in patients with dry eye disease. 2014;252(6):857–872. DOI:https://doi.org/10.1007/s00417-014-2618-2.

13. Nakayama T, Watanabe A, Rajak S, Yamanaka Y, Chie Sotozono Congenital nasolacrimal duct obstruction continues trend for spontaneous resolution beyond first year of life. 2020; 104(8):1161–1163. DOI:<https://doi.org/10.1136/bjophthalmol-2019-314980>.
14. Ozlen Rodop Ozgur, Levent Akçay, Tutas, N. and Onur Karadağ Management of acquired punctal stenosis with perforated punctal plugs. 2015;29(3):205–209. DOI:<https://doi.org/10.1016/j.sjopt.2015.04.001>.
15. Seyed Ali Mirshahvalad, Mohammadreza Chavoshi, Mohsen Bahmani Kashkouli, Fallahi, B., Alireza Emami-Ardakani and Reyhaneh Manafi-Farid Diagnostic value of lacrimal scintigraphy in the evaluation of lacrimal drainage system obstruction: A systematic review and meta-analysis. [online] Publish Ahead of Print; 2022. DOI:<https://doi.org/10.1097/mnm.0000000000001578>.
16. Dutton JJ. Evaluation and Imaging of Lacrimal Drainage Disorders. 2019;433–448. DOI:https://doi.org/10.1007/978-3-030-14092-2_38.
17. Mohsen Bahmani Kashkouli, Hoor Mirzajani, Mansooreh Jamshidian-Tehrani, Sahab Shahrzad and Mostafa Soltan Sanjari Fluorescein Dye Disappearance Test. 2015;31(4): pp.296–299. DOI:<https://doi.org/10.1097/iop.0000000000000308>.
18. Garaszczuk IK, Montes Mico R, Iskander DR, Expósito AC. The tear turnover and tear clearance tests – a review. Expert Review of Medical Devices. 2018;15(3):219–229. DOI:<https://doi.org/10.1080/17434440.2018.1435271>.
19. Goel R, Jain S, Malik K, Nagpal S, AG A, Kumar S, Kishore D. Oculoplasty for general ophthalmologists. Expert Review of Ophthalmology. 2015;10(2):197–210. DOI:<https://doi.org/10.1586/17469899.2015.1007129>.
20. Barna S, Garai I, Kornel Kukuts, Gesztelyi R, Toth LS, Kemeny-Beke A. Clinical utility of SPECT/CT and CT-dacryocystography-enhanced dacryoscintigraphy in the imaging of lacrimal drainage system obstruction. 2019;33(10):746–754. DOI:<https://doi.org/10.1007/s12149-019-01385-2>.
21. Isloor SD. Lacrimal Drainage Surgery. [online] Google Books. JP Medical Ltd; 2014. Available:https://books.google.gr/books?hl=el&lr=&id=IK7TAwAAQBAJ&oi=fnd&pg=PA17&dq=lacrimal+scintigraphy+involves+the+injection+of+a+radioactive+tracer+into+the+lacrimal+drainage+system&ots=6D6Lyx-jO7&sig=la2kz8lcTsmq1CiVDW_aSoYWB7o&redir_esc=y#v=onepage&q&f=false [Accessed 31 May 2023].
22. Badawy AM. Surgical Versus non Surgical Techniques for Punctal Occlusion as Alternative Methods for Treatment of Dry-eye in Dogs. Journal of Advanced Veterinary Research. 2017; 7(4):108–115. Available:<https://www.advetresearch.com/index.php/avr/article/view/272> [Accessed 31 May 2023].
23. Vagge A, Desideri L, Nucci P, Serafino M, Giannaccare G, Lembo A, Carlo Enrico Traverso Congenital Nasolacrimal Duct Obstruction (CNLDO): A Review. 2018; 6(4):96–96. DOI:<https://doi.org/10.3390/diseases6040096>.
24. Mansur C, Pfeiffer ML, Bitá Esmaeli. Evaluation and Management of Chemotherapy-Induced Epiphora, Punctal and Canalicular Stenosis, and Nasolacrimal Duct Obstruction. 2017;33(1):9–12. DOI:<https://doi.org/10.1097/iop.0000000000000745>.
25. Hanife Tuba Akcam and Onur Konuk. Mechanical transnasal endoscopic dacryocystorhinostomy versus transcanalicular multidiode laser dacryocystorhinostomy: long-term results of a prospective study. 2021;36(2):349–356. DOI:<https://doi.org/10.1007/s10103-020-03038-7>.
26. Singh S, Ali M. A Review of Diagnostic and Therapeutic Dacryoendoscopy. 2019;35(6):519–524. DOI:<https://doi.org/10.1097/iop.0000000000001404>.
27. Mimura M, Ueki M, Oku H, Sato B, Ikeda T. Indications for and effects of Nunchaku-style silicone tube intubation for primary acquired lacrimal drainage obstruction. 2015;59(4):266–272. DOI:<https://doi.org/10.1007/s10384-015-0381-5>.

28. Bidhan Chandra Ray, Datta, D. and Pijush Pal Roy. Long Term Success of Endo-DCR with Silicone Tube Catheter (STC) versus Endo-DCR without STC: A Comparative Study. 2016;24(2):88–93. DOI:<https://doi.org/10.47210/bjohns.2016.v24i2.83>.
29. Cena I, Matteo Alicandri-Ciufelli and Federico Maria Gioacchini. Epiphora and Indications to Surgery. 2016;45–52. DOI:https://doi.org/10.1007/978-3-319-20633-2_5.
30. Penttilä E, Smirnov G, Tuomilehto H, Kaarniranta K, Seppä J. Endoscopic dacryocystorhinostomy as treatment for lower lacrimal pathway obstructions in adults: Review article. Allergy & Rhinology. 2015;6(1):12–19. DOI:<https://doi.org/10.2500/ar.2015.6.0116>
31. Azhar RA, Bochner BH, James WF, Catto, Goh A, Kelly JM, Patel HD, Pruthi RS, Thalmann GN, Desai MM. Enhanced recovery after urological surgery: A contemporary systematic review of outcomes, key elements, and research needs. 2016;70(1): 176–187. DOI:<https://doi.org/10.1016/j.eururo.2016.02.051>.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/115075>