



Zinc Nutrition in Rice Based Cropping Systems: A Comprehensive Review

**Boddu Archana ^{a*}, Bindhu J.S ^b, Shalini Pillai P ^a,
Jacob John ^b and Meera A.V ^b**

^a Department of Agronomy, College of Agriculture, Kerala Agricultural University, Vellayani, Pin-695522, India.

^b Integrated Farming System Research Station, Kerala Agricultural University, Karamana, Thiruvananthapuram, Pin-695002, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JEAI/2024/v46i52413

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/114940>

Review Article

Received: 28/01/2024

Accepted: 02/04/2024

Published: 06/04/2024

ABSTRACT

Rice-based cropping systems being the most practiced cropping system in India, it suffers zinc deficiency as majority of the Indian soils are deficient in the Zn micronutrient. Zn is considered the fourth most important yield limiting nutrient of Agricultural crops. Apart from plant nutrition Zn also plays crucial role in human nutrition. Zn plays a vital role in the physiological function of plants and it is associated with phytohormone activity, protein synthesis and carbohydrate metabolism. To understand better zinc nutrition and its associated benefits from optimum nutrition and ways to manage Zn in rice-based cropping systems, the present review critically examines nutrient mobilization and transformation of Zn in rice-based cropping systems of India.

Keywords: *Rice-based cropping systems; phytohormone; protein synthesis; carbohydrate metabolism; optimum nutrition.*

*Corresponding author: E-mail: boddu-2020-21-046@student.kau.in;

J. Exp. Agric. Int., vol. 46, no. 5, pp. 574-583, 2024

1. INTRODUCTION

Zinc is considered the key element affecting the growth, development and life of living organisms along with playing an essential role in physiological activities [1,2,3]. Zn deficiency in people results in symptoms such as loss of appetite, anorexia, loss of taste and smell, and other symptoms. It can also have an impact on the immune system, leading to arteriosclerosis and anemia [4]. Growth retardation, congenital fetal abnormalities, and pregnancy and birthing problems have all been linked to zinc deficiency [5].

Micronutrient deficits, particularly those of Zn, are one of the main causes of economic loss in developing nations [6]. According to recommendations, an average individual needs 11 mg of zinc per day for males and 8 mg of zinc per day for females [7].

Zinc is a crucial element needed to improve crop growth and output; of all micronutrients, Zn is regarded as the most important nutrient [8]. More than 300 enzymes, including carbonic anhydrase, alcoholic dehydrogenase, alkaline phosphate, carboxy peptidase, phospholipase, superoxide dismutase (SOD) and RNA polymerase depend on zinc for their structural and functional integrity. These enzymes regulate a variety of biochemical and physiological processes including cell division, protein synthesis, photosynthesis, gene transcription, nucleic acid metabolism and pollen development [9].

Rice is a major staple food cereal on which half the world's population is dependent for energy and 20% protein requirements [10]. Around 50%

of Asia's population, or roughly 2 billion people, are in danger of Zn insufficiency, though Asia covers about 87% of the world's rice-growing area and consumes 90% of all rice produced [11,12]. Agronomic biofortification of Zn could be a potential option to address the problem of Zn deficiency [13].

The primary cropping system used in India is the rice-based cropping system (RBCS), which rotates crops with vegetables, fodder crops, pulses and oilseeds. The productivity of a system is significantly influenced by how nutrients are managed during different seasons. Any farming system can be made more productive by using micronutrients [14]. Zn is thought to be the micronutrient that is needed the most.

2. RICE-BASED CROPPING SYSTEMS (RBCS)

Rice is the predominant crop in the world meeting 20% of the human food calories and India is considered as the largest producer occupying ~ 45 million ha under cultivation [14]. Due to varying climatic situations during the crop year the annual crops are grown in progression. The crops may either be grown along with rice as inter-crop or in sequence according to the climatic conditions and feasibility to maximize resource utilization [15].

Rice-based cropping system can be described as, in a farming practice considering rice as the major crop is succeeded by other adaptable crops. In India, rice-based cropping system is the major cropping system where, the major crop is rice followed by pulses, cereals, oilseeds, etc. [14].

Table 1. Rice based cropping systems followed in Indian states [14]

Agro-climatic region	Cropping system followed
Western himalayan	Rice-wheat, rice-potato-potato
Eastern himalayan	Rice fallow, rice-rice, rice-pulses/oilseeds
Lower gangetic plain	Rice-rice, rice-wheat, rice-potato-jute/vegetables
Middle gangetic plain	Rice-wheat, rice-maize, rice-potato-sunflower
Upper gangetic plain	Rice-wheat, sugarcane-ratoon-wheat
Trans gangetic plain	Rice-wheat
Eastern plateau and hills	Rice-blackgram, rice-niger/linseed, rice-vegetables
East coast plain and hills	Rice-groundnut-greengram, Rice-greengram/blackgram, rice-rice
West coast plains and hills	Rice-rice
Andaman and Nicobar	Rice- fallow

The central principle in considering a crop in a cropping system lies with; a. the biological traits of the crop and how they interact with and are influenced by its physical, chemical, and ecological environment, b. the crop to be grown in sequence after the major crop, c. the management practices to be followed for the cropping system [16]. The major crop in cropping systems was chosen based on the agro-climatic and socio-economic circumstances of the area. In India, rice-rice, rice-wheat, rice-pulse, and rice-potato are the most common rice-based cropping systems [17].

3. NUTRIENT MOBILIZATION OF ZINC IN RBCS

The movement of the nutrient Zn in the crop sequence of the cropping system can be understood as nutrient mobilization. Cropping systems following the integrated mode of nutrient management were found to have a significant effect on soil fertility. Inclusion of pulse crops like green gram and vegetable crops in the rice-based cropping system along with supplementing nutrients through different sources of biofertilizers and organic manures significantly improves crop nutrient absorption by increasing the availability of soil nutrients, the total zinc nutrient uptake by the rice and wheat [18].

The organic sources of nutrients and biofertilizers supply considerable amounts of the major and micronutrients continuously throughout the crop growth period unlike the inorganic sources, the nutrients are available to the plants in the earlier and the later stages of growth, organic sources provide long-term availability of nutrients [19,20,21].

Change from the current, conventional cropping system to an enhanced cropping system including potato and cucumber between two seasons of rice has resulted in 49% higher rice equivalent yield due to the greater profits gained under higher yield of cucumber and potato [22]. Rice-non rice crop-based cropping systems were more profitable compared to rice-rice cropping systems [23].

In a rice-based cropping system, crop intensification results in a rice equivalent yield that is two to four times greater [24,25]. The rate of increase of REY depends on the type of crop, the number of crops involved, and the ecosystem in which the crop is grown. An improved cropping

system consisting of transplanted Aman rice followed by potato, cucumber and transplanted Aus rice has received 49% higher rice equivalent yield compared to the existing cropping system consisting of transplanted Aman rice followed by potato and transplanted Boro rice [22].

The Nutrient Index (NI) gauges reveals how well soil can supply nutrients to plants [26]. Cropping systems with pulse as a component will have a higher nutrient index than vegetables as a component [27]. Supply of the recommended dose of fertilizers along with the inclusion of pulse crops as a component/sequence in a cropping system will result in the improved nutrient response of the crop to Zn [28]. To achieve higher productivity and profitability from a cropping system the need-based application of Zn is essential along with the application of inorganic sources of fertilizers as the nutrients will be present in the available forms for plants to get maximum benefit [29].


Thus, the application of both organic and inorganic sources of nutrients will make the nutrients more available as is the case with DTPA-extractable Zn content in post-harvest soil of both rice and wheat, Customized fertilizer application had shown increased Zn content by 95% and 158% over 100% RDF in soils of wheat and rice, respectively [30].

4. STATUS OF ZN IN INDIAN SOILS

Fertility of Indian soils is generally poor with progressively emerging micronutrient deficiencies due to their catalyzed removal under agricultural intensification. According to latest estimates, out of about 188.4 thousand tonnes (Tt) of micronutrients removed by 263 Mt of food grains produced, nutrient removal for Zn is 23.9 Tt and in Indian soils, total Zn content ranges from 7-2960 mg/kg and available Zn content ranges from 0.01-52.9 mg/kg. Currently, 36.5% of soil samples nationwide have deficiencies in available Zn, with acute, latent, and total deficiencies occurring in 8, 29, and 15% of the country, respectively [31]. Zn is currently regarded as the fourth-most significant nutrient for crops in India that limits productivity. By 2025, zinc deficiency in Indian soils is predicted to rise from 42 to 63% [32]. According to a study of Indian soils, out of 2.52 lakh soil samples examined, 49% were found to be low in the available Zn spread throughout 20 states. There is widespread zinc deficiency throughout all agro-ecological zones (AEZ). Maximum Zn deficit

extended, up to 60%, in AEZ 6 and 10 soils. Agro-ecological zones 3, 4, 7, 8, 13, and 14 had soils that were 50–60% zinc deficient. In states like Punjab, Haryana, Uttar Pradesh, Andhra Pradesh, Bihar, and Madhya Pradesh, the severity of Zn deficiency is decreasing.

The extent of zinc deficiency was found to be 55, 47, and 36% in Trans-northern, Central, and Eastern sections of IGP alluvial plains, according to analysis of 90419 surface soil samples from AEZ 4, 9, 13, and 16 of the Indo-Gangatic alluvial plains [33].



Map 1. Map showing area having Zinc deficient soils

Table 2. Critical limits of available Zn for agricultural soils of India

	Available Zn (mg kg ⁻¹)
Acute deficient	≤ 0.30
Deficient	> 0.30 – ≤ 0.60
Latent deficient	> 0.60 – ≤ 0.90
Marginal sufficient	> 0.90 – ≤ 1.20
Adequate	> 1.20 – ≤ 1.80
High	> 1.80




Fig. 1. Deficit by state (% of soil samples) available Zn [34].

5. PHYSIOLOGICAL FUNCTIONS OF ZN IN PLANTS

Zinc is the vital nutrient having its role in normal growth and productivity of the plants [35]. The nutrient was also found to have its role in structurally and functionally maintaining the cell membranes [36]. Optimum Zn nutrition helps in the improvement of the photosynthetic activity by 50%-70% and thus increases the chlorophyll content, net photosynthetic rate, transpirational rate, and stomatal conductance [37,38]. Zn application improves plant activity under drought conditions by enhancing the photosynthetic process [39]. A deficiency of zinc in plants results in reduction of chlorophyll content and also affects the ratio of chl a: chl b. Reduced photosynthetic activity leads to reduced growth, leaf area (up to 72%) and reduced number of leaves [40]. Zinc plays role in phytohormone activity, enzyme activation, protein synthesis, carbohydrate metabolism and noticeable role in defending against phytopathogens [41,35]. Zinc is identified as the primary requisite of enzyme classes such as transferases, ligases, oxidoreductases, isomerases, lyases, hydrolases and is also a constituent of ribulose 1,5-biphosphate carboxylase (RuBpc) involved in photosynthesis. As the element is involved in different categories of enzymes the functions, indirectly involved in the metabolism of carbohydrates, proteins, auxins; the deficient conditions affect the productivity of the plant [42,43]. The enzyme carbonic anhydrase which governs the fixation of carbohydrates by conversion of carbon dioxide to bicarbonates is managed by zinc [44]. The most rudimentary effect of zinc in protein metabolism is its activity

in maintaining the stability and functioning of genetic material [38]. Living membranes are maintained by zinc, cell membrane integrity is often regulated by the availability of zinc which drives the detoxification of membrane-damaging ROS and thus involved in the plant water relations [45]. So, in the absence of the Zn sulphhydryl groups and lipids in the membrane undergoes injury and organic compounds leaks through the cells, especially root cells exposed to certain situations leads to root diseases in Zn deficient plants [46,47]. The defense mechanism of plants against the diseases is often correlated to the Zn nutrition through which the plants attain resistance by modifying the physiology and morphology of the plants [44].

5.1 Factors Affecting the Availability of Zinc

Parent material from which the soils are derived has a significant effect on the availability of Zn. The soils originating from gneisses, granite, sandstone, and limestone are found to be lower in the available Zn contents [48]. The major concern in the availability of Zn lies in its fixed/ insoluble/ unexchangeable forms [35].

Availability of Zn is highly reliant on pH, with an increase in pH above 6, Zn availability decreases with an increase in pH. Precipitation of Zn into insoluble forms of Zn, $ZnSiO_4$ or $ZnFe_2O_4$ will take place leading to reduced soluble Zn / available Zn.

Zn deficiency is more likely to be noticed in alkaline soils than in acid soils. In soils with high hydroxyl (OH^-) ions, the response of the plants to

external application of the Zn is negligible. Under alkaline situations Zn is precipitated as $Zn(OH)_2$ or $ZnCO_3$ [48,49]. Soils with higher carbonate contents will relatively exhibit unavailability of the Zn due to absorption of Zn onto soil particles and turned to unexchangeable forms [50].

Zn deficiency is more prevalent in soils with low organic matter. In high organic matter-containing soils available Zn is accordingly high [51].

Humic acid and fulvic acids are the identified stable organic compounds comparably contain more functional group ions like OH, COOH, and SH which show an affinity to Zn^{2+} [35]. Fulvic acids generally form chelates with Zn increasing the mobility and solubility of Zn. In case of the high organic soils like peat and muck binding of zinc to humic substances takes place and makes Zn unavailable [52].

Soil texture also affects the Zn availability, sandy soils with lighter texture exhibit more deficiency in available Zn compared to the heavier textured clay soils. Clay soils with high CEC values certainly have more affinity to retain Zn [51,35].

Soil moisture significantly affects the availability of nutrients to the plants as the diffusion of nutrients to roots is affected by soil moisture. Zn nutrition is considered to be at stake in arid and semi-arid regions due to the coincidence of prolonged water deficit periods during the growing season. Whereas, in water-logged situations Zn co-precipitates with iron and aluminium making Zn unavailable to the plants [51,35]. Compared to dry soils, flooded soils are more prone to Zn deficiency because in the prior case, Zn reacts with free sulfide. In case of rice delayed application of Zn until after flooding rice decreases the fixing of zinc with sesquioxides [53,49].

Fluctuations in temperature towards the colder side below $16^\circ C$ during the growing period will hinder the zinc availability to the plants due to change in the rate of Zn mineralization. Besides plants suffer Zn deficiency as a result of unfavourable climatic situations [50].

5.2 Available Sources of Zn

The following order describes the reduction in solubility of several zinc minerals: $Zn(OH)_2$ (amorphous) > $Zn(OH)_2$ > $ZnCO_3$ (smithsonite) > ZnO (zincite) > $Zn(PO_4)_2 \cdot 4H_2O$ (willemite) > soil Zn > $ZnFe_2O_4$ (franklinite). Zinc forms

soluble complexes with chloride, phosphate, nitrate, and sulfate ions however, the neutral sulfate ($ZnSO_4$) and phosphate ($ZnHPO_4$) species are the most significant and contribute to the overall concentration of zinc in solution. When acidifying fertilizers like ammonium sulphate [$(NH_4)(SO_4)_2$] are employed, the $ZnSO_4$ complex accounts for the enhanced availability of zinc and may increase the solubility of Zn^{2+} in soils [54]. Most commonly available forms of zinc sulfate are $(ZnSO_4 \cdot H_2O)$ -33% Zn - Zinc sulfate monohydrate, $(ZnSO_4 \cdot 7H_2O)$ -21% Zn- Zinc sulfate heptahydrate and chelated form as $(ZnEDTA)$ -9% Zn [55].

5.3 Nano Zn- ZnO

Zinc oxide nanoparticles in a colloidal solution are utilized as fertilizer. A plant nutrient known as nano fertilizer is more than just a fertilizer because it not only gives plants the nutrients, they need, but also restores the soil to its organic state without the negative effects of chemical fertilizers. Nano fertilizers have the benefit of being able to be applied in extremely small quantities [56]. ZnO as a fertilizer source will aid in improved growth and yield resulting in an enhanced qualitative and quantitative value of the crop [57].

5.4 Zinc Deficiency Symptoms in Plants

In general, the deficiency symptoms of Zn include: shortening of internodes leading to stunted growth, and malformation of leaves which will be visible as little leaf and rosette leaf formation. Usually, the deficient symptoms tend to be visible in younger leaves because under deficient conditions Zn gets immobile in plants. Necrosis symptom is often seen in the middle-aged leaves, interveinal chlorosis is another predominant symptom visible under Zn-deficient situations.

In rice, chlorosis of the basal leaves, loss of turgidity in leaves leading to wilting, bronzing of the leaves, and death of the young seedlings is seen. In younger seedlings of age 2-4 weeks after transplanting or sowing, chlorosis of the youngest leaves in the midrib at the base is seen. The appearance of brown spots on the older leaves, which eventually enlarge and coalesce to form brown-colored leaves [51]. Other predominant symptoms include reduced tillering, stunted growth, loss of turgidity and plants falling on the surface of the water and floating.

In many cases, the Zn deficiency can be mistaken for the deficiency of N, Mg, Mn, or Fe. So, plant analysis is often recommended to detect Zn deficiency.

6. EFFECTIVE ZINC MANAGEMENT METHODS IN RICE BASED CROPPING SYSTEMS

Soil conditions and rice production systems play a crucial role in deciding the method of application of zinc to the plants to maximize benefits. The most promising methods followed in rice-based cropping systems are soil application, seed priming, zinc encapsulated fertilizers, foliar application, seed coating, seedling root dip method, bio-fertilizer-based application.

Soil application of Zn fertilizers is found to be effective in promoting the growth, yield, and biofortification of the grains. Basal application of zinc has improved the yield by 18% compared to control [58]. Plants are capable of absorbing soluble compounds and gases through leaves. The application of foliar sprays makes the leaves absorb them and export them to the growing tissues. Jose [59] reported that higher grain and straw yields of rice were recorded with foliar spray of ZnSO₄ at 0.1% at different growth stages. Combined soil + foliar applied Zn resulted in maximum kernel yield and kernel Zn concentration in fine rice [60]. Improved yield components and nutrient recovery through Zn foliar nutrition are observed by [61].

Seed coating with nutrients like zinc using plant-based substances as adhesives is also a possible method to maximize the benefits. Zulfiqar et al. [58] Reported 13 % increase in yield following seed coating with Zn in rice.

By partially hydrating seeds, a practice known as "seed priming," germination-related metabolic processes can start to take place but radicle emergence is prevented. This is a feasible method to raise the Zn content in seeds to improve yield-attributing characteristics and yields because of the tiny amounts of Zn required for seed priming and the simplicity of the procedure [32]. Seed priming has a significant effect on yield and grain Zn contents in rice under different production systems and zinc application methods [60].

The growth, Zn nutrient content, and Zn nutrient uptake of rice were all found to be improved by

inoculating it with arbuscular mycorrhizal fungus [62]. Replacing 25% of inorganic nutrients with bio-fertilizers has shown promising nutrient uptake and performance of zinc in rice [61].

7. CONCLUSION

The review of earlier studies highlights that India is highly at risk of zinc malnutrition and the majority of soils are deficient in nutrients in the first place and zinc nutrition is often neglected in agriculture. Correcting the zinc deficiency in rice-based cropping systems will be a significant endeavor to achieve the benefits of biofortification through fertilization because rice-based cropping systems are widely used in India and rice is the main diet for the majority of Indians. As availability of Zn to plants is dependable on various factors related to the soil pH, organic matter, moisture availability etc. and considering the factors responsible for unavailability from the available sources of Zn, effective zinc management methods like seed priming, foliar application etc. has to be adopted to deal with zinc deficiency.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Fageria NK, Baligar VC, Clark RB. Micronutrients in crop production. Adv. Agron. 2002;77:189–272.
2. Stanton C, Sanders D, Krämer U, Podar D. Zinc in plants: integrating homeostasis and biofortification. Mol. Plant. 2002;15, 65–85.
3. Wu Q, Liu C, Wang Z, Gao T, Liu Y, Xia, Y. Zinc regulation of iron uptake and translocation in rice (*Oryza sativa* L.): Implication from stable iron isotopes and transporter genes. Environ. Pollut. 2022; 297:118818.
4. Gondal AH, Zafar A, Dua-e-Zainab, Toor MD, Sohail S, Ameen S, Ijaz AB, Imran C.B, Hussain I, Haider S, Ahmad IA, Rehman B, Younas, N. A detailed review study of zinc involvement in animal, plant and human nutrition. Ind. J. Pure App. Biosci. 2021;9(2):262-271.
5. Black RE. Micronutrients in pregnancy. Br. J. Nutr. 2001;85:193-197.
6. Stein AJ. Rethinking the measurement of under nutrition in a broader health context: Should we look at possible causes or

- actual effects?.Glob. Food. Secur. 2014; 3(3&4):193-199.
7. NIH [National Institutes of Health]. Zinc fact sheet for health professionals; 2021. . Available:<https://ods.od.nih.gov/factsheets/Zinc-HealthProfessional/>. [10 November 2021].
 8. Samreen T, Humaira HUS, Ullah S, Javid, M. Zinc effect on growth rate, chlorophyll, protein and mineral contents of hydroponically grown mung beans plant (*Vigna radiata*). Arab. J. Chem. 2017; 10(1):1802–1807.
 9. Hacisalihoglu G. Zinc (Zn): The last nutrient in the alphabet and shedding light on Zn efficiency for the future of crop production under suboptimal Zn. Plants. 2020;9:1471.
 10. Rehman H, Aziz T, Farooq M, Wakeel A, Rengel Z. Zinc nutrition in rice production systems: A review. Plant Soil. 2020;361: 203-226.
 11. Chen XP, Zhang YQ, Tong YP, Xue YF, Liu DY, Zhang W, Cui ZL. Harvesting more grain zinc of wheat for human health. Sci. Rep. 2017;7(1):7016
 12. FAO [Food and Agriculture Organization]; 2018. Available:<https://www.fao.org/food-systems/en/>.
 13. Zulfiqar U, Hussain S, Maqsood M, Ishfaq M, Ali, N. 2021. Zinc nutrition in puddle transplanted and direct seeded production systems to enhance productivity, zinc use efficiency and grain biofortification of fine aromatic rice. J. Plant Growth Regul. 2017; 40:1539-1556.
 14. Tuti DM, Kumar M, Saha S, Singh A. 2018. Rice-based cropping systems for enhancing productivity of food grains in India: decadal experience of AICRP. Indian Farming. 2017;68(01):27–30.
 15. Sarwar N, Atique-ur-Rehman, Ahmad S, Hasanuzzaman, M. Rice-Based Cropping Systems.(eds) Modern Techniques of Rice Crop Production. Springer, Singapore; 2022.
 16. Jensen ES, Peoples MB, Nielsen HH. Faba bean in cropping systems. Field Crops Res. 2010;115:203-216.
 17. Sravan US, Murthy KVR. Enhancing productivity in rice-based cropping systems. Intech Open. 2018;2010:59-75.
 18. Verma RK, Shivay YS, Choudhary M, Ghasal PC, Madar, R. Nutrient mobilization and crop assimilation as influenced by nutrient management strategies under direct seede basmati rice (*Oryza sativa*)-based cropping systems. Indian J. Agric. Sci. 2020;90(10):1894-1901.
 19. Singh KK, Singh YV, Sharma SK. Influence of biofertilizers and urea application on grain yield and quality attributes in rice (*Oryza sativa* L.) cultivars. J Soil Water Conserv. 2010;9(4):271–76.
 20. Garai TK, Datta JK, Mondal NK. Evaluation of integrated nutrient management on boro rice in alluvial soil and its impacts upon growth, yield attributes, yield and soil nutrient status. Arch. Agron. Soil Sci. 2013; 60(1): 1–14.
 21. Verma RK, Shivay YS, Ghasal PC. Effect of different cropping systems and nutrient sources on growth, productivity and economics of direct seeded basmati rice (*Oryza sativa*). Indian J. Agric. Sci. 2017; 87(10):1377–83.
 22. Alam MJ, Al-Mahmud A, Islam MA, Hossain M.F, Ali MA, Dessoky ES, El-Hallous EI, Hassan, MM, Begum N, Hossain, A. Crop diversification in rice-basedcropping systems improves the system productivity, profitability and sustainability. Sustainability. 2021;13:6288.
 23. Nagoli SB, Basavanneppa MA, Sawargaonkar GL, Biradar DP, Biradar SA, Tevari, P. Diversification of Rice-rice (*Oryza sativa* L.) cropping systems for productivity, profitability and resource use efficiency in Tunga Bhadra Project Command Area. BEPLS. 2017;6:108–114.
 24. Samant TK. System productivity, profitability, sustainability and soil health as influenced by rice based cropping systems under mid central table land zone of Odisha. Int. J. Agric. Sci. 2015;7:746–749.
 25. Jat RA, Dungrani RA, Arvadia MK, Sahrawat KL. Diversification of rice (*Oryza sativa* L.)-based cropping systems for higher productivity, resource-use efficiency and economic returns in south Gujarat. India. Arch. Agron. Soil Sci. 2015;58:561-572.
 26. Singh G, Sharma M, Manan J, Singh G. Assessment of soil fertility status under different cropping sequences in District Kapurthala. J Krishi vigyan. 2016;5(1): 1-9.
 27. Singh SR, Yadav P, SinghD, Bahadur L, Singh SP, Yadav AK., Mishra A, Yadav PPS, Kumar, S. 2020. Impact of different crop grown systems on land nutrient index, microbial diversity and soil quality. Land Degrad Dev. 2016;32(14): 3973-3991.

28. Shankar T, Banerjee M, Malik GC, Dutta S, Maiti D, Maitra S, Alharby H, Bamagoos A, Hossain, A, Ismail IA. The Productivity and Nutrient Use Efficiency of Rice–Rice–Black Gram Cropping Sequence Are Influenced by Location Specific Nutrient Management. Sustainability. 2021;13:3222.
29. Xu X, He P, Pampolino MF, Johnston AM, Qiu S, Zhao S, Chuan L, Zhou W. Fertilizer recommendation for maize in China based on yield response and agronomic efficiency. Field Crop. Res. 2014;157:27–34.
30. Kumar M, Singh SK, Patra A. Effect of different nutrient sources on yield and biochemical properties of soil under rice–wheat cropping sequence in middle Gangetic alluvial plain, J. Plant Nutr. 2021; 44(15):2310-2330.
31. Shukla AK, Behera SK, Pakhre A, Chaudhari SK. Micronutrients in soils, plants, animals and Humans. Indian J. Fertilizers. 2019;14(4):30-54.
32. Sharma M, Parmar DK, Kumar P, Sankhyan NK, Kumar P, Butail NP. Efficacy of zinc nutrition for improving zinc content at different growth stages of rainfed pea-maize sequence in North Western Himalaya. J. Plant Nutr. 2021; 44(20):1-11.
33. IISS [Indian Institute of Soil Science]. 2021. Extent of micro- and secondary nutrient deficiencies in Indian soils and their correction. Available: <https://iiss.icar.gov.in/#modal-one>. [11 November 2021].
34. Shukla AK, Behera SK, Prakash C, Tripathi A, Patra AK, Dwivedi BS, Trivedi V, Rao CS, Chaudhari SK, Das S, Singh AK. 2021. Deficiency of phyto-available sulphur, zinc, boron, iron, copper and manganese in soils of India. Sci. rep. 2021; 11:19760.
35. Suganya A, Saravanan A, Manivannan N. Role of Zinc Nutrition for Increasing Zinc Availability, Uptake, Yield, and Quality of Maize (*Zea Mays* L.) Grains: An Overview. Commun Soil Sci Plant Anal. 2020;51:15.
36. Hafeez-ur Rehman, Aziz T, Farooq M, Wakeel A, Rengel, Z. Zinc nutrition in rice production systems: a review. Plant Soil. 2012;361:203–226.
37. Liu J, Zhu R, Xu T, Xu, Y, Ge F, Xi, Y, Zhu J, He H. Co-adsorption of phosphate and zinc (II) on the surface of ferrihydrite. Chemosphere. 2016;144:1148–1155.
38. Rudani L, Vishal P, Kalavati, P. The importance of zinc in plant growth-A review. Int. Res. J. Nat. Appl. Sci. 2018;5: 38–48.
39. Hassan MU, Aamer M, Chattha MU, Haiying T, Shahzad B, Barbanti L, Nawaz M, Rasheed A, Afzal A, Liu Y, Guoqin H. The critical role of zinc in plants facing the drought stress. Agric. 2020;10(9):396.
40. Hajiboland R, Amirzad, F. Drought tolerance in Zn-deficient red cabbage (*Brassica oleracea* L. var. capitata f. rubra) plants. Hortic. Sci. 2010;37:88-98.
41. Grungreiff K, Gottstein T, Reinhold, D. 2020. Zinc deficiency-An independent risk factor in the pathogenesis of haemorrhagic stroke?. Nutrients. 2010;12:3548.
42. Romheld V, Marschner H. Function of micronutrients in plants. In Micronutrients in agriculture, ed. JJ Mortvedt FR Cox LM. Shuman, and RM. Welch. Madison, USA: Soil Science Society of America, Book Series No. 4. 1991;297–328
43. Singh B, Kumar S, Natesan A, Singh BK, Usha K. Improving Zinc deficiency of cereals under zinc deficiency. Current Science. 2005;88:36–44.
44. Kandil EE, El-Banna AAA, Tabl DMM, Mackled MI, Ghareeb RY, Al-Huqail AA, Ali HM, Jebri J, Abdelsalam NR. Zinc Nutrition Responses to Agronomic and Yield Traits, Kernel Quality, and Pollen Viability in Rice (*Oryza sativa* L.). Front. Plant Sci. 2022;13:791066.
45. Ghazi S, Diab AM, Khalafalla MM, Mohamed RA. Synergistic effects of selenium and zinc oxide nanoparticles on growth performance, hemato-biochemical profile, immune and oxidative stress responses, and intestinal morphometry of Nile tilapia (*Oreochromis niloticus*). Biol. Trace Elem. Res. 2022;200:364–374.
46. Mousavi SR, Galavi M, Rezaei M. Zinc importance for crop production - a review. Int. J. Agron. Plant Prod. 2013;4:64-68.
47. Sadeghzadeh B. A review of zinc nutrition and plant breeding. J. Soil Sci. Plant Nutri. 2013;13:905–27.
48. Barak P, Helmke PA. The chemistry of Zinc. In Zinc in soils and plants, ed. A.D. Robin. Dordrecht: Kluwer Academic Publishers; 1993;90-106
49. Suganya A, Saravanan A. DTPA – Zn in pH varied soils under simulated moisture conditions as influenced by graded levels of Zn in combination with zinc solubilizing bacteria. Biosci. Trends. 2015;8:812–15.

50. Alloway BJ. Micronutrients and crop production. In Micronutrient deficiencies in global crop production, 2008;1–39.
51. Hafeez B, Khanif YM, Saleem M. Role of zinc in plant nutrition – a review. Am. J. Exp. Agric. 2013; 3(2): 374-91.
52. Katyal JC, Randhawa NS. 1983. Micronutrients FAO Fertilizer and Plant Nutrition Bulletin 7. Rome: Food and Agriculture Organization of the United Nations.
53. Suganya A, Saravanan A. DTPA – Zn in soil under simulated moisture conditions as influenced by graded levels of Zn in combination with zinc solubilizing bacteria. Biosci. Trends. 2014; 7:3968–71.
54. Arunachalam P, Kannan P, Prabhukumar G, Govindaraj M. Zinc deficiency in Indian soils with special focus to enrich zinc in peanut. African J. Agric. Res. 2013;8(50): 6681-6688.
55. Gite D, Gite H, Darvhankar M, Gite P. Zinc sulphate a potential micronutrient for wheat growth: A review. Pharma Innovation. 2021;10(5):473-475.
56. Sabir S, Arshad M, Chaudhari SK. Zinc oxide nanoparticles for revolutionizing agriculture: synthesis and applications. Sci. World J. 2014;1-8.
57. Azam M, Bhatti HN, Khan A, Zafar L, Iqbal M. Zinc oxide nano-fertilizer application (foliar and soil) effect on the growth, photosynthetic pigments and antioxidant system of maize cultivar. Biocatal. Agric. Biotechnol. 2022;42:102343.
58. Zulfiqar U, Hussain S, Maqsood M, Ishfaq M. Zinc nutrition to enhance rice productivity, zinc use efficiency and grain biofortification under different production systems. Crop Ecol. Manag. Quality. 2020;10:1-11
59. Jose A. Agronomic biofortification of zinc in rice (*Oryza sativa* L.). M.Sc. (Ag) thesis, Kerala Agricultural University, Thrissur. 2019;119.
60. Ali MF, Adnan M, Khan, Asif M, Perveen S, Ammar A, Ali A, Hanif MS, Ali U, Rehman FU, Ilahi H, Raza A. Agronomic zinc biofortification in rice production systems. Plant Cell Biotechnol. Mol. Biol. 2021;22(25&26):103-113.
61. El-Sobky EEA., Taha AE, El-Sharnouby M, Sayed SM, Elrys AS. Zinc-biochemical co-fertilization improves rice performance and reduces nutrient surplus under semi-arid environmental conditions. Saudi J Biol Sci. 2022;29(3):1653-1667.
62. Purakayastha TJ, Chhonkar PK. Influence of vesicular-arbuscular mycorrhizal fungi (*Glomus etunicatum* L.) on mobilization of zinc in wetland rice (*Oryza sativa* L.). Biol Fertil Soils. 2001; 33: 323-327.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/114940>