



Prevalence of Schistosomiasis among Pupils of Umunya Central School, Oyi Local Government, Anambra State, Nigeria

Esiaba, G. C. ^{a*}, Chikwendu, J. I. ^a, Obinwa I. C. ^a
and Okoye, F.C. ^a

^a Department of Parasitology and Entomology, Nnamdi Azikiwe University, Awka, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpr/2024/v13i3285>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/116309>

Original Research Article

Received: 02/03/2024
Accepted: 06/05/2024
Published: 14/05/2024

ABSTRACT

Background: Schistosomiasis is a disease of medical importance caused by the parasitic worm *Schistosoma haematobium*. It is responsible for morbidities such as blood and protein deficiency in those infected. The study on the prevalence of urogenital schistosomiasis and contributory risk factors was carried out among pupils of Umunya Central School, Oyi Local Government, Anambra State between November and December, 2021.

Methods: The children were between the ages of 4-15 years. One hundred and twenty respondents of children were selected and their urine samples were collected and examined for the presence of haematuria and proteinuria using a combi 9 strip before the filtration technique was used to detect the presence of the parasite. Pupils data were collected using Google Forms researcher's questionnaire.

*Corresponding author: Email: esiabagrnt@gmail.com;

Cite as: Esiaba, G. C., Chikwendu, J. I., Obinwa I. C., & Okoye, F.C. (2024). Prevalence of Schistosomiasis among Pupils of Umunya Central School, Oyi Local Government, Anambra State, Nigeria. *International Journal of Pathogen Research*, 13(3), 46–55. <https://doi.org/10.9734/ijpr/2024/v13i3285>

Results: Out of the 120 (62 males and 58 females) children examined, a total prevalence of 0% was gotten. The pupils had a good source of water although a few of them (8%) still drank from the rivers in the community which could predispose them to the disease. Knowledge of the disease was very little among the children and many of them visited the rivers for different activities (78%).
Conclusion: Health education of the children and residents for improved perception of the behavioural and socio-economic activities associated with the disease will help reduce the burden of disease transmission.

Keywords: Schistosomiasis; schistosoma haematobium; local government; prevalence.

1. INTRODUCTION

Schistosomiasis also known as bilharzia or snail fever is an infectious disease caused by blood trematode of the genus *Schistosoma*. It is the most important and prevalent water-borne disease, the parasite was first observed in Cairo in 1851 by a young pathologist, Theodor Bilharz in the blood of mesenteric veins of a young man on autopsy. There are three major species under the genus *Schistosoma*, that can infect humans, and these are; *Schistosoma haematobium*, *Schistosoma mansoni* and *Schistosoma japonicum* [1] while other species include; *Schistosoma mekongi* and *Schistosoma intercalatum*.

Furthermore, schistosomiasis is among the set of 17 neglected tropical infectious diseases and it majorly affects people who live in these areas (tropics). It is also the second leading endemic parasitic disease after malaria on the list of parasitic diseases in 76 countries worldwide [2] and the disease has been implicated to be the cause of over 280,000 deaths each year in Africa alone [3]. Nigeria, with 20 million people requiring schistosomiasis preventive chemotherapy, is ranked first among the countries of the world endemic for schistosomiasis [4]. The Federal Ministry of Health in 2015 recorded that Nigeria had the highest burden of schistosomiasis in the world. The overall prevalence was 9.5% and it was present in all 20 states surveyed but was higher in the northern part of Nigeria.

The aetiological agent of urogenital Schistosomiasis is the infective stage i.e. cercaria, of *S. haematobium*, a digenic trematode of the phylum Platyhelminthes, whose intermediate hosts are some species of Gastropod Snails in the genus *Bulinus* [5]. These snails are found in freshwater, therefore proximity of communities to freshwater sources and irrigated lands predisposes them to the infective stage of the parasite (Cercariae). Some human activities also play a huge role in

predisposing humans to infection by this parasite. Schistosome eggs are excreted by humans in faeces and urine. After hatching, miracidia infects specific snails which develops into cercariae. *Schistosoma cercariae* penetrates the skin of humans during domestic (for example, washing clothes or food materials) and Recreational activities (for example., bathing and swimming in infected freshwater bodies). Other factors that facilitate the transmission of schistosomiasis include; proximity to freshwater bodies, socioeconomic factors which influence occupational activities (e.g., poor individuals without running water at home are likely to contact freshwater bodies) and climate change [6]. Lack of access to improved sanitation facilities contributes to open defecation into freshwater bodies which also enhances the transmission of Schistosomiasis. Human-induced ecological transformations like dam construction and irrigation scheme developments are becoming the major risk factors for the resurgence of the parasitic disease.

Symptoms of schistosomiasis are caused by the body's reactions to the worm's eggs. Urogenital schistosomiasis can present as chronic, which is the most common or acute haematuria (blood in urine), and dysuria (painful urination) in the early symptoms of the disease and it is worth noting that the severity of the disease is dependent on the intensity of the infection [7]. In women, urogenital schistosomiasis may present with genital lesions, vaginal bleeding, pain during sexual intercourse and nodules in the vagina. It could also induce pathology of the seminal vesicles, prostate, and other organs in men. The disease could also have other long-term irreversible consequences, including infertility [8]. In children, schistosomiasis can cause anaemia, stunting and a reduced ability to learn, although the effects are usually reversible with treatment [8]. The best ways of preventing schistosomiasis include avoiding swimming in freshwater containing cercariae, drinking safe water, and swimming in treated pools. Control measures

could include massive drug administration to communities and targeted treatment of school-age children (CDC, 2017). Diagnosis of infection is confirmed by the identification of eggs in stools or urine specimens. Antibodies and/or antigens detected in blood or urine samples are also indications of infection. For urogenital schistosomiasis, a filtration technique using nylon, paper or polycarbonate filters is the standard diagnostic technique. Children with *S. haematobium* almost always have microscopic blood in their urine which can be detected by chemical reagent strips.

1.1 Significance of Study

Previous studies have shown that schoolchildren are highly susceptible to the infection. Although past studies indicate the prevalence of urogenital schistosomiasis in Anambra State (2.9% and 5.5% for microscopy and hematuria respectively) [9], but current information on the prevalence of urogenital schistosomiasis in Umunya is lacking. The school children, representing a greater percentage of the community and being the group more susceptible to the disease's morbidity, makes them an important group for carrying out this investigation. Carrying out this study would help determine if there would be an urgent need for the government to start an integrated, targeted and effective schistosomiasis control programme among these pupils to help improve their health.

1.2 Aim and Objectives

Thus, this study was aimed at investigating the prevalence and contributory risk factors of urogenital schistosomiasis among pupils in Umunya Central School in Oyi local Government area of Anambra state, Nigeria.

2. MATERIALS AND METHODS

2.1 Study Area

This study was carried out in the Umunya community in Oyi Local Government Area of Anambra State, Southeast Nigeria. Umunya is a town situated at the centre of Anambra State and it lies within latitudes of 6° 12' 34" N and longitudes 6° 54' 51" E. It is a town of ten villages namely Ezi Umunya, Okpu, Ojobi, Umuebo, Amaezike, Ajakpani, Odumodu-Ani, Isioye, Odumodu-Enu and Ukunu. These villages are sub-grouped into Ezi-lfite and Ikenga sub-

divisions, the tri-partite heritage of all Gadite H/Igbos commonly referred to in anthropological history as ERI-AWKA Igbo. The climate of the area is typically tropical with distinct wet and dry seasons with a temperature range of 23-35°C between June and December but rises to 25-37°C between January and April. Farming and trading over the years has become the major occupation of the people while some of them are transport workers and civil servants. They rely on borehole water and well water as their source of water.

2.2 Study Design

The study design was a cross-sectional design to determine the prevalence and contributory risk factors of urogenital schistosomiasis among pupils in Umunya Central School, Oyi local government area in Anambra State. The study countered both surveys of the pupils and laboratory studies of their urine specimens. One urine sample was collected from each pupil. The study was carried out between October and December 2021.


2.3 Sample Population

The study population comprises of a total number of 171 Pupils from Primary 1 to Primary 6, 120 pupils were selected on a stratified random basis from Umunya Central School in Oyi Local Government Area, Anambra State. The pupils were selected between the ages of 4-15 years from primary one to primary six.

2.4 Sample Size

About 171 pupils make up the study population being pupils from Primary 1 to Primary 6. Sample size formula, $n = \frac{N}{1+N(e^2)}$. Where n is minimum sample size, N is the study population, e is the error term, 0.05 at 95% confidence interval, was used to determine the sample size for the study.

Minimum Sample size, $n = \frac{171}{1+171(0.05^2)} = \frac{171}{1+0.4275} = \frac{171}{1.4275} = 119.78 = 120$. Sample size n=120 pupils were purposively selected for the study. To evenly distribute the number of students per class, the minimum sample size was divided by the number of classes: $\frac{120}{6} = 20$. Therefore 20 urine samples were taken from each class (Primary 1 to Primary 6) making a total of 120 samples which were used for the study.



Map 1. Map of Anambra state showing Umunya

2.5 Administration of Questionnaires

A structured, Google-form researcher's questionnaire was used for data collection. Data on pupils' age, gender, activities carried out in water bodies, parent's occupation as well as their view and knowledge of schistosomiasis, which was referred to as "Okpo mammiri" in their local

parlance, was obtained using Google forms. Data obtained was analyzed subsequently.

2.6 Collection of Urine Sample

The urine was collected between 10 am - 2 pm to coincide with the peak of excretion of *S. haematobium* eggs. Sterile 20ml plastic universal

specimen bottle with screw-caps was used to collect terminal urine samples from the selected pupils. The samples were appropriately labelled with identification numbers and were transported to the lab where the analysis was done.

2.7 Determination of Biochemical Parameters

Reagent strips (Medi-Test Combi 9) were dipped into the Urine inside the universal containers. Various parameters such as glucose, protein, blood, urobilinogen, nitrite, bilirubin and pH were evaluated according to the manufacturer's instructions.

2.8 Parasitological Examination using the Filtration Technique

The filtration technique was used in other to recover ova from the urine samples [10].

2.9 Apparatus

A packet of polycarbonate membrane filters, forceps, 10ml syringes, 13mm filter holder, cotton wool, beaker, Lugol's iodine, slide, coverslip and a microscope.

2.10 Procedure

- Using the blunt-ended (untoothed) forceps, a polycarbonate membrane filter was picked and placed carefully on the filter support of the filter holder (13mm diameter).
- The filter holder was then attached to the end of a 10ml Luer syringe.
- The plunger was removed from the syringe before the syringe was filled to the 10ml mark with well-mixed urine after which the plunger was replaced.
- The syringe was then held over a beaker and the urine sample was gently passed through the filter.
- The filter holder was removed and unscrewed before the blunt-ended forceps were used to carefully remove the membrane filter which was then transferred in such a way as to leave the eggs intact on the filter paper. The filter paper was faced upwards while being placed on the slide.
- A drop of Lugol's iodine was added to the slide and covered with a cover slip.

- Using the x10 objective with the condense and the iris closed sufficiently to give good contrast, the entire filter was examined systematically by adjusting the microscope until a clear image was obtained.

2.11 Determination of Intensity of Infection

The number of eggs was counted and reported as egg per 10ml of urine. 1-10 eggs/10ml urine signifies light infection, 11-49 eggs/10ml of urine as mild infection and >50eggs/10ml of urine as heavy infection.

Microscopic examination of the samples was performed at the Laboratory of Parasitology and Entomology, Nnamdi Azikiwe University Awka, Anambra state.

2.12 Data Analysis

The data obtained from the sample examination and questionnaires were represented on tables concerning age group, sex, prevalence and other risk factors such as activities carried out in natural bodies of water and Knowledge of the disease (Urogenital Schistosomiasis). Analysis was done using the Chi-square statistical method.

3. RESULTS

Six classes were selected for the study (Primary 1 to 6) and 20 samples were taken each from the selected classes summing 120 samples which were examined from Umunya Central School. Out of the 120 samples examined 0 (0%) tested positive for urogenital schistosomiasis.

3.1 Prevalence According to Sex

Out of the 120-sample collected and examined, 62 were males and 58 were females and there were no positive samples. This shows that there was no significant association between males and females in the prevalence of schistosomiasis.

3.2 Prevalence According to Age

The pupils were between the ages of 4 to 15 and of these age groups, no positive samples were observed.

Table 1. Prevalence of urogenital schistosomiasis by sex

Sex	No. Examined	No. Infected (%)
Male	62 (51.7%)	0
Female	58 (48.3%)	0
Total	120 (100%)	0

Table 2. Prevalence of urogenital schistosomiasis by age group

Age interval	No. Examined	No. Infected (%)
4 – 6	15	0
7 – 9	54	0
10 – 12	50	0
13 – 15	1	0
Total	120	0

Table 3. Risk factors of Schistosomiasis as observed among the pupils

Risk Factors	Observation	Percentage
Activities carried out in Natural water body	Bathing: 50	42%
	Washing clothes: 29	24%
	Defecation/Urination: 6	5%
	Fishing: 3	2%
	Washing food materials:6	5%
	No contact with river: 26	22%
Knowledge of the Disease (Urogenital Schistosomiasis)	Know about it: 23	19%
	Ignorant: 97	81%
Source of drinking water	Shallow wells: 11	9%
	Tankers: 11	9%
	Streams: 10	8%
	Borehole: 48	38%
	Pipe borne water: 45	36%
Number of Dewormed	Dewormed: 92	77%
	Not Dewormed: 28	23%




Fig. 1. Showing the pupil's source of drinking water




Fig. 2. Showing Number of Dewormed pupils




Fig. 3. Showing pupils knowledge of the disease




Fig. 4. Showing activities carried out by pupils in natural water bodies

3.3 Risk factors of Schistosomiasis Observed among the Selected Children

From the questionnaire given to assess the risk factors promoting the spread of schistosomiasis, the following risk factors were observed. A greater number of children indicated that they make use of Borehole water (48%). Most of them make use of pipe-borne water (45%), while a few of them (10%) still make use of rivers as a source of drinking water. 78% of them indicated that they still visit the rivers around them to carry out different activities, while 22% indicated that they don't go to the rivers for any activity. 81% of them were ignorant of what schistosomiasis was all about, its causes and transmission while the remaining 19% admitted to having heard of schistosomiasis or "Okpo mammiri" as they called it in their local Parlance. 77% representing a greater percentage of the pupils indicated to have recently taken worm medicines, while 23% of the pupils have not been dewormed.

4. DISCUSSION

Schistosomiasis is a disease of medical importance. It is prevalent in Nigeria where it is responsible for grave economic losses [11].

These results showed a total 120 school-aged pupils, aged 4-15 years old were investigated in Umunya.

The findings of this study reveal the prevalence of urinary schistosomiasis among pupils in Umunya Central School, Oyi Local Government Area, Anambra State was 0% out of the 120 examined pupils. The prevalence in this work was relatively low compared to other studies reporting schistosomiasis in other places for instance Yvonne et al. [12] reported that the prevalence of urogenital schistosomiasis in Anambra North was 6.0%, Which may be because the rivers and streams are not close to the schools which limits students' activities in them. Aribodor et al. [13] also recorded a prevalence of 2.1% among school pupils in Nsugbe community, Anambra state. In Ebonyi state, prevalence was up to 49.7% in Ohaukwu Local Government Area of the state (Obionu et al., 2014). Studies in other parts of Anambra and Maidugri reported a lower prevalence from a 2013 survey reported a prevalence of 9.5% (Federal Ministry of Health, 2015).

The 0% prevalence could be a result of some factors which were observed during the research. From the study, it was observed that 45(36%) and 48(38%) of the study population used pipe-

borne water and boreholes as their major source of drinking water. This could be a contributory factor to the low prevalence of schistosomiasis, as this generally reduces their contact with freshwater bodies and its use as a source of drinking water thereby reducing the possibility of infection [14].

Deworming using Praziquantel Preventive Chemotherapy (PC) has proven to be very effective and schistosomiasis can be easily controlled and treated with a single annual dose of praziquantel, which can also reverse up to 90% of the damage done by the parasite [7]. From the collected data, a greater percentage (77%) of the pupils indicated to have always partaken in the annual deworming programme in the school which is pioneered by the government.

5. CONCLUSION AND RECOMMENDATION

Transmission of schistosomiasis in communities is enhanced by the resident's use of unsafe water sources of water for both personal and domestic activities as well as behavioural and socio-economic tendencies that promote risky contacts with potentially parasite-infested bodies. It has also been observed that constant use of bore-holes, pipe borne and participation in annual praziquantel chemotherapy has reduced the prevalence of urogenital schistosomiasis to zero. Nevertheless, as was observed, ignorance of the disease was high among the children, therefore, to maintain the zero level of prevalence; health education and awareness should be created among the pupils and residents of the communities for improved perception of the behavioural and socio-economic activities associated with the disease. Children's activities should also be regulated as they stand a higher chance of infection.

ETHICAL APPROVAL AND CONSENT

A letter of introduction was obtained from the Department of Parasitology and Entomology, Nnamdi Azikiwe University, Awka. An advocacy visit was made to first inform parents, teachers and students about the research project and written consent and informed assent were obtained from parents and pupils respectively prior to the commencement of the research. The volunteers (participants) were also made aware

of the aim of the research, additionally, volunteers were assured that the results would be solely used for research purposes.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Colley DG, Bustinduy AL, Secor E, King CH. Human schistosomiasis. *The Lancet*. 2014;28;383(9936):2253–2264.
2. Mubyazi G, Donohue R, Madon S, Mashoto K, Michael E, Malecela M. Biosocial determinants of persistent Schistosomiasis among schoolchildren in Tanzania despite repeated treatment, *Tropical Medicine and Infectious Disease*. 2017;2(4):61-70.
3. Chala B, Torben W. An epidemiological trend of urogenital schistosomiasis in Ethiopia. *Frontiers in public Health*. 2018;5(6):60-70.
4. World Health Organisation (WHO). PCT databank Schistosomiasis; 2019. https://www.who.int/neglected_diseases/preventive_chemotherapy/sch/en
5. Tolulope EA, Stephen DA, Kingsley ME. The current epidemiological status of urogenital schistosomiasis among primary school pupils in Kastina State, Nigeria: An imperative for a scale up of water and Sanitation initiative and mass administration of medicines with Praziquantel. *PLoS Neglected Tropical Diseases*. 2018;12(7):7-15.
6. McCreesh N, Nikulin G, Booth M. Predicting the effects of climate change on *Schistosoma mansoni* transmission in eastern Africa. *Parasites & vectors*. 2015;8:1-9.
7. Carter care. Schistosomiasis control programme; 2015. <http://www.cartercenter.org>.
8. WHO, schistosomiasis; 2021. <https://www.who.int/health-topics/schistosomiasis?>
9. Ndukwe YE, Obiezue RN, Aguzie IN, Anunobi JT, Okafor FC. Mapping of urinary schistosomiasis in Anambra State, Nigeria. *Annals of Global Health*. 2019;2'85(1):52.
10. Obisike VU, Amuta EU, Audu AB, Kwenev SA, Comparison of polycarbonate filter

- paper and sedimentation methods in diagnosing *Schistosoma haematobium* infection in Makurdi, Benue, Nigeria. South Asian Journal of Parasitology. 2019;2(1):1-6.
11. Chikwendu JI, Onekutu A, Ogbonna IO, Amuta EU. Association between *Schistosoma mansoni* Infection rates in humans and in *Biomphalaria pfeifferi* snails in Akwanga, Nasarawa State, Nigeria. South Asian Journal of Research in Microbiology. 2019;5(1):1-8.
 12. Ndukwe, Yvonne, Obiezue, Rose, Aguzie, Ifeanyi, Toochukwu Joy, Anunobi, Okafor, Fabian. Mapping of urinary schistosomiasis in Anambra State, Nigeria. Annals of Global Health. 2019; 85(52):10.5334.
 13. Aribodor DN, Basse SA, Yoonuan T, Sam-Wobo SO, Aribodor OB, Ugwuanyi IK. Analysis of schistosomiasis and soil-transmitted helminths mixed infections among pupils in Enugu State, Nigeria: Implications for control. Infections Disease and Health. 2019;24(2):98-106.
 14. Amuta EU, Onekutu A, Ogbonna IO, Chikwendu JI, Water contact activities affecting prevalence and distribution of intestinal schistosomiasis in two endemic communities gwanje and mada hills, Akwanga, Nigeria. Trends in Applied Sciences Research. 2020;15:275-280.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/116309>