



Effect of Different Growing Media and Varying Levels of Gibberellic Acid on Seed Germination and Vigour of Acid Lime (*Citrus aurantifolia* Swingle.) cv.Kagzi Lime

Aishwarya ^{a++*} and Vijay Bahadur ^{a#}

^a Department of Horticulture, Sam Higginbottom University of Agriculture Technology and Sciences, Naini, Prayagraj, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i81124>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119413>

Original Research Article

Received: 06/05/2024

Accepted: 09/07/2024

Published: 11/07/2024

ABSTRACT

The present investigation "Effect of different growing media and varying levels of gibberellic acid on seed germination and vigour of acid lime (*Citrus aurantifolia* Swingle.) cv.Kagzi lime" was laid out on the experimental site of Department of Horticulture, Sam Higginbottom University of Agriculture Technology & Sciences, Naini, Prayagraj (UP), during 2023. The present investigation was carried out in randomized block design with 13 treatment which were replicated thrice. The treatments

⁺⁺ M.Sc. Fruit Science;

[#] Associate professor;

^{*}Corresponding author: E-mail: aishdubey631@gmail.com;

Cite as: Aishwarya, and Vijay Bahadur. 2024. "Effect of Different Growing Media and Varying Levels of Gibberellic Acid on Seed Germination and Vigour of Acid Lime (*Citrus Aurantifolia* Swingle.) cv.Kagzi Lime". *Journal of Advances in Biology & Biotechnology* 27 (8):91-96. <https://doi.org/10.9734/jabb/2024/v27i81124>.

were T₀ (Control), T₁ (Sand + FYM + GA3 150mg/l), T₂ (Sand + FYM + GA3 200mg/l), T₃ (Sand + FYM + GA3 250mg/l), T₄ (Sand + Vermicompost + GA3 150mg/l), T₅ (Sand + Vermicompost + GA3 200mg/l), T₆ (Sand + Vermicompost + GA3 250mg/l), T₇ (Sand + Coco peat + GA3 150mg/l), T₈ (Sand + Coco peat + GA3 200mg/l), T₉ (Sand + Coco peat + GA3 250mg/l), T₁₀ (Sand + Coco peat + Vermicompost + GA3 150mg/l), T₁₁ (Sand + Coco peat + Vermicompost + GA3 200mg/l), T₁₂ (Sand + Coco peat + Vermicompost + GA3 + 250mg/l). On the basis of our experimental findings it can be concluded that the treatment T₄ (Sand + Vermicompost + GA3 150mg/l) Was found to be best in the terms of germination percentage growth parameters and survival percentage of Acid lime.

Keywords: FYM; vermicompost; coco peat; GA3, kagzi lime.

1. INTRODUCTION

The cultivation of acid lime (*Citrus aurantifolia* Swingle), commonly known as Kagzi lime, is of significant economic importance in many tropical and subtropical regions worldwide. This citrus fruit is esteemed for its tart flavor, aromatic qualities, and versatility in culinary and medicinal applications. Maximizing the yield and quality of acid lime crops is a primary concern for growers, necessitating research into various factors influencing seed germination and plant vigor. *Citrus aurantifolia*, commonly known as acid lime, is a small evergreen tree belonging to the Rutaceae family. This research is justified by several reasons. Understanding the optimal conditions for seed germination and seedling vigor is crucial for enhancing the productivity and quality of acid lime crops. By examining the influence of different growing media, such as soil, vermicompost or coco peat, on seed germination and vigor, this research can provide valuable insights into selecting the most conducive substrate for acid lime cultivation.

Growing media play an important role in germination of seed. It is a substrate that provides the required elements and physical support to the seed. Media should also have good water holding capacity, drainage and other physical and chemical properties. During seed germinations, the role of GA3 in the induction of synthesis of α - amylase and other hydrolytic enzymes among monocots and certain dicots is well documented. GA3 appears mainly to induce the activity of the gluconeogenic enzymes during early stages of seed germinations [1].

Farmyard manure (FYM), often referred to as "green manure," is a traditional organic fertilizer derived from decomposed animal waste, primarily from cattle, horses, sheep, or poultry. It is a valuable source of nutrients, organic matter, and beneficial microorganisms, making it an essential component of sustainable agriculture

and soil management practices. FYM is rich in essential plant nutrients such as nitrogen (N), phosphorus (P), potassium (K), and micronutrients like calcium, magnesium, and sulfur [2].

Vermicompost is an organic fertilizer and soil amendment produced through the decomposition of organic materials by earthworms, typically species such as *Eisenia fetida* and *Lumbricus rubellus*. This process, known as vermicomposting, yields a nutrient-rich, humus-like material prized for its ability to improve soil health and enhance plant growth. The vermicomposting process begins by providing a suitable habitat for earthworms, often in containers or specialized vermicomposting systems [3].

2. MATERIALS AND METHODS

The present investigation "Effect of different growing media and varying levels of gibberellic acid on seed germination and vigour of acid lime (*Citrus aurantifolia* Swingle.) cv.Kagzi lime" was laid out on the experimental site of Department of Horticulture, Sam Higginbottom University of Agriculture Technology & Sciences, Naini, Prayagraj (UP), during 2023. The present investigation was carried out in randomized block design with 13 treatment which were replicated thrice. The treatments were T₀ (Control), T₁ (Sand + FYM + GA3 150mg/l), T₂ (Sand + FYM + GA3 200mg/l), T₃ (Sand + FYM + GA3 250mg/l), T₄ (Sand + Vermicompost + GA3 150mg/l), T₅ (Sand + Vermicompost + GA3 200mg/l), T₆ (Sand + Vermicompost + GA3 250mg/l), T₇ (Sand + Coco peat + GA3 150mg/l), T₈ (Sand + Coco peat + GA3 200mg/l), T₉ (Sand + Coco peat + GA3 250mg/l), T₁₀ (Sand + Coco peat + Vermicompost + GA3 150mg/l), T₁₁ (Sand + Coco peat + Vermicompost + GA3 200mg/l), T₁₂ (Sand + Coco peat + Vermicompost + GA3 + 250mg/l). Seeds were kept in GA3 solutions for 18 hours. The

research work was carried out the objective of growing media and gibberellic acid on germination and survival percentage of kagzi lime. Data was analyzed using OPSTAT.

3. RESULTS AND DISCUSSION

The effect different growing media and varying levels of gibberellic acid on seed germination of kagzi lime is very obvious and consistent. There was significant difference among the different treatments. Among the treatment applied the maximum seed germination percentage was obtained in the treatment T₄ (Sand + Vermicompost + GA₃ 150mg/l) with 85.07 % followed by treatment T₅ (Sand + Vermicompost + GA₃ 200mg/l) with 83.12 % which was statistically superior over control T₀ (Control) with 72.36 %. The results are in conformity with the findings of Farooqui et al. [4] in sapota.

The effect different growing media and varying levels of gibberellic acid on seed vigor index of kagzi lime is very obvious and consistent. There was significant difference among the different treatments. Among the treatment applied the maximum seed vigor index was obtained in the treatment T₄ (Sand + Vermicompost + GA₃ 150mg/l) with 1538 followed by treatment T₅ (Sand + Vermicompost + GA₃ 200mg/l) with 1408 which was statistically superior over control T₀ (Control) with 1055. Together, GA₃ and vermicompost synergistically promote acid lime seed germination by regulating cellular processes, ensuring optimal conditions for germination, and providing essential nutrients and growth-promoting substances.

The maximum plant height of kagzi lime was recorded in treatment T₄ (Sand + Vermicompost + GA₃ 150mg/l) with (33.92) cm followed by treatment T₅ (Sand + Vermicompost + GA₃ 200mg/l) with (33.82) cm and the minimum plant height was recorded in T₀ (control) with (23.01) cm. Gibberellic acid (GA₃) plays a crucial role in increasing the plant height of acid lime at the cellular level. GA₃ is a plant growth regulator that primarily stimulates cell elongation and division. At the cellular level, GA₃ interacts with specific receptors in the cell membrane, initiating signal transduction pathways that activate genes responsible for cell elongation. Similar result was obtained by Chaudhary et al. [5].

The maximum Number of leaves of kagzi lime was recorded in treatment T₄ (Sand + Vermicompost + GA₃ 150mg/l) with (16.87)

followed by treatment T₅ (Sand + Vermicompost + GA₃ 200mg/l) with (15.53) and the minimum Number of leaves was recorded in T₀ (control) with (11.53) . At the cellular level, GA₃ interacts with receptors on the cell membrane, initiating signaling cascades that activate transcription factors responsible for leaf primordia formation. Additionally, GA₃ stimulates the expression of genes encoding proteins crucial for leaf growth and expansion. By promoting cell division and expansion, GA₃ facilitates the formation of new leaves. A similar finding was reported by Choudhary and Chakrawar [6] in Rangpur lime.

The maximum Plant spread of kagzi lime was recorded in treatment T₄ (Sand + Vermicompost + GA₃ 150mg/l) with (8.08 and 7.76) cm followed by treatment T₅ (Sand + Vermicompost + GA₃ 200mg/l) with (8.97 and 8.09) cm and the minimum Plant spread was recorded in T₀ (control) with (4.17 and 4.19) cm. At the cellular level, organic manure improves soil structure and nutrient availability, facilitating better root penetration and lateral root growth, which contribute to increased plant spread. Together, GA₃ and organic manure synergistically promote cellular processes that enhance plant spread, leading to healthier and more vigorous acid lime plants with increased lateral expansion. Similar result was obtained by Chaudhary et al. [5].

The maximum stem diameter of kagzi lime was recorded in treatment T₄ (Sand + Vermicompost + GA₃ 150mg/l) with (9.28) mm followed by treatment T₅ (Sand + Vermicompost + GA₃ 200mg/l) with (9.09) mm and the minimum Stem Diameter was recorded in T₀ (control) with (8.25) mm. Gibberellic acid (GA₃) and organic manure play integral roles in increasing the stem diameter of acid lime at the cellular level. GA₃, a plant growth regulator, promotes cell elongation and division, influencing stem thickness. At the cellular level, GA₃ interacts with receptors on the cell membrane, initiating signaling pathways that activate genes responsible for cell expansion and secondary growth. Organic manure, such as compost or vermicompost, enriches the soil with nutrients and beneficial microorganisms, supporting robust root development and overall plant vigor. A similar result was obtained by Bhardwaj et al. [7] in seedling growth of papaya.

The maximum Leaf area of kagzi lime was recorded in treatment T₄ (Sand + Vermicompost + GA₃ 150mg/l) with (10.33) cm² followed by treatment T₅ (Sand + Vermicompost + GA₃ 200mg/l) with (10.15) cm² and the minimum Leaf

Table 1. Effect of different growing media and varying levels of gibberellic acid on Seed germination, seed vigour index, plan height, no. of leaves, plant spread, stem diameter, leaf area, chlorophyll content and Survival percentage of kagzi lime

Symbol	Seed Germination (%)	Seed Vigour Index	Plant height (cm)	No. of leaves	Plant spread		Stem diameter	Leaf area (cm ²)	Chlorophyll Content	Survival percentage
					N-S	E-W				
T ₀	72.36	1055	23.01	11.53	4.17	4.19	8.25	8.92	3.29	63.79
T ₁	80.45	1203	30.45	13.53	5.49	6.37	8.79	9.18	3.55	74.47
T ₂	80.67	1367	32.66	13.49	5.39	6.28	8.44	10.02	3.65	76.57
T ₃	82.82	1390	31.32	13.87	6.12	7.6	8.79	9.63	3.73	75.37
T ₄	85.07	1538	33.92	16.87	8.08	7.76	9.28	10.33	5.44	88.72
T ₅	83.12	1408	33.82	15.53	8.97	8.09	9.09	10.15	5.07	85.11
T ₆	81.00	1300	31.09	12.20	7.27	6.21	8.62	9.27	4.63	78.39
T ₇	82.67	1242	29.93	14.20	7.69	6.47	8.50	9.21	4.04	80.02
T ₈	79.87	1276	32.86	14.53	7.9	6.54	8.75	9.41	3.83	79.58
T ₉	81.32	1313	32.26	14.20	6.93	6.78	8.71	9.44	3.68	82.62
T ₁₀	79.63	1358	32.32	15.29	5.88	6.69	8.70	9.55	4.00	78.73
T ₁₁	80.65	1387	31.64	14.87	7.28	7.49	8.77	10.04	3.87	78.97
T ₁₂	82.43	1398	33.45	14.64	7.46	7.13	8.79	9.85	3.97	76.98
F-Test	S	S	S	S	S	S	S	S	S	S
SEd(±)	1.04	36.87	0.059	0.085	0.026	0.024	0.039	0.18	0.220	0.220
CD@5%	2.19	73.74	0.118	0.175	0.054	0.051	0.078	0.453	0.643	0.643
CV	4.16	110.61	0.177	0.255	0.078	0.072	0.117	0.576	0.965	0.965

area was recorded in T₀ (control) with (8.92) cm². Organic manure, such as compost or vermicompost, enriches the soil with nutrients and beneficial microorganisms, supporting vigorous plant growth. At the cellular level, organic manure enhances nutrient availability and metabolic processes, facilitating optimal leaf development and expansion. By synergistically promoting cellular processes, GA₃ and organic manure enhance leaf area in acid lime plants, resulting in increased photosynthetic capacity and overall plant vigor. Similar results was obtained by Haribhau et al. [8] in rangpur lime.

The maximum Chlorophyll Content was obtained in the treatment T₄ (Sand + Vermicompost + GA₃ 150mg/l) with 5.44 followed by treatment T₅ (Sand + Vermicompost + GA₃ 200mg/l) with 5.07 which was statistically superior over control T₀ (Control) with 3.29. Gibberellic acid (GA₃) has a complex effect on chlorophyll. While it can stimulate leaf growth, which could indirectly lead to more chlorophyll production due to more leaf area, high GA₃ concentrations can actually decrease chlorophyll content. This is because GA₃ may trigger chlorophyll breakdown in favor of other pigments. So, GA₃'s role in chlorophyll content is dependent on factors like concentration and plant species. Similar result was obtained by Chaudhary et al. [5].

The maximum Survival percentage was obtained in the treatment T₄ (Sand + Vermicompost + GA₃ 150mg/l) with 88.72 % followed by treatment T₅ (Sand + Vermicompost + GA₃ 200mg/l) with 85.11 % which was statistically superior over control T₀ (Control) with 63.79 %. A winning combination for acid lime survival can be achieved with GA₃ and organic manures [9,10]. GA₃, applied before sowing, acts like a germination booster, leading to more uniform and earlier sprouting, giving seedlings a crucial head start. Organic manures, like compost, enrich the soil with vital nutrients like nitrogen and phosphorus, while also improving water retention and aeration. Similar result was obtained by Prajapati [11] Joshi et al. [12] in Acid lime and Dilip et al. [13].

4. CONCLUSION

On the basis of our experimental findings it can be concluded that the treatment T₄ (Sand + Vermicompost + GA₃ 150mg/l) Was found to be best in the terms of germination percentage growth parameters and survival percentage of Acid lime.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that generative AI technology have been used during writing or editing of manuscript.

Details of the AI usage are given below:
Chat GPT v.3.5.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Suthar D, Kotecha A, Thounaojam, Amarjeet. Effect of growing media and gibberellic acid on seed vigour and survival of acid lime (*Citrus aurantifolia* Swingle.) cv. Kagzi lime. 2021;291-293.
2. Bhatt, Rajan Naresh RK, Tiwari, Himanshu Singh PK, Das, Dhritiman Kumar, Manoj Banik, Rahul Tomar, Gaurav. Farm Yard Manure as a Resource for Integrated Nutrient Management. 2023;12:672-681. DOI:10.5281/zenodo.7826550.
3. Hakeem, Rahmatullah, Shrivastava, Purnima, Bhat, Umar, Patel, Chirag. Research paper on vermiculture, eco-friendly vermicompost and vermiwash also its influence in the soil nutrient, plant growth, and enhance yield especially Apple productivity in the Union territory (UT) of Jammu and Kashmir. SSRN Electronic Journal. 2021;8:366-373.
4. Farooqui AA, Nalawadi UG, Sulladmath UV. Effect of growth regulators on the germination of sapota (*Achras zapota* Mill foseberg) seeds. The Mysore J Agric Sci. 1991;5:341-343.
5. Chaudhary, Asha Ahlawat, Timur & Kumar, Sanjeev & Jena, Suchismita & Patel, Dharmishtha. Effect of Gibberellic Acid on Germination and Vigour of Kagzi Lime Seedlings. Current Journal of Applied Science and Technology. 2019; 1-8. DOI:10.9734/cjast/2019/v38i630411.
6. Choudhary BK, Chakrawar VK. Effect of seed treatment using some chemicals on shoot and root growth of Rangpur Lime. J Maharashtra Agric. Univ. 1982;1(1):66- 60.
7. Bhardwaj RL. Effect of growing media on seed germination and seedling growth of papaya cv. 'Redlady'. Academic J. 2008;8 (4):178-184.

8. Katkar Ankush Haribhau, MS Paikra, UB Deshmukh, LK Ramteke, Dikeshwar Nishad, Manish Taram and Aaradhana Rathore. Effect of GA3 and growing media on seed germination, growth and vigour of Rangpur lime (*Citrus limonia* Osbeck). The Pharma Innovation Journal. 2022; 11(12): 3056-3061.
9. Scott SJ, Jones RA, Williams W. Review of data analysis methods for seed germination 1. Crop science. 1984;24(6): 1192-9.
10. Miransari M, Smith DL. Plant hormones and seed germination. Environmental and experimental botany. 2014;99:110-21.
11. Prajapati DD. Influence of growth regulators on germination of jackfruit (*Artocarpus heterophyllus* Lam.) seed. M. Sc Thesis, Navsari Agricultural University, Navsari, Gujarat; 2013.
12. Joshi PS, Sahoo AK, Bhoyar RK, Meshram PC. Effect of various plant growth promoting substances on seedling growth of acid lime. Trends in Biosciences. 2015; 8(19):5222-5225.
13. Dilip WS, Singh D, Moharana D, Rout S, Patra SS. Effect of gibberellic acid (GA) different concentrations at different intervals on seed germination and seedling growth of rangpur lime. Journal of Agroecology and Natural Resource Management. 2017;4(2): 157-165.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/119413>