



Profiling Bioactive Compounds in the Bark of *Cinchona officinalis* L. and Anti-arthritic Properties Thereof

N. Senthilkumar ^{a*} and R. Sumathi ^a

^a ICFRE- Institute of Forest Genetics and Tree Breeding, R.S. Puram, Coimbatore, Tamil Nadu. India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajb2t/2024/v10i3208>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119655>

Original Research Article

Received: 08/05/2024

Accepted: 11/07/2024

Published: 16/07/2024

ABSTRACT

The bark samples of *C. officinalis* were collected from The Nilgiris district of Tamil Nadu. The collected samples were processed and extracted using methanol solvent. The extracts showed high yield of secondary metabolites especially alkaloids, phenols, flavonoids, steroids, terpenoids, sterols, quinones and tannins. Alkaloid content was very high in *C.officinalis* followed by phenol, flavonoid, steroid and Tannin. The secondary metabolites present in *C. officinalis* (Bark) were isolated, purified, identified and characterized using differential Chromatographic techniques such as TLC, HPLC and GC-MS/ MS. The compounds identified using High Performance Liquid Chromatography (HPLC) technique in bark of *C. officinalis* in the retention times of 1.750, 2.250, 4.016 and 7.266 minutes were quinine, isoquinidine and cinchonidine. The compounds were then characterized using GC- MS/ MS technique with reference to NIST library. The compounds were eluted from the bark of *C. officinalis* in the retention time of 9.569, 10.488, 11.814, 13.593, 14.282, 16.542, 16.967, 23.494, 24.954, 25.151, 25.988, 26.752, 26.982, 27.906, 28.277, 31.454, 33.604,

*Corresponding author: E-mail: senthilnk@icfre.org;

Cite as: Senthilkumar, N., and R. Sumathi. 2024. "Profiling Bioactive Compounds in the Bark of *Cinchona Officinalis* L. And Anti-Arthritic Properties Thereof". *Asian Journal of Biotechnology and Bioresource Technology* 10 (3):15-22. <https://doi.org/10.9734/ajb2t/2024/v10i3208>.

and 35.744 minutes. By correlating the retention times with (NIST) mass spectral library the eluted compounds were confirmed as Phthalic acid, Hexadecanoic acid, 1,2,3,5-Cyclohexanetetrol, Cinchonab-9-one, 9,12-Octadecadienoic acid (Z,Z)-, Heptadecanoic acid, Quinine, Cinchonine, Cinchonidine, Phenol, Quinine, and gamma.-Sitosterol. The phytochemicals present in the *C. officinalis* (Bark) samples were evaluated for their biological properties viz., anti-inflammatory activity and anti-arthritic activity in in-vitro conditions. Anti-inflammatory properties of plant samples were tested by protein denaturation method using diclofenac (75 mg/ ml concentration) as positive control. The methanol extract of *C.officinalis* (Bark) showed anti-inflammatory properties of range between 17.0 and 57.3%. Anti-arthritic activity of bark extract of *C. officinalis* (75 mg/ ml concentration) was evaluated by albumin denaturation method using diclofenac (75 mg/ ml concentration) as positive control. The methanol extract of *C.officinalis* (Bark) showed anti-arthritic properties of range between 18.4 and 58.5%. From the study it is understood that the alkaloids and polyphenols present in the methanolic extract of bark of *C. officinalis* were responsible for their biological properties. Hence, *C. officinalis* can serve as an alternative to synthetic drugs available in the market to treat arthritis subject to safety (toxicity) studies in humans. Based on the *In vitro* results, it was confirmed that the bark of *C.officinalis* possesses comparatively high biological values. Thus an herbal aerosol for arthritis may be formulated using the bark extract of *C.officinalis* as key ingredient and the therapeutic product will be of cost effective, easy to use and pocket friendly.

Keywords: *Cinchona officinalis*; arthritis; phyto-compounds; HPLC; GCMS.

1. INTRODUCTION

Arthritis is the leading cause of disability in adults worldwide, limiting the daily activities for billions of people [1]. Globally more than 100 types of arthritis are there. The most common arthritis are osteoarthritis, rheumatoid arthritis, psoriatic arthritis, and septic arthritis, which are autoimmune diseases, often originate from the joints (Aiyalu et al. 2017). Arthritis affects people of all ages and has been known to mankind since ancient times. Scientists found evidence of arthritis in the knees of Nefertari's mummy. Mitch Leslie reports in science that the study of old knees provides evidence supporting this model and suggests that arthritis may not be the fate of old age.

The symptoms of arthritis may include joint pain, stiffness, tenderness, lack of movement, grating inside the joints, fever, chills and redness. Ultimately, because of the avascular nature of cartilage, once damage has occurred, it cannot be repaired, thus making a cure essentially impossible [2]. Treatment aims to control pain, reduce joint damage and improve or maintain quality of life. It includes physical therapy, medication and patient education and support. Ancient medicine is also used to treat arthritis. Clinical studies conducted in recent years have helped shed light on this disease. Although there is no cure for arthritis, researchers have found many treatments to control symptoms. Treatment of arthritis depends on the nature and severity of the underlying disease. The main goal is to

reduce pain and improve the function of affected joints before more serious problems occur.

NSAIDs are anti-inflammatory, analgesic, and antipyretic agents, and they are typically used to reduce chronic pain, decrease stiffness and improve function in patients with osteoarthritis, Rheumatoid arthritis and other forms of arthritis [3]. In addition, steroidal drugs, Immunosuppressant's, Corticosteroid medicines and synthetic drugs such as hydroxychloroquine, leflunomide, methotrexate and sulfasalazin are often prescribed medicine for arthritis. Though the synthetic drugs are efficient in pain relief, long term intake of these drugs can cause wide range of side effects. Studies of older adults show that chronic NSAIDs use increases the risk of peptic ulcer disease, acute renal failure, and stroke/ myocardial infections (Marcum et al., 2010). Although research has not proven that specific herbs or supplements can treat arthritis, many herbs can reduce joint pain. There are so many herbal products available in the market for oral intake and topical application including tablets, powder, syrup, oil and spray etc. The existing topical formulations have the disadvantages like difficulty in applying the formulation to respective site, staining cloth due to greasy nature, more chances of contamination and more wastage due to inaccuracy of dose (Aiyalu et al., 2017).

Cinchona officinalis possess many biological properties which includes antimalarial [4], Lupus erythematosus [5], SARS COVID 19 [6] and

arthritis. Considerably less number of research works is carried out to substantiate the efficacy of *C. officinalis* against arthritis. In this work the bark of *C. officinalis* was selected due to the potentially active biological ingredients present in it. The present study focused on the profiling active compounds in bark extract of *C. officinalis* and evaluation of anti-inflammatory and anti-arthritic activities thereof. The phytochemicals present in the bark extract of *C. officinalis* were qualitatively and quantitatively estimated using standard protocols followed by characterization using differential chromatographic techniques such as thin layer Chromatography (TLC), High Performance Liquid Chromatography (HPLC) and Gas Chromatography coupled with double Mass Spectrometer (GC-MS/ MS). Subsequently anti-inflammatory and anti-arthritic activities of *C. officinalis* (Bark) were also evaluated to determine its ant-arthritic efficacy.

2. MATERIALS AND METHODS

The bark samples of *C. officinalis* were collected from the original sources at Medicinal Plants Development Area (11°25'37" N Latitude and 76°43'48" E Longitude), Cinchona Village, Doddabetta, The Nilgiris district, Tamil Nadu (INDIA). Qualitative and quantitative evaluation of methanol extract of bark of cinchona was performed by following the standard protocols to detect and quantify the phytochemicals present in it [7] (Harborne, 1984). Plant metabolites like alkaloids, phenols, flavonoids and Tannins were isolated and characterised using differential chromatographic techniques such as TLC, HPLC and GC-MS/ MS [8]. The biological properties like anti-inflammatory and anti-arthritic activities of the secondary metabolites were determined by Protein denaturation [9] and Albumin denaturation (Vaijanthimala et al., 2019) methods respectively.

3. RESULTS AND DISCUSSION

The bark of *C. officinalis* collected from The Nilgiris district of Tamil Nadu was authenticated by a taxonomist at Institute of Forest Genetics and Tree Breeding (IFGTB), Coimbatore, Tamil Nadu. The percentage yield of bark of *C.officinalis* was found to be 21.1% using methanol as solvent. Plants and herbs are rich sources in phytochemicals that prevent/treat or reduce many health issues. Qualitative evaluation of phytochemicals in the methanol

extract of bark sample *C.officinalis* revealed the presence of plant metabolites such as proteins, alkaloids, phenols, flavonoids, steroids, Terpenoids, Sterols and Quinones. Alkaloids such as colchicine, colchicoresin, democolchicine found in *Colchicum autumnale* (Colchicum) [10], strychnine, brucine and brucine N-oxide in *Strychnos nux-vomica* (Nux-vomica) [11], barbaloin and Aloe emodin in *Aloe vera* were reported to have anti-inflammatory activity [12,13].

3.1 Thin Layer Chromatography Analysis

Phytochemicals such as alkaloids, phenols, flavonoids and tannins find major applications in therapeutics. Hence these compounds were isolated and characterized using differential chromatographic techniques.

The spots (Compounds) observed on TLC plates for *C. officinalis* are 1 and 2 in case of alkaloids, phenols and flavonoids and 3 in tannins. RF value range 0 indicates the solvent polarity is very low and 1 indicating solvent polarity is very high. The stronger a compound is bound to the adsorbent, the slower it moves up the TLC plate. A non-polar compound moves up the plate more rapidly (higher RF value) than the polar compounds (lower RF value). The RF value so obtained for all the compounds in average is very close to 1 which means the polarity of the compounds is very low (non- polar compounds).

3.2 High Performance Liquid Chromatography Analysis

Pratiwi et al. [14] reported that the commercially vital alkaloid quinine was isolated from cell cultures of Cinchona species. HPLC has proved as one such most useful technique used for the separation of quinoline alkaloids. The HPLC chromatogram denoted the prominent peaks of the eluted compounds of *C. officinalis* (Bark) in the retention time of 1.750, 2.250, 4.016 and 7.266 minutes. The eluted compounds were identified as quinine, isoquinidine and cinchonidine. Reece and Peikert [15] used HPLC method for rapid quantification of plasma levels of quinidine, dihydroquinidine and 3-hydroxyquinidine. Similarly, Murauer and Ganzera [16] isolated dihydroquinidine, dihydroquinine, quinidine, quinine, cinchonine and cinchonidine from the bark of cinchona.

Table 1. Qualitative and quantitative analysis of phytochemicals in methanol extract of *C.officinalis* (Bark)

S.No	Phytochemical constituents	Name of the conformation test	Phytochemical screening	Phytochemical quantification Mg/g
1	Alkaloids	Mayer's Test	+	0.24
2	Phenols	Lead acetate Test	+	0.82
3	Flavonoids	H ₂ SO ₄ Test	+	1.61
4	Steroids	Salkowski Test	+	2.86
5	Terpenoids	Salkowski Test	+	0.13
7	Sterols	Salkowski Test	+	22.79
8	Quinones	H ₂ SO ₄ Test	+	33.40

Table 2. Retention factor of alkaloids, phenols, flavonoids and tannins in the plant samples

Compounds	<i>C. officinalis</i> (Bark)			Standard
	1	2	3	
Alkaloids	0.96	0.73	-	0.94
Phenols	0.98	0.8	-	0.96
Flavonoids	0.94	0.34	-	0.94
Tannins	0.92	0.65	0.26	0.91




Fig. 1. TLC separation of alkaloids, phenols, flavonoids and tannins of *C. officinalis* bark methanol extract




Fig. 2. HPLC Chromatogram plot of *C. officinalis* (Bark)

3.3 Gas Chromatography and Mass Spectrometry Analysis

The phytochemicals in the plant samples were characterised based on the interpretation of mass spectrum of GC-MS/MS in reference with the database of National Institute of Standard and Technology (NIST). The NIST 20 EI Library has 350,643 spectra for 306,869 compounds and over 447 k literature and experimentally determined GC methods and Retention indexes for 139,963 compounds. The name, molecular weight and structure of the components in the plant samples were ascertained by comparing the spectrum of unknown compounds with the spectrum of known compounds stored in the NIST library. Gas chromatography and Mass spectroscopy analysis of methanol extract of *C. officinalis* (Bark) was performed.

The chromatogram denoted the prominent peaks of the eluted compounds of *C. officinalis* (Bark) in the retention time of 9.569, 10.488, 11.814, 13.593, 14.282, 16.542, 16.967, 23.494, 24.954, 25.151, 25.988, 26.752, 26.982, 27.906, 28.277, 31.454, 33.604, and 35.744 minutes. By correlating the retention times with (NIST) mass spectral library the eluted compounds were confirmed as Phthalic acid, Hexadecanoic acid, 1,2,3,5-Cyclohexanetetrol, Cinchonab-9-one, 9,12-Octadecadienoic acid (Z,Z)-, Heptadecanoic acid, Quinine, Cinchonine, Cinchonidine, Phenol, Quinine, and gamma-Sitosterol. The eluted bioactive compounds present in the plant samples are responsible for their antimicrobial, antioxidant, anti-inflammatory and anti-arthritis activity. Phytotherapies approved for clinical treatments motivates the search for plant-based phytochemicals with therapeutic potential (Piovezana Bossolani et al., 2018). The botanicals especially flavonoids such as luteolin and by β -sitosterol isolated from the methanol extract of *Cissus quadrangularis* stem reported to have anti-inflammatory activity Rasale, [17].

3.4 Investigation of anti- Inflammatory Activity of bark extract of *C. officinalis*

Anti-inflammatory activity of bark extract of *C. officinalis* (75 mg/ ml concentration) was evaluated by protein (BSA) denaturation method using diclofenac (75 mg/ ml concentration) as positive control. The methanol extract of *C.officinalis* (Bark) have showed anti-inflammatory properties of range between 17.0 and 57.3%. Among the phytochemical components isolated in plants, flavonoids [18], alkaloids [19] and saponins [20] are reported to have effective anti-inflammatory activity. This activity is attributed to the greater antioxidant properties of polyphenols and presence of alkaloids in the plant samples. Hence these natural active compounds can be used as an alternative to synthetic, steroidal and non-steroidal drugs available in the market to treat pain and inflammation.

3.5 Investigation of Anti- Arthritic Activity of bark extract of *C. officinalis*.

Anti-arthritic activity of bark extract of *C. officinalis* (75 mg/ ml concentration) was evaluated by albumin denaturation method using diclofenac (75 mg/ ml concentration) as positive control. The methanol extract of *C.officinalis* (Bark) have showed anti-arthritic properties of range between 18.4- 58.5%. Based on the results obtained it was confirmed that the bioactive compounds present in the bark extract of *C. officinalis* inhibit the denaturation of proteins. Thus, the bark extract of *C. officinalis* can be used to treat arthritis subject to animal and human studies. Wang et al. [21] observed that the arthritis activity of *Urtica atrichocaulis* was the combined action of multiple phenolic constituents. Souto et al. [22] observed that the alkaloids such as isoquinoline, colchicines and aconitine isolated from plants were found effective against for arthritis treatment. Phytochemical compounds present in *C. officinalis* are most related for treatment of Arthritis [23-26].

Table 3. Anti-inflammatory activity of Diclofenac and *C.officinalis* (Bark)

Concentration	Diclofenac (%)	<i>C. officinalis</i> (Bark)- %
20 μ l	42.6	17.0
40 μ l	53.2	20.6
60 μ l	62.0	28.7
80 μ l	70.5	49.2
100 μ l	89.8	57.3




Fig. 3. GC-MS/ Chromatogram plot of bark of *C. officinalis*

Table 4. Anti-arthritic activity of Diclofenac, and *C.officinalis* (Bark)

Concentration	Diclofenac (%)	<i>C. officinalis</i> (Bark)- %
20 µl	49.2	18.4
40 µl	54.6	23.7
60 µl	63.7	30.4
80 µl	71.3	49.8
100 µl	90.8	58.5

4. CONCLUSION

The bioactive compounds present in the methanolic extract of bark of *C. officinalis* are responsible for various biological properties including anti-arthritic activity. The study revealed the anti-inflammatory and anti-arthritic properties of bark extract of *C.officinalis* by inhibiting the denaturation of proteins. All the active ingredients present in bark of *cinchona* was cited

for the anti-arthritic activity. Therefore, the bark extract of *C.officinalis* was used in the development of therapeutic product. The present work concludes to recommend to develop the formulation of an aerosol for arthritis using natural ingredients and the product is under development at Ceras Pharmaceuticals Private Limited. The product will be further tested among healthy individuals for its efficiency and toxicity.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Briggs AM, Cross MJ, Hoy DG, Sánchez-Riera L, Blyth FM, Woolf AD, March L. Musculoskeletal health conditions represent a global threat to healthy aging: A report for the 2015 World Health Organization world report on ageing and health. *The Gerontologist*. 2016;56(Suppl 2):S243–S255.
2. Hani D El-Gabalawy, Peter E Lipsky. Why do we not have a cure for rheumatoid arthritis?. *Arthritis Res*. 2002;4 Suppl 3(Suppl 3):S297-301. DOI: 10.1186/ar568
3. Simon LS. Nonsteroidal anti-inflammatory drugs and their risk: A story still in development. *Arthritis Res Ther*. 2013;15Suppl 3(Suppl 3):S1. DOI: 10.1186/ar4173
4. Permin H, Norn S, Kruse E, Kruse PR. On the history of Cinchona bark in the treatment of Malaria. *Dan MedicinhistArbog*. 2016;44:9-30.
5. Lubov JE, Jamison AS, Baltich Nelson B, Amudzi AA, Haas KN, Richmond JM. Medicinal plant extracts and natural compounds for the treatment of cutaneous lupus erythematosus: A systematic review. *Front Pharmacol*. 2022;13:802624. DOI: 10.3389/fphar.2022.802624. PMID: 35431950; PMCID: PMC9008762.
6. Arumugam S, Vadivel K, Dhaked DK, Prasanna VS, Yellurkar ML, Das P, Manoharan R, Thomas AJ, Singh L, Singh S, Velayutham R. Antiviral Activity of Cinchona officinalis, a Homeopathic Medicine, against COVID-19. *Homeopathy*. 2024;113(1):16-24. DOI: 10.1055/s-0043-1770339. Epub 2023 Sep 6. PMID: 37673083.
7. Harborne JB. *Phytochemical Methods*. 11th edition. New York, NY, USA: Chapman & Hall; 1984.
8. Senthilkumar N, Murugesan S, Vijayalakshmi KB. GC-MS-MS analysis of *Trichiliaconnaroides* (Wight & Arn.) Benth (Meliaceae): A tree of ethnobotanical records, *Asian Journal of Plant Science and Research*. 2012;2(2):193-197.
9. Williams LAD, O'Connar A, Latore L, Dennis O, Ringer S, Whittaker JA, Conrad J, Vogler B, Rosner H, Kraus W. The *In vitro* anti-denaturation effects induced by natural products and non-steroidal compounds in heat treated (Immunogenic) Bovine Serum Albumin is proposed as a screening assay for the detection of anti-inflammatory compounds, without the use of animals, in the early stages of the drug discovery process. *West Indian Med J*. 2008;57(4):327-331.
10. Sultana S, Asif M, Naveed A, Waqas M, Rehman S. Comprehensive review on ethanobotanical uses, phytochemistry and pharmacological properties of *Melia azedarach* Linn. *Asian Journal of Pharmaceutical Research and Health Care*. 2014;6(1):26-32.
11. Bhati R, Anupama Singh, Vikas Anand Saharan, Veerma Ram, Anil Bhandari. *Strychnosnux-vomica* seeds: Pharmacognostical standardization, extraction, and antidiabetic activity. *J Ayurveda Integr Med*. 2012;3(2):80–84.
12. Shelton RM. Aloe vera: Its chemical and therapeutic properties. *Int J Dermatol*. 1991;30(10): 679–83.
13. Kshirsagar AD, Panchal PV, Harle UN, Nanda RK, Shaikh HA. Antiinflammatory and anti-arthritic activity of anthraquinone derivatives in rodents. *Int. J. Inflam*. 2014;690596 1-12.
14. Pratiwi DR, Sumaryono Sari PT, Ratnadewi D. Cinchona cells performance in *in vitro* culture: Quinine alkaloid production with application of different elicitors. *IOP Conference Series: Earth and Environmental Science*. 2018;185: 012029. DOI:10.1088/1755-1315/185/1/012029
15. Reece PA, Peikert M. Simple and selective high-performance liquid chromatographic method for estimating plasma quinine. *Journal of Chromatography B: Biomedical Sciences and Applications*. 1980;181(2): 207-217.

16. Muraier A, Ganzera M. Quantitative determination of major alkaloids in Cinchona bark by Supercritical Fluid Chromatography. J Chromatogr A. 2018;1554:117-122. DOI: 10.1016/j.chroma.2018.04.038
17. Rasale PL. Phytochemical and pharmacological review of fracture healing drug asthisamharak used in ancient Indian medicine Int J Pharm Med Biol Sci. 2014;3(3):28-35.
18. Serafini M, Peluso I, Raguzzini A. Flavonoids as anti-inflammatory agents. Proc Nutr Soc. 2010;69(3):273-8.
19. Liu W, Zhang Y, Zhu W, Ma C, Ruan J, Long H, et al. Sinomenine inhibits the progression of rheumatoid arthritis by regulating the secretion of inflammatory cytokines and monocyte/ macrophage subsets. Front Immunol. 2018;9:2228.
20. Lee CJ, Moon SJ, Jeong JH, Lee S, Lee MH, Yoo SM, et al. Kaempferol targeting on the fibroblast growth factor receptor 3-ribosomal S6 kinase 2 signaling axis prevents the development of rheumatoid arthritis. Cell Death Dis. 2018;9(3).
21. Wang M, Li K, Nie Y, Wei Y, Li X. Antirheumatoid arthritis activities and chemical compositions of phenolic compounds-rich fraction from *urticaatrichocaulis*, an endemic plant to China. Evid Based Complement Alternat Med. 2012;2012:818230. DOI: 10.1155/2012/818230
22. Souto AL, Tavares JF, Da Silva MS, Diniz MDFFM, Athayde-Filho D, Filgueiras P, Barbosa Filho JM. Anti-inflammatory activity of alkaloids: an update from 2000 to 2010. Molec. 2011;16(10): 8515-8534.
23. Anderson LAAT Keene, Phillipson JD. J. Pharm. Pharmacol. 1982;34-45:6.
24. Anderson LAAT Keene, Phillipson JD. Planra Med. 1982a;46;25.
25. Piovezana Bossolani GD, Silva BT, Colombo Martins Perles JV, Lima MM, Vieira Frez FC, Garcia de Souza SR, et al. Rheumatoid arthritis induces enteric neurodegeneration and jejunal inflammation, and quercetin promotes neuroprotective and anti-inflammatory actions. Life Sci. 2019;238:116956.
26. Aiyalu R, Govindarjan A, Ramasamy A. Formulation and evaluation of topical herbal gel for the treatment of arthritis in animal model. Brazilian Journal of Pharmaceutical Sciences. 2016;52(3):493-507. Available:<http://dx.doi.org/10.1590/S1984-82502016000300015>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/119655>