



Kernel Trait and Genetic Variability Analysis in BC₂F₁ Maize Lines Derived from *Zea nicaraguensis* Crosses

Senthilkumar V.^a, Priya Garkoti^a, Varalakshmi S.^a,
Thotla Naresh^a and Narendra Kumar Singh^{a*}

^a Department of Genetics and Plant Breeding, G. B. Pant University of Agriculture & Technology, Pantnagar-263145 (Uttarakhand), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i81139>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/120100>

Original Research Article

Received: 13/05/2024
Accepted: 15/07/2024
Published: 17/07/2024

ABSTRACT

Maize (*Zea mays* L.) is an important cereal crop with significant applications in food, feed and industrial applications. This study aimed to increase maize yield and quality by evaluating core traits in introgressed *Zea nicaraguensis* individual's recipients of LM13 and LM14 maize inbred lines. Kernel type, shape, color, dimensions and weight were assessed in 35 BC₂F₁ plants using standardized descriptor guidelines. Analysis of genotypic and phenotypic coefficients of variation, heritability and genetic progress showed significant genetic diversity and the influence of environmental factors. Significant sources of variability highlighted by principal component analysis were kernel color, endosperm color, and 100-seed weight. High heritability and genetic progress in kernel color and length-to-width ratio indicate great selection potential. Reintroducing the diversity

*Corresponding author: E-mail: narendrksingh2@gmail.com;

Cite as: V., Senthilkumar, Priya Garkoti, Varalakshmi S., Thotla Naresh, and Narendra Kumar Singh. 2024. "Kernel Trait and Genetic Variability Analysis in BC₂F₁ Maize Lines Derived from *Zea Nicaraguensis* Crosses". *Journal of Advances in Biology & Biotechnology* 27 (8):268-79. <https://doi.org/10.9734/jabb/2024/v27i81139>.

lost in cultivated maize inherent in *Zea nicaraguensis* has the potential to improve both its agronomic performance and quality. Future research should aim to identify genetic markers and examine the interaction between genetic and environmental influences to optimize maize varieties for specific environments and consumer preferences, ultimately contributing to global food security and agricultural sustainability.

Keywords: Maize; *Zea nicaraguensis*; introgression breeding; kernel traits; genetic diversity.

1. INTRODUCTION

Maize (*Zea mays* L.) is the most important cereal crop in the world, holding wide applications in food, feed, and industrial uses [1]. As a staple food for millions of people, especially in developing countries, maize plays a critical role in food security and nutrition. It is also an important component of animal feed and serves as a raw material for various industrial products, including biofuels, starches and sweeteners [2]. It provides economic and nutritional importance; continuous improvement of maize yield and quality traits is imperative to meet the needs of a growing global population [3]. Kernel characteristics such as type, shape, colour, dimension, and weight are the main determinants of maize commercial value [4-6]. These are important not only for the appearance and marketability of maize but relate to its nutritional quality and hence suitability for different uses [7]. For instance, kernel size and weight are directly linked to grain yield, while kernel colour can indicate the existence of specific nutrients or anti-nutritional factors [8].

The genetic basis of kernel traits is complex, involving multiple genes and environmental interactions. Therefore, understanding the genetic variability underlying these traits is critical to developing effective breeding programs [9]. Introducing genetic diversity into breeding populations is a cornerstone of crop improvement. In maize, the use of wild relatives has proven to be an effective approach to broadening the genetic base and introducing new traits not found in cultivated varieties [10]. Wild relatives of maize, such as *Zea nicaraguensis*, possess unique traits that can enhance water-logging tolerance and nutritional quality [11,12]. Incorporating these traits into cultivated maize can lead to the development of superior genotypes with improved agronomic performance [13].

This investigation focuses on the BC₂F₁ population derived from crosses between LM13 and LM14 inbred maize lines and *Zea nicaraguensis*. To analyse the diversity of kernel traits in the descriptive analysis, genetic variability parameter analysis and principal component analysis were used [14,15]. In this population, we found a significant level of kernel variability, which included characteristics such as kernel type, shape, color, dimensions, and weight. These characteristics were selected because of their importance in determining yield, quality, and marketability of maize [16]. This analysis aids in understanding trait relationships and identifying primary factors for selection of kernel traits [17]. These study reports advance knowledge of maize genetic variability and provide a foundation for wild introgression breeding programs to improve kernel traits, agronomic performance, nutritional quality, and commercial value [18].

2. MATERIALS AND METHODS

2.1 Experimental Design and Plant Material

This experiment was conducted at the N.E.B. Crop Research Centre, GBPUA&T Pantnagar (Coordinates: 28.97°N, 79.41°E, Average Rainfall: 1443.9 mm). LM13 and LM14 maize inbred lines were crossed with the wild relative *Zea nicaraguensis* to develop F₁ hybrids. These were then backcrossed with the recurrent parents to produce BC₁F₁ and subsequently BC₂F₁ generation.

A total of 35 BC₂F₁ plants from each biparental cross combination were analysed for kernel traits including kernel type, shape of upper surface of kernel, kernel colour, endosperm colour, kernel width, kernel length, K L/B ratio, and 100 seed weight. All traits of data collection were observed based on CIMMYT Maize descriptor guidelines. This is schematically breeding process shown in Fig. 1.

Table 1. Kernel characteristics of parent materials: LM13, LM14, and *Zea nicaraguensis*

Parent Materials	LM13	LM14	<i>Zea nicaraguensis</i>
Kernel Type	Flint	Flint	Wild
Shape of Upper Surface of Kernel	Round	Round	Level
Kernel Colour (Top of grain)	Yellow	Yellow	Brown
Endosperm Colour	Pale Yellow	Pale Yellow	White
Kernel Length (mm)	9.00	7.00	6.00
Kernel Breadth(mm)	6.00	5.00	2.00
K L/B ratio	1.50	1.40	3.00
100 Seed Weight (g)	22.00	20.00	8.25




Fig. 1. Breeding Scheme for BC₂F₁ Generation from Crosses of LM13 and LM14 with *Zea nicaraguensis*

2.2 Statistical Analysis

To analysed the genetic and phenotypic variability of kernel traits, various statistical methods were employed. Descriptive statistics, including the computation of the range of kernel traits, offered a basic understanding of the degree of variation found in the BC₂F₁ population. Variability analysis, including the Genotypic Coefficient of Variation (GCV), Phenotypic Coefficient of Variation (PCV), Heritability (H²), and Genetic Advance (GA), was conducted using the 'Agricolae' package in R software version 4.1.2. PCA was performed to identify key traits contributing to variability. This

involved data standardization, extraction of eigenvalues and percentage of variance, and biplot visualization with the 'Factoextra' package. These combined approaches aimed to clarify the genetic basis of kernel traits and identify promising genotypes for breeding programs, leveraging the genetic diversity introduced by *Zea nicaraguensis* [14,15].

3. RESULTS AND DISCUSSION

3.1 Descriptive Statistics Analysis

Qualitative trait analysis: The data collected were differentiated between two populations:

LM13 BC₂F₁ and LM14 BC₂F₁. The results revealed notable variations in kernel types, with LM13 BC₂F₁ exhibiting a balanced distribution of semi-dent (7), semi-flint (13), and flint kernels (15), while LM14 BC₂F₁ showed a significant preference for semi-dent kernels (14), with fewer semi-flint (7) and flint kernels (8). This divergence suggested genetic or environmental factors influencing the dominance of certain kernel types [17].

The shape of the kernel surface varied significantly between populations. LM14 BC₂F₁ was dominated by indented kernels (3), whereas LM13 BC₂F₁ had none, highlighting the impact of specific genetic or environmental influences. Both populations predominantly had rounded kernels, with LM13 BC₂F₁ having 13 and LM14 BC₂F₁ having 22, indicating a common genetic basis. The absence of strongly pointed kernels in LM14 BC₂F₁, compared to their presence in LM13 BC₂F₁ having 7, suggests possible genetic divergence outcomes favouring different kernel shapes in these populations [17].

Endosperm colour analysis indicated a strong preference for yellow in both populations, with LM13 BC₂F₁ having 15 yellow kernels and LM14 BC₂F₁ having 16 yellow kernels. This suggested that yellow endosperm may have been selected for its nutritional and aesthetic benefits [19]. In contrast, LM14 BC₂F₁ showed a higher frequency of orange endosperm (5 kernels) compared to LM13 BC₂F₁ (1 kernel), indicating genetic diversity. The lower occurrence of white endosperm (4 in LM13 BC₂F₁ and 2 in LM14 BC₂F₁) and cream endosperm (5 in LM13 BC₂F₁ and 1 in LM14 BC₂F₁) across both populations suggested that these traits were less desirable.

Kernel colour (Top of grain) data reinforced these findings, with both populations predominantly showing yellow kernels (23 in LM13 BC₂F₁, 21 in LM14 BC₂F₁). The slight variation in white kernel frequencies between the populations (5 in LM13 BC₂F₁, 7 in LM14 BC₂F₁) suggested residual genetic variation [16].

These findings emphasized the importance of understanding the genetic basis and environmental interactions influencing these traits, which could guide future breeding and selection strategies for improved *Zea mays* derived varieties [20]. Further research was needed to analyse the genetic mechanisms and environmental factors that contribute to trait variation and provide valuable

insights for optimizing breeding programs to develop varieties tailored to specific environments and consumer preferences [21]. All of these results are shown schematically in Fig. 2.

3.2 Quantitative Trait Analysis

The quantitative analysis of kernel dimensions in *Zea mays* introgression individuals elucidated key differences and similarities between two populations, LM13 BC₂F₁ and LM14 BC₂F₁. The LM13 BC₂F₁ population exhibited an average kernel breadth of around 5.0 units, slightly higher than LM14 BC₂F₁ having 4.5 units. Both populations have similar kernel lengths, averaging 7.5 mm units, suggesting stable genetic factors for this trait. The kernel length to breadth (K L/B) ratio varies from 1.0 to 1.8 in LM13 BC₂F₁ and from 1.0 to 2.0 in LM14 BC₂F₁, indicating that kernels in LM14 BC₂F₁ are slightly more elongated. These findings highlighted significant genetic diversity and suggest that both genetic and environmental factors influence kernel dimensions [17]. Understanding these traits can inform breeding programs aimed at optimizing kernel size and shape for improved yield and adaptability in *Zea mays* varieties [16].

Focusing on the 100 seed weight of two populations: LM13 BC₂F₁ and LM14 BC₂F₁. The data is presented in Fig. 3, showing the variation in 100 seed weight across 35 samples for each population. The 100 seed weight for LM13 BC₂F₁ and LM14 BC₂F₁ populations shows significant variability. The weight ranges from approximately 13.00 grams to nearly 30.00 grams across the samples. The variation in 100 seed weight between the two populations suggests a considerable genetic diversity introduced through the introgression process. The broad range of seed weights indicates that both populations possess a wide array of genetic backgrounds, which is beneficial for breeding programs aiming to enhance specific traits. In both populations, there are instances of higher and lower seed weights, reflecting the influence of both genetic and environmental factors. The presence of heavier seeds in both populations can be advantageous for traits associated with yield and vigour, whereas the lighter seeds might indicate less favourable conditions or genetic combinations that need to be considered in breeding strategies [22]. The quantitative analysis of the 100 seed weight in *Zea mays* introgression individuals reveals

significant genetic diversity and variability within the LM13 BC₂F₁ and LM14 BC₂F₁ populations. This diversity is crucial for breeding programs as it provides a broad genetic base to select for

desirable traits such as seed weight, which is directly related to crop yield and quality [23]. The results are shown graphically in Fig. 3.




Fig. 2. Kernel Trait Distribution in LM13 BC₂F₁ and LM14 BC₂F₁




Fig. 3. Kernel dimensions and 100 Seed Weight (g) Distribution in LM13 BC₂F₁ and LM14 BC₂F₁

3.3 Genetic Variability Analysis

The analysis of genetic variability parameters for kernel traits in *Zea nicaraguensis* introgression individuals revealed significant genetic variability between the LM13 BC₂F₁ and LM14 BC₂F₁ populations. For 100 seed weight, LM13 BC₂F₁ shows a remarkably high genotypic coefficient of variation (GCV) of 37.80, indicating substantial genetic influence, whereas LM14 BC₂F₁ has a GCV of 29.69. The phenotypic coefficient of variation (PCV) is higher in LM13 BC₂F₁ (24.82) than in LM14 BC₂F₁ (12.23), suggesting greater environmental impact in the former. Kernel length exhibits notable GCV values of 12.80 (LM13

BC₂F₁) and 12.27 (LM14 BC₂F₁), with corresponding PCV values of 22.10 and 23.40, highlighting considerable genetic diversity. Additionally, kernel breadth shows high PCV values in both populations, emphasizing environmental influence. These findings indicate a broad genetic base and significant environmental interaction, essential for breeding programs targeting enhanced yield and adaptability [24]. The considerable genetic variability, particularly in 100 seed weight and kernel dimensions, underscores the potential for selecting superior genotypes, thereby aiding in the development of improved *Zea nicaraguensis* varieties [25-27].




Fig. 4. GCV and PCV Values of Introgression Individuals

Table 2. Heritability and Genetic Advance Values of Introgression Individuals

Traits	LM13 BC2F1		LM14 BC2F1	
	Broad Sense Heritability (H ²)	Genetic Advance (GA)	Broad Sense Heritability (H ²)	Genetic Advance (GA)
Kernel type	0.023	3.464	0.032	3.464
Shape of upper surface of kernel	0.023	10.387	0.067	10.610
Endosperm colour	0.115	6.270	0.072	7.179
Kernel colour	0.314	6.270	0.151	7.179
Kernel Breadth	0.027	5.797	0.021	4.805
Kernal Length	0.016	10.909	0.015	9.461
K L/B Ratio	0.064	15.134	0.060	14.482
100 Seed weight	0.013	2.068	0.051	2.440

Heritability and Genetic advance: The analysis of broad-sense heritability (H^2) and genetic advance (GA) for kernel traits in *Zea nicaraguensis* introgression individuals revealed notable findings between the LM13 BC₂F₁ and LM14 BC₂F₁ populations, results are depicted in the Table 2. Kernel type exhibited low heritability (0.023 for LM13 BC₂F₁ and 0.032 for LM14 BC₂F₁) with moderate GA, indicating limited genetic control. The shape of the upper surface of the kernel showed higher heritability in LM14 BC₂F₁ (0.067) compared to LM13 BC₂F₁ (0.023), suggesting better genetic predictability in LM14 BC₂F₁. Endosperm colour had moderate heritability (0.115 for LM13 BC₂F₁ and 0.072 for LM14 BC₂F₁) with substantial GA, indicating moderate genetic influence and significant improvement potential through selection. Kernel colour displayed the highest heritability (0.314 for LM13 BC₂F₁ and 0.151 for LM14 BC₂F₁) with considerable GA, suggesting strong genetic control and substantial selection potential. Kernel breadth and length showed low heritability and moderate GA, highlighting significant environmental influence. The K L/B ratio had moderate heritability (0.064 for LM13 BC₂F₁ and 0.060 for LM14 BC₂F₁) with high GA, indicating considerable genetic variability and improvement potential. The 100 seed weight exhibited low heritability (0.013 for LM13 BC₂F₁ and 0.051 for LM14 BC₂F₁) and low GA, suggesting limited genetic control and higher environmental impact. Overall, kernel colour and K L/B ratio emerged as significant traits with strong genetic control and high improvement potential. These findings

emphasized the importance of both genetic and environmental factors in trait expression, guiding breeding programs to focus on traits with high heritability and genetic advance for developing superior *Zea nicaraguensis* varieties [17].

3.4 Principal Component Analysis Results

PCA of LM13 BC₁F₂ Population: The PCA biplot represents the relationships between various maize kernel traits and the individual samples. Dim1 explains 28.9% of the variance and Dim2 explains 18.1%, together capturing 47% of the total variance.

Traits such as Kernel Length, Kernel Type, HSW (Hundred Seed Weight), and K L/B ratio are positively correlated with Dim1, while Kernel Breadth shows a negative correlation. Endosperm colour is negatively correlated with Dim2, whereas the Shape of upper surface of kernel is positively correlated. Samples on the right side of the biplot (e.g., samples 8, 6, 10, 21, 23, 30) have higher values for traits like Kernel Length, Kernel Type, HSW, and K L/B ratio. Samples on the left (e.g., samples 22, 27) have higher Kernel Breadth, and those at the bottom (e.g., samples 25, 28, 26) show intense Endosperm colour. Samples near the center exhibit average trait values. This analysis visualizes the trait distribution and patterns among the maize samples, aiding in the identification of trait relationships and sample groupings.

Table 3. Principal component analysis results of LM13 BC₂F₁

Component	Eigenvalue	Percentage of Variance
Comp 1	2.308	28.854
Comp 2	1.447	18.086
Comp 3	1.338	16.719
Comp 4	1.085	13.561
Comp 5	0.658	8.231
Comp 6	0.630	7.878
Comp 7	0.515	6.440
Comp 8	0.018	0.230

Table 4. Principal Component Analysis Results of LM14 BC₂F₁

Component	Eigenvalue	Percentage of Variance
Comp 1	2.516	31.452
Comp 2	1.621	20.268
Comp 3	1.225	15.317
Comp 4	0.951	11.89
Comp 5	0.627	7.843
Comp 6	0.548	6.855
Comp 7	0.506	6.325
Comp 8	0.003	0.045




Fig. 5. PCA Biplot for LM13 BC₁F₂ individuals




Fig. 6. PCA Biplot for LM14 BC₁F₂ individuals

PCA of LM14 BC₁F₂ Population: The PCA biplot you've provided illustrates the relationships between various maize kernel traits and individual samples, with Dim1 explaining 31.5% of the variance and Dim2 explaining 20.3%,

together accounting for 51.8% of the total variance.

Traits of kernel length, kernel type, HSW (hundred seed weight), and K L/B ratio have

positive loadings on Dim1, indicating that samples positioned to the right of the biplot (e.g., samples 1, 15, 19, 21, 31) are associated with higher values for these traits. Kernel breadth is negatively correlated with Dim1, suggesting that samples (e.g., samples 3, 4, 11, 13) exhibited higher kernel breadth. Dim2 is positively correlated with traits like kernel length and K L/B ratio, while endosperm colour had been negative loading, indicating that samples located at the bottom (e.g., samples 5, 18, 25, 27, 30) tend to have intense endosperm colour and specific kernel colour characteristics. The central grouping of samples suggested average values for most traits. This biplot effectively visualizes the distribution and patterns of traits among the maize samples, aiding in understanding their relationships and identifying trait groupings [14].

4. CONCLUSION AND FUTURE PROSPECTS

This study comprehensively analysed kernel traits of BC₂F₁ maize lines obtained from crosses between the inbred lines LM13 and LM14 with *Zea mays*. It is an inference that genetic variability is relatively high within the traits concerning the type, shape, colour, dimensions, and weight of kernels. Genetic diversity, the main effect, and interaction of environmental factors were found to be responsible for shaping all of these traits, as described both descriptively and statistically. The traits concerned with kernel colour and kernel length-to-breadth ratio showed high heritability, along with high genetic advance in most cases, indicating high potential under effective selection programs.

These critical characteristics causing the most variability within the BC₂F₁ populations were further emphasized by the results from the PCA. With this knowledge of critical traits, selection efforts can be put forth by breeders with the most influence on yield, quality, and marketability. The estimated heritability and genetic variability values obtained in the present study indicate much potential for improvement in maize with well-informed breeding strategies. Indeed, introgressions of gene material related to wild species, such as *Zea mays*, proved beneficial to maize by contributing novel traits that could positively influence improvement for new agronomic performance and adaptability [17].

Future work should focus on revealing the genetic mechanisms underlying these kernel characteristics to facilitate the estimation of gene action. Investigations on the environmental factors responsible for the expression of these traits will further guide the breeding programs in such a way that suitable genetic materials are developed for various agro-ecological zones. This can be aided with the molecular methods like MAS and genomic selection. The genes underlying the expression of traits can be effectively mapped using QTL analysis. Also, broadening the study to incorporate more advanced backcross generations and multi-location trials helps in affirming stability and adaptability in the chosen traits. This is ultimate research that would pave the way for developing superior maize varieties to meet the demands in achieving food security, nutrition, and industrial applications to ensure sustainable maize production.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Murdia LK, Wadhvani R, Wadhawan N, Bajpai P, Shekhawat S. Maize utilization in India: An overview. American Journal of Food and Nutrition. 2016;4(6): 169-176.
2. Klopfenstein TJ, Erickson GE, Berger LL. Maize is a critically important source of food, feed, energy and forage in the USA. Field Crops Research. 2013;153:5-11.
3. Shiferaw B, Prasanna BM, Hellin J, Bänziger M. Crops that feed the world 6. Past successes and future challenges to the role played by maize in global food security. Food Security. 2011;3:307-327.
4. Almeida, Rafael Nunes de, André Soares de Castro. Potential of grain physical traits to the study of variability in maize. Journal

- of Scientific Research and Reports. 2022; 28(11):153-60.
Available:<https://doi.org/10.9734/jsrr/2022/v28i111712>.
5. Russell WA. Genetic improvement of maize yields. *Advances in Agronomy*. 1991, Jan 1;46:245-98.
 6. Kist V, Bernardi Ogliairi J, de Miranda Filho JB, Alves AC. Genetic potential of a maize population from Southern Brazil for the modified convergent–divergent selection scheme. *Euphytica*. 2010, Nov;176:25-36.
 7. Xu Y, Skinner DJ, Wu H, Palacios-Rojas N, Araus JL, Yan J, Crouch JH. Advances in maize genomics and their value for enhancing genetic gains from breeding. *International Journal of Plant Genomics*. 2009;2009(1):957602.
 8. Hassan ZM, Sebola NA, Mabelebele M. The nutritional use of millet grain for food and feed: A review. *Agriculture & Food Security*. 2021;10:1-14.
 9. Liu Z, Garcia A, McMullen MD, Flint-Garcia SA. Genetic analysis of kernel traits in maize-teosinte introgression populations. *G3: Genes, Genomes, Genetics*. 2016; 6(8):2523-2530.
 10. Nawaz R, Riaz U, Gouhar H, Mukhtar M, Arshad A, Hussain A, Amjad I. Harnessing genetic diversity for sustainable maize production. *Journal of Physical, Biomedical and Biological Sciences*. 2023;2023(1):15-15.
 11. Mano Y, Omori F. Flooding tolerance in interspecific introgression lines containing chromosome segments from teosinte (*Zea nicaraguensis*) in maize (*Zea mays* subsp. *mays*). *Annals of Botany*. 2013;112(6): 1125-1139.
 12. Varalakshmi S, Sahoo S, Singh NK, Pareek N, Garkoti P, Senthilkumar V, Nankar AN. Marker–trait association for protein content among maize wild accessions and coix using SSR markers. *Agronomy*. 2023;13(8):2138.
 13. Sahoo S, Adhikari S, Joshi A, Singh NK. Use of wild progenitor teosinte in maize (*Zea mays* subsp. *mays*) improvement: present status and future prospects. *Tropical Plant Biology*. 2021;14(2):156-179.
 14. Okporie EO. Characterization of maize (*Zea mays* L.) germplasm with principal component analysis. *Agro-Science*. 2008; 7(1):66-71.
 15. Jilo T, Tulu L, Birhan T, Beksisa L. Genetic variability, heritability and genetic advance of maize (*Zea mays* L.) inbred lines for yield and yield related traits in southwestern Ethiopia. *Journal of Plant Breeding and Crop Science*. 2018;10(10): 281-289.
 16. Jaradat AA, Goldstein W. Diversity of maize kernels from a breeding program for protein quality: I. Physical, biochemical, nutrient, and color traits. *Crop Science*. 2013;53(3):956-976.
 17. Prasanna BM. Diversity in global maize germplasm: Characterization and utilization. *Journal of Biosciences*. 2012; 37(5):843-855.
 18. Araus JL, Slafer GA, Royo C, Serret MD. Breeding for yield potential and stress adaptation in cereals. *Critical Reviews in Plant Science*. 2008;27(6):377-412.
 19. Kajla P, Chaudhary V, Dewan A, Goyal N. Colored cereals: Botanical aspects. In *Functionality and Application of Colored Cereals*. Academic Press. 2023;1-25.
 20. Lorant A. Plasticity and genetic adaptation as contributors to the evolutionary history of cultivated maize and its wild relatives (Doctoral dissertation, Université Paris Saclay (COMUE); 2018).
 21. Smale M, Bellon MR, Aguirre Gomez JA. Maize diversity, variety attributes, and farmers' choices in Southeastern Guanajuato, Mexico. *Economic Development and Cultural Change*. 2001; 50(1):201-225.
 22. Hossain F, Muthusamy V, Bhat JS, Zunjare RU, Kumar S, Prakash NR, Mehta BK. Maize breeding. In *Fundamentals of Field Crop Breeding*. Singapore: Springer Nature Singapore. 2022;221-258.
 23. Anjorin FB, Ogunniyan DJ. Comparison of growth and yield components of five quality protein maize varieties. *International Journal of Agriculture and Forestry*. 2014;4(1):1-5.
 24. Yadav OP, Hossain F, Karjagi CG, Kumar B, Zaidi PH, Jat SL, Dhillon BS. Genetic improvement of maize in India: retrospect and prospects. *Agricultural Research*. 2015;4:325-338.
 25. Bioversity International; International Maize and Wheat Improvement Center (CIMMYT) Key access and utilization descriptors for maize genetic resources. 2009;6.
 26. Shrestha J. Agro-morphological characterization of maize inbred lines. *Wudpecker Journal of Agricultural Research*. 2013;2(7):209-211.

27. Solomon Oyeniya, Omotola, Bunmi Olasanmi, Samuel Adelowo Olakojo, Solomon Tayo Akinyosoye. Genetic Diversity and responses of some selected yellow maize genotypes to stem borer (Sesamia Calamistis Hampson) infestation. Journal of Experimental Agriculture International. 2019;39(1):1-10. Available:<https://doi.org/10.9734/jeai/2019/v39i130322>.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/120100>