



Seaweed-enriched Cookies: A Nutritional and Functional Perspective

**R. B. Hajare ^a, A. U. Pagarkar ^{a*}, A. S. Desai ^a, J. M. Koli ^a,
P. E. Shingare ^a, Soorya Gopan ^a, S. B. Satam ^a
and A. N. Sawant ^a**

^a College of Fisheries, Shirgaon, Ratnagiri, Dr. B. S. Konkan Krishi Vidyapeeth, Dapoli, Maharashtra.
415629, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajb2t/2024/v10i4225>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/127013>

Review Article

**Received: 12/09/2024
Accepted: 14/11/2024
Published: 22/11/2024**

ABSTRACT

Seaweed and seaweed-based products are gaining global attention due to their health-benefiting properties. For the last three decades, seaweeds have been used as nutraceutical food, as they provide dietary benefits. It offers a rich source of nutrients, including minerals, vitamins, and fibre, while also possessing unique functional properties such as water and oil absorption. Due to their functional characteristics, the incorporation of seaweed into bakery products is becoming a novel strategy to develop nutritionally enriched value-added products. Cookies are one of the most popular bakery products, with high consumer preference. The incorporation of seaweed offers a

*Corresponding author: E-mail: aupcof@gmail.com;

promising avenue for enhancing the nutritional value of cookies. It improves the physical, sensorial, and nutritional properties of cookies. This review seeks to explore the significance of seaweed consumption, the development of value-added cookies incorporating seaweed, and the impact of seaweed on the overall quality of cookies.

Keywords: Seaweed; cookies; bakery products; nutritionally enriched cookies.

1. INTRODUCTION

Seaweeds are marine, multicellular eukaryotes characterized by their photosynthetic nature, chlorophyll content, and distribution spanning from the intertidal zone to depths of approximately 300 meters [1]. It represents a valuable renewable resource within the marine ecosystem [2]. This macroalgae is classified into three primary groups: Rhodophyta (red algae), Phaeophyta (brown algae), and Chlorophyta (green algae). Globally, approximately 10,300 species of seaweed have been documented, with a total annual production of 38 million tonnes as of 2022 [3,4].

The Indian coastline is characterized by a rich diversity of seaweed populations. In 2021, India's seaweed production reached 34,000 tonnes (CMFRI, 2022). The seaweed flora of India exhibits a high degree of diversity, thriving in various coastal habitats such as rocky beaches, mudflats, estuaries, coral reefs, and lagoons. Approximately 770 seaweed species have been identified along the Indian coastline, comprising 420 red algae, 184 green algae, and 166 brown algae [2,5].

The occurrence of seaweeds is recorded abundantly along the coastal regions of Tamil Nadu, Gujarat, Andhra Pradesh, Orissa, West Bengal, Kerala, Maharashtra, Lakshadweep and Andaman Nicobar Islands. Along the Maharashtra (Mumbai, Ratnagiri), Goa, Karnataka (Karwar), Kerala (Vizhinjam, Varkala), and Orissa (Chilika) rich seaweed beds are reported [6].

Traditionally seaweed has been used as food, folk remedies, dyes and fertilizers. For the last three decades, seaweeds have been used as nutraceuticals or functional foods, as they give dietary benefits [7]. Seaweeds are recognized as a significant source of nutrients and possess numerous health-promoting properties. Maybe due to their nutritional benefits, seaweeds have been regarded as a food of gods and kings [8]. At present, over 56 countries produce seaweed worldwide with 97% of the total production

comes from aquaculture. Out of the total world's seaweed production, only 30-38% of seaweed is utilized for human consumption [4].

Though the Indian coastline is an abundant source of seaweed having a potential of 9.7 million tonnes with 342 production sites (CMFRI, 2022), it has very low share in global production. The commercial exploitation of seaweeds in India began in 1966, primarily for the production of agar and alginate [8]. While phycocolloids (agar, alginate, and carrageenan) have been widely utilized in various industries, the use of seaweeds as a direct food source remains limited within the country.

Bakery products are the oldest and traditional stable diet of many countries globally with huge contribution in the food processing industry. It comprises the third largest segment of the packaged and convenience food (ready-to-eat and ready-to-cook products) market [1,9]. The bakery made products contributes 1.14% to the country's gross domestic product and engages more than 1.6 million people directly and 120 million indirectly on its supply chains in India [10].

Cookie is one of the popular bakery products and can be defined as a chemically leavened product which is baked in an oven for usually a short time span. The nutrient availability, inexpensiveness, palatability, compactness and convenience makes cookies an ideal bakery product. In comparison with other baked products like bread, cakes, etc. cookies have low moisture content which makes it free from microbial spoilage and have a longer shelf life [11].

The integration of seaweeds into bakery products has emerged as a focal point of research given their established nutritional and functional attributes. Studies have consistently demonstrated that the addition of seaweeds can substantially enhance the phytochemical content, antioxidant activity, and dietary fiber in bakery products. Moreover, seaweed incorporation can positively influence the nutritional, textural, and sensory properties of a diverse range of bakery items [12,13].

The primary objective of this study is to conduct a comprehensive review of the nutritional significance and health-promoting attributes of seaweeds, their integration into cookie formulations, and the subsequent impact of seaweed incorporation on the nutritional, physical, and sensory properties of cookies.

2. BENEFITS OF SEAWEED CONSUMPTION ON HUMAN HEALTH

Seaweed, a nutrient-dense marine plant, offers a wealth of health benefits due to its unique composition. It is a great source of vitamins (A, B1, B2, B9, B12, C, D, E, and K), essential minerals (calcium, iron, iodine, magnesium, phosphorus, potassium, zinc, copper, manganese, selenium, and fluoride), dietary fibres, protein, essential amino acids and polyphenols [14].

Rajapakse and Kim [15] reviewed the effects of seaweed consumption on human digestive health. They stated that the presence of dietary fibres in seaweed helps in reducing the risk of colorectal cancer and suppressing gastrointestinal inflammation. According to global epidemiological studies, countries where seaweeds are regularly consumed have significantly fewer instances of obesity and dietary-related disease [7].

Also, some health-promoting properties are shown by seaweeds, i.e., anticancer, antiviral, antifungal, antidiabetic, antihypertensive, immunomodulatory, anticoagulant, anti-inflammatory, antioxidant, UV protective, neuroprotective, dyslipidaemia, bone-health, heart-health and mental-health benefits [1].

3. CHALLENGES ASSOCIATED WITH SEAWEED BAKERY PRODUCT

Though incorporation of seaweed into bakery products can be a novel and health-beneficial approach in food industry, some challenges are associated with the production and sale of the bakery products. These challenges are mentioned below:

Availability of seaweed: Seaweed is not as widely available as other ingredients, which can make it difficult to find and obtain.

Strong flavour: Seaweed has a strong flavor that can be overpowering for some consumers. This can be a challenge when creating seaweed

bakery products that are palatable to a wide range of people.

Regulatory challenges: There may be regulatory challenges associated with the production and sale of seaweed bakery products, such as food safety regulations and labeling requirements.

4. INCORPORATION OF SEAWEEDS IN BAKERY PRODUCTS

Seaweeds have gained attention for their potential use in bakery products due to their nutritional and functional properties. Studies have shown that incorporating seaweed powder, typically from species like *Kappaphycus alvarezii*, *Ulva lactuca*, and *Gracilaria corticata*, can enhance the nutritional profile of bakery products. Seaweeds increase dietary fibre, protein, polyphenols, and antioxidant capacity, making products like bread, cakes, cookies, and buns more nutritionally robust.

For instance, Mamat et al. [16,17] demonstrated that incorporating red seaweed powder improves water absorption in dough and increases moisture and fibre content in buns, though higher seaweed levels may decrease protein content. Sensory evaluations suggest that products with moderate seaweed incorporation are more acceptable, as higher concentrations may negatively impact flavour. Turuk and Banerjee (2023) found that bakery products containing *G. corticata* seaweed tend to have better nutritional outcomes than those with *U. 126 ornema*, further supporting the idea that seaweed-enriched bakery items are healthier alternatives to traditional ones. Overall, seaweeds offer a promising avenue for creating nutritionally enhanced bakery products with added health benefits.

Quitral et al. [18] emphasized that finely ground seaweed can form stable emulsions in dough, enhancing the functional qualities of bakery products. However, the optimal incorporation levels are critical, as excessive seaweed can impair sensory qualities, particularly flavour. Commonly recommended levels are around 4–10% for various bakery products.

5. SEAWEED ENRICHED COOKIES

The incorporation of seaweed into cookies has been explored as a means of enhancing their nutritional, functional, and sensory properties. Various types of seaweed, including *Caulerpa racemosa*, *Bifurcaria bifurcata*, *Ulva*

lactuca respectively, and others, have shown potential to improve the quality of baked products by increasing protein, fibre, and antioxidant content, while also affecting dough and texture properties. Numerous studies have demonstrated the effectiveness of different seaweed species in cookie formulations, contributing to the development of healthier and more functional food products.

5.1 Incorporation of Seaweed in Cookies

- **Sea Grapes (*Caulerpa racemosa*):** Kumar et al. [1] prepared semi-sweet biscuits supplemented with seaweed (*Caulerpa racemosa*) powder under the institutional project of ICAR-CIFT to assess its impact on physical, functional, and antioxidant properties. The inclusion of *C. racemosa* significantly increased water and oil absorption capacities, as well as solvent retention capacities of the flour mix. Protein, fiber, phenolic content, and antioxidant activities were increased with the increasing level of seaweed powder. Hence, the incorporation of seaweed powder showed a positive effect on the overall quality of the cookies.
- ***Bifurcaria bifurcate*:** Arufe et al. [19] incorporated *Bifurcaria sornemannii* seaweed powder into chestnut flour dough at 3%, 6%, and 9%. The seaweed addition enhanced dough stability and reduced its mechanical weakening. Gelatinized starch, used during mixing, improved dough recovery properties.
- ***Ulva 127ornema*:** Jenifer and Kanjana [20] used *Ulva 127ornema* in seaweed-based biscuits to improve the anthropometric profile of malnourished children. Various formulations of *Ulva 127 ornema* were tested, and the biscuits were supplemented to malnourished children for 2 months, resulting in improved nutritional status.
- **Korean Coastal Seaweeds:** Oh et al. [21] studied the impact of seaweeds native to the Korean coast on cookies. They prepared seaweed flour using four different species, including *Sargassum fulvellum*, *Enteromorpha linza*, *Codium fragile* and *Hizikia fusiforme*. In the preparation of cookies, 5% wheat flour was replaced with seaweed flour. The study found that seaweed incorporation improved solvent retention capacities, and cookies with *Hizikia fusiforme* were highly preferred in sensory evaluations.

- ***Gelidium spinosum*:** Poulouse et al. [22] used *Gelidium spinosum* seaweed to extract stigmasterol, which was nano-emulsified and incorporated into biscuit dough. The resulting biscuits had higher nutritional content, reduced hardness, and were sensorily acceptable.
- ***Ulva intestinalis*:** Mohibullah et al. [23] prepared seaweed cookies with the inclusion of different percentages of powdered and fragmented forms of *Ulva intestinalis*. They found that all the studied parameters of cookies including moisture content, spread factor, baking loss, pH, cookie density, colour, texture properties and volatile basic nitrogen were found within acceptable limits. They mentioned that cookies prepared with the inclusion of 2.5% powdered and 5% fragmented forms of seaweed can be considered healthy nutritional diets due to their high sensorial and nutritional properties.
- ***Portieria hornemannii*:** Sivaraman et al. [24] combined *Portieria hornemannii* seaweed with Tilapia fish protein concentrate (FPC) in cookie formulations. Cookies with 2.5% seaweed and 5% FPC showed high overall acceptability, enhanced nutritional value, and sensory attributes.

6. EFFECT OF SEaweEDS ON PHYSICAL, SENSORIAL, AND NUTRITIONAL QUALITY OF COOKIES

6.1 Sensorial Properties

Results of the sensory evaluation of seaweed cookies vary based on the species and concentration of incorporated seaweed.

Kumar et al. [1] substituted refined wheat flour with seaweed powder at different levels, including 0, 1, 5, and 10%. A significant decrease in sensory attributes was observed at higher levels (5 and 10%) of seaweed incorporation as compared to 1% of seaweed powder-supplemented biscuits.

Similarly, Mohibullah et al. [23] prepared seaweed cookies with the inclusion of 1, 2.5, 5 and 10% of powdered and fragmented forms of *Ulva intestinalis*. Seaweed inclusion levels of 1-5% were acceptable to sensory panellists, with 2.5% powdered and 5% fragmented forms of seaweed concentrations selected for further studies due to their favourable properties.

Table 1. Summary of work conducted on seaweed incorporated cookies and their characteristics

A. Incorporation of different levels seaweed in cookies by different authors, used baking temperature and time					
Authors	Kumar et al. [1]	Arufe et al. [19]	Oh et al. [21]	Poulose et al. [22]	Mohibullah et al. [23]
Species of seaweed	<i>Caulerpa racemosa</i>	<i>Bifurcaria bifurcate</i>	<i>Hizikia fusiforme</i>	<i>Gelidium spinosum</i>	<i>Ulva intestinalis</i>
Level of seaweed incorporation	0,1, 5 and 10%	3, 6 and 9%	5%	Extract of seaweed: Stigmasterol was incorporated	2.5% powdered and 5% fragmented form
Baking temperature (°C)	190	180	160	160	180
Baking time (minutes)	14-15	20	10	20 ± 5	12
B. Effect of seaweed incorporation on proximate composition of cookies when compared with control cookies used					
Moisture (%)	Not significantly changed (from 4.19 to 4.03)	--	--	Decreased	Decreased (from 3.57 to 3.02)
Protein (%)	Increased (from 7.69 to 9.01)	--	--	Increased	Increased (from 6.88 to 11.78)
Fat (%)	Not significantly changed (from 14.48 to 13.77)	--	--	Increased	Not significantly changed (from 25.39 to 25.04)
Ash (%)	Increased (from 1.28 to 2.23)	--	--	--	Increased (from 1.45 to 4.81)
Carbohydrate (%)	Decreased (from 71.7 to 68.75%)	--	--	Decreased	--
C. Effect of seaweed incorporation on physical parameters of cookies when compared with control cookies used					
Hardness (N)	Decreased (from 17.57 to 11.29)	Increased (from 0.82 to 1.01)	--	Decreased (from 12.80 to 8.87)	Not significantly changed (from 2.3 to 2.82 kg/cm ²)
Colour	L* (lightness/darkness)	Decreased (from 66.6 to 41.1)	--	Decreased (from 70.69 to 45.18)	Decreased (44.68)
	a* (redness/greenness)	Decreased (from 10.6 to 3.26)	--	Increased (from -3.73 to -0.08)	Not significantly changed
	b* (yellowness/blueness)	Decreased (from 32.0 to 22.9)	--	Decreased (from 20.85 to 11.76)	Increased
Water absorption capacity	Increased (from 1.24 to 1.52)	Increased (in dough)	Increased (in dough)	--	--
Oil absorption capacity	Increased (from 0.87 to 1.11)	--	--	--	--

Solvent retention capacity	Lactic acid	Increased (from 1.52 to 1.56)	--	Decreased (from 110.3 to 91.20%)	--	--
	Na ₂ CO ₃	Increased (from 1.37 to 1.59)	--	Decreased (from 80.62 to 78.11%)	--	--
	Sucrose	Increased (from 1.98 to 2.38)	--	Increased (from 122.4 to 124.7%)	--	--

Oh et al. [21] conducted the sensory evaluation of cookies prepared by replacing 5% wheat flour with seaweed flour made up of four different seaweed species viz., *Sargassum fulvellum*, *Enteromorpha linza*, *Codium fragile* and *Hizikia fusiforme*. In sensory evaluation they found that, cookies with the incorporation of 5% seaweed (*Hizikia fusiforme*) powder by replacing wheat flour were greatly preferred with the highest sensory scores.

Effect of seaweed (*Portieria hornemannii*) powder and tilapia fish protein concentrate (FPC) on cookies were studied by Sivaraman et al. [24]. Cookies were prepared using 5% FPC and different percentages of seaweed viz., 2.5%, 5%, 7.5% and 10%. Sensory evaluation showed that cookies prepared with 2.5% seaweed and 5% FPC obtained the highest overall acceptability.

Overall, cookies with lower seaweed content tend to score higher in sensory evaluation. The incorporation of high concentrations of seaweed into cookies can lead to a decrease in sensory scores due to the introduction of strong, characteristic marine flavours and textures. Seaweed's unique sensory attributes, including astringency and bitterness, can negatively impact the overall acceptability of the final product [25]. Quitral et al. [18] mentioned that the incorporation of seaweed in cookies should be less than 5%.

6.2 Physical Properties

Seaweed incorporation generally enhances the water and oil absorption capacities of doughs [1], increases dough stability [19], and improves baking properties. For example, gelatinized starch mixed with seaweed improves dough recovery properties, and seaweed can increase the spread factor and reduce the baking loss [24]. Mohibullah et al. [23] found that cookies with *Ulva intestinalis* had acceptable physicochemical characteristics, including moisture content, spread factor, texture, and baking loss. Arufe et al. [19] noted that the hardness of cookies increased with more than 3% seaweed.

6.3 Nutritional Quality

Seaweed-enhanced cookies tend to have improved nutritional profiles, including higher protein, fiber, phenolic content, and antioxidant activities [1,19]. The addition of seaweeds also increases dietary fiber [24] and essential

nutrients such as ash and lipid content [23]. Seaweed-based biscuits supplemented to malnourished children showed significant improvements in anthropometric measures [20]. Additionally, Poulouse et al. [22] demonstrated that seaweed biscuits containing stigmasterol had anti-diabetic properties, making them a promising functional food for diabetic patients.

7. CONCLUSION

Seaweed is a nutrient-dense food source that exhibits several health-promoting properties. The incorporation of seaweed in cookies is an innovative approach for developing a nutritionally enhanced, value-added product. The inclusion of seaweed significantly enhances the nutritional, functional, and sensory properties of cookies. Seaweed can be incorporated into cookies in various forms, including fragmented, powdered, and extracted. Cookies with higher seaweed concentrations may exhibit lower sensory scores due to the introduction of strong marine flavours and textures. This suggests that a moderate level of seaweed incorporation, typically less than 5%, is optimal for maintaining sensory acceptability. Though the higher level of seaweed inclusion negatively affects the sensory properties, the incorporation of a proper amount of seaweed in cookies can enhance dough properties, improve baking outcomes, and significantly elevate the nutritional profile of cookies. These findings highlight the potential of seaweed as a functional ingredient for developing healthier and more nutritious baked goods.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

ACKNOWLEDGEMENT

The authors wish to thank respected vice-chancellor and all the university authorities, Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli for providing the facilities to carry out this study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kumar A, Krishnamoorthy E, Devi HM, Uchoi D, Tejpal CS, Ninan G, Zynudheen AA. Influence of sea grapes (*Caulerpa racemosa*) supplementation on physical, functional, and anti-oxidant properties of semi-sweet biscuits. *Journal of Applied Phycology*. 2018;30:1393-1403.
2. Rao PVS, Mantri VA. Indian seaweed resources and sustainable utilization: Scenario at the dawn of a new century. *Current sci*. 2006;91(2):164-174.
3. Chakraborty K, Gopalakrishnan A, Mohanty BP. Seaweeds as treasury to develop novel marine-derived biomolecules and nutraceuticals. *RDWU Bulletin of Science*. 2022;3(1):10-22.
4. FAO. The State of World Fisheries and Aquaculture 2024. Blue Transformation in action. Rome. 2024;264.
5. Joshi A, Desai AY, Mulye V. Seaweed resources and utilization: an overview. *Biotech Express*. 2015;2:22.
6. Rao PS, Periyasamy C, Kumar KS, Rao AS, Anantharaman P. Seaweeds: distribution, production and uses. *Bioprospecting of algae*. Society for Plant Research. 2018;59-78.
7. Shannon E, Abu-Ghannam N. Seaweeds as nutraceuticals for health and nutrition. *Phycologia*. 2019;58(5):563-577.
8. Escrig JA, Goni Combrodon I. Nutritional evaluation and physiological effects of edible seaweeds. *Arch Latinoam Nutr*. 1999;49(2):114-120.
9. Kulshreshtha SK, Kumar A, Passah DRS. Indian Experience in Ensuring Sustainability in Baking Industry. In *Baking Business Sustainability Through Life Cycle Management*. 2023;147-167.
10. Govt. of India. Ministry of Micro, Small and Medium Enterprises - Annual Report. 2017-2018. Available:<https://msme.gov.in/sites/default/files/MSME-AR-2017-18-Eng.pdf>.
11. Dinnah A, Guyih MD, Eke MO. Vitamin Content and Storage Studies of Cookies Produced from Wheat, Almond and Carrot Flour Blends. *Asian Food Science Journal*. 2020;15(3):31-41.
12. Cox S, Abu-Ghannam N. Incorporation of Himanthalia Elongata Seaweed to Enhance the Phytochemical Content of Breadsticks Using Response Surface Methodology (RSM). *International Food Research Journal*. 2013;20:1537-1545.
13. Roohinejad S, Koubaa M, Barba FJ, Sajjoughian S, Amid M, Greiner R. Application of seaweeds to develop new food products with enhanced shelf-life, quality and health-related beneficial properties. *Food research international*. 2017;99 Pt 3:1066-1083.
14. Lomartire S, Marques JC, Gonçalves AM. An overview to the health benefits of seaweeds consumption. *Marine Drugs*. 2021;19(6):341.
15. Rajapakse N, Kim SK. Nutritional and digestive health benefits of seaweed. *Advances in food and nutrition research*. 2011;64:17-28.
16. Mamat H, Matanjun P, Hamid M, Yeoh WC. The effect of seaweed on the quality of bakery product. In *Emeritus Prof. Dr. Faizah Binti Mohd Sharoum Chairman Umt International Annual Symposium on Sustainability Science and Management*. 2016;42:745.
17. Mamat H, Matanjun P, Ibrahim S, Md. Amin SF, Abdul Hamid M, Rameli AS. The effect of seaweed composite flour on the textural properties of dough and bread. *Journal of Applied Phycology*. 2014; 26:1057-1062.
18. Quitral V, Sepúlveda S, Gamero-Vega G, Jimenez P. Seaweeds in bakery and farinaceous foods: A mini-review. *International Journal of Gastronomy and Food Science*. 2022;28: 100403.
19. Arufe S, Chenlo F, Sineiro J, Moreira R. Effect of brown seaweed addition and starch gelatinization on gluten-free chestnut flour doughs and cookies. *Journal of Food Measurement and Characterization*. 2019;13(4):2571-2580.
20. Jenifer A, Kanjana K. Effect of seaweed based biscuit supplementation on anthropometric profile of malnourished children residing at Tuticorin. *J Sci Technol*. 2019;4:2349-5456.
21. Oh H, Lee P, Kim SY, Kim YS. Preparation of cookies with various native seaweeds found on the Korean coast. *Journal of Aquatic Food Product Technology*. 2020; 29(2):167-174.
22. Poulouse N, Sajayan A, Ravindran A, Chandran A, Priyadarshini GB, Selvin J, Kiran GS. Anti-diabetic potential of a stigmasterol from the seaweed *Gelidium spinosum* and its application in the formulation of nanoemulsion conjugate for the development of functional

- biscuits. *Frontiers in nutrition*. 2021;8:694362.
23. Mohibbullah M, Amin A, Talha MA, Baten MA, Rana MM, Sabuz AA, Newaz AW, Choi JS. Physicochemical and nutritional characteristics of cookies prepared with untapped seaweed *Ulva intestinalis*: An approach to value addition as a functional food. *Foods*. 2023;12(1):205.
24. Sivaraman L, Deshmukhe G, Xavier KAM, Hirekudel SK, Desai AS, Balange AK. Effect of incorporation of seaweed (*Portieria hornemanni*) and Tilapia fish protein concentrate on the nutritional, physical and sensory attributes of cookies. *International Journal of Food Science and Technology*. 2023;58(8):4415-4424.
25. Cassani L, Gomez-Zavaglia A, Jimenez-Lopez C, Lourenço-Lopes C, Prieto MA, Simal-Gandara J. Seaweed-based natural ingredients: Stability of phlorotannins during extraction, storage, passage through the gastrointestinal tract and potential incorporation into functional foods. *Food Res Int*. 2020;137:109676. Available:<https://doi.org/10.1016/j.foodres.2020.109676>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/127013>